THE COBORDISM OF INVOLUTIONS ON ORIENTABLE MANIFOLDS

R. J. ROWLETT

ABSTRACT. This note calculates the cobordism of smooth, oriented manifolds with orientation-reversing involution, and also the 2-torsion in the cobordism of such manifolds with orientation-preserving involution. In particular, it is shown that the forgetful homomorphism maps both groups monomorphically into the cobordism of unoriented manifolds with involution.

1. Statement of results. Let $\mathcal{O}_*$ be the cobordism theory of smooth oriented manifolds M with involution T ; we do not require T to be orientation preserving. Then $\mathcal{O}_* = \mathcal{O}_*^- \oplus \mathcal{O}_*^+$, where the equivalence class $[M, T]$ is in $\mathcal{O}_*^-$ (respectively, $\mathcal{O}_*^+$), if T is orientation reversing (respectively, preserving). The notation agrees with that of [4].

In this note we determine $\mathcal{O}_*^- \oplus \text{torsion } \mathcal{O}_*^+$. Rosenzweig [5] proved that the torsion of $\mathcal{O}_*^+$ is all of order 2. In $\mathcal{O}_*^-$ every class is of order 2, for if T is orientation reversing,

$$(M \cup M, T \cup T) \cong \partial(M \times I, T \times \text{Id})$$

under the diffeomorphism $\text{Id} \cup T$.

$\mathcal{O}_*^+$ was also studied by Conner [2], and $\mathcal{O}_*$ has recently been studied by Lee and Wasserman [4], who show how classes of $\mathcal{O}_*$ are detected by equivariant characteristic numbers. Our results complement these two papers, and we offer a short proof (Lemma 4) of a result which appears in both.

THEOREM 1. *Let I_* be the cobordism of unoriented manifolds with involution [3, Chapter IV]. The forgetful map $\rho: \mathcal{O}_* \rightarrow I_*$ is monic on 2-torsion.*

From [6] it follows that the torsion is mapped monomorphically into the Wall bordism group $\tilde{W}_*(\bigvee_{j=1}^{\infty} TBO(j))$ (see [7, Chapter VIII]). An element of this group is represented by a vector bundle $\nu \rightarrow F$, whose disk bundle $D\nu$ is a Wall manifold [8], together with a classifying map

Received by the editors November 14, 1972.

AMS (MOS) subject classifications (1970). Primary 57D85; Secondary 57D85.

Key words and phrases. Equivariant cobordism, orientation-preserving involution, orientation-reversing involution, Wall manifold.

© American Mathematical Society 1973

$(D\nu, S\nu) \rightarrow (\bigvee TBO(j), \infty)$. Define $\partial'[D\nu] = [Di^* \nu]$, where $i: N \subset F$ includes a submanifold dual to $w_1 F + w_1 \nu$. Then $Di^* \nu$ is dual to $w_1 D\nu$, hence is oriented.

THEOREM 2. $\mathcal{O}_*^- \cong \text{Im } \partial' \cap \tilde{W}_*(\bigvee_{j=1}^\infty TBO(2j+1))$.

Let $p: \tilde{W}_*(\bigvee TBO(2j)) \rightarrow W_*$ map $[D\nu]$ to the class of the projective space bundle $P\nu$, and $q: \tilde{W}_*(TBO(1)) \rightarrow \tilde{W}_*(\bigvee TBO(2j))$ map $[D\nu]$ to $[D(\nu \oplus 1)]$. Then $pq=0$ and $\partial'(\text{Im } q) \subseteq \text{Im } q$, so p and ∂' induce

$$p': \tilde{W}_*(\bigvee TBO(2j))/\text{Im } q \rightarrow W_*$$

and

$$\partial'': \tilde{W}_*(\bigvee TBO(2j))/\text{Im } q \rightarrow \tilde{W}_*(\bigvee TBO(2j))/\text{Im } q.$$

THEOREM 3. torsion $\mathcal{O}_*^+ \cong \text{Im } \partial'' \cap \text{Ker } p'$.

These theorems are used to describe Z_2 -generators of $\mathcal{O}_*^-$ and torsion $\mathcal{O}_*^+$. The usefulness of the latter description is limited by lack of knowledge of $\text{Ker } p'$.

I wish to thank Professors Lee and Wasserman for supplying an advance copy of [4], and also Professor R. E. Stong for his advice and encouragement.

2. Wall manifolds with involution. A *Wall manifold with involution*, (W, T, f) , is an unoriented manifold W with an involution T and a map $f: W \rightarrow S^1$ such that $fT=f$ and $f^*(\iota)=w_1 W$, where $\iota \in H^1(S^1; Z_2)$ is the generator. To construct examples one uses:

LEMMA 1 (See [6, (2.5)]). *If $D\nu$ is a Wall manifold (respectively, orientable) and $\dim \nu$ is even, then $P\nu$ is a Wall manifold (respectively, orientable).*

Let W_*^I and $W_*^{F'}$ be the cobordism groups of Wall manifolds with involution and with fixed point free involution, respectively. Following are the chief results of [6]. The forgetful map $W_*^I \rightarrow I_*$ is monic. Clearly $\text{Im } \rho \subseteq W_*^I$. If $d: W_*^I \rightarrow \mathcal{O}_*$ assigns to $[W, T, f]$ the class of $(N, T|N)$, where $N \subset W$ is an invariant submanifold dual to $w_1 W$, then the sequence

$$\mathcal{O}_* \xrightarrow{\rho} W_*^I \xrightarrow{d} \mathcal{O}_* \xrightarrow{2} \mathcal{O}_*$$

is exact. Define $d' = \rho d: W_*^I \rightarrow W_*^I$. The map $s: W_*^I \rightarrow \tilde{W}_*(\bigvee TBO(j))$, given by classifying the normal bundle to the fixed set, is monic. The map $\partial: \tilde{W}_*(\bigvee TBO(j)) \rightarrow W_*^{F'}$, given by $\partial[D\nu] = [S\nu, \text{antipodal map}]$, is epic.

We give a new proof of the last fact by constructing a splitting map. Recalling [6, (3.1)], if $[M, f] \in W_*(RP^\infty)$ let $\pi: \tilde{M} \rightarrow M$ be the double cover classified by f . If $[M, f] \in N_*(S^1)$ let $\pi: \tilde{M} \rightarrow M$ be the double cover corresponding to $f^* \iota + w_1 M$. In either case $\tilde{M}$ is a Wall manifold with

involution Δ , the deck transformation reversing sheets of $\tilde{M}$. This defines monomorphisms

$$k : W_*(RP^\infty) \rightarrow W_*^F, \quad j : N_*(S^1) \rightarrow W_*^F$$

with $k+j$ epic and $\text{Im } k \cap \text{Im } j \cong W_*$, identified as the set of classes $[W \times S^0, \text{Id} \times \Delta]$ for $[W] \in W_*$.

Now let $\lambda \rightarrow M$ be the line bundle corresponding to π . If $[M, f] \in N_*(S^1)$, $D\lambda$ is a Wall manifold. Define

$$\delta[\tilde{M}, \Delta] = [D\lambda] \in \tilde{W}_*(\vee TBO(n)).$$

If $[M, f] \in W_*(RP^\infty)$, let $i : N \subset M$ include a submanifold defining the double cover. $D(i^*\lambda)$ is a Wall manifold; by Lemma 1 so is $P = P(i^*\lambda \oplus 1)$. Define

$$\delta[\tilde{M}, \Delta] = [D(i^*\lambda \oplus 1)] + [D(1 \rightarrow M)].$$

LEMMA 2. $\partial\delta = \text{Id} : W_*^F \rightarrow W_*^F$.

PROOF. For $x \in \text{Im } j$ it is obvious that $\partial\delta(x) = x$. If $[M, f] \in W_*(RP^\infty)$ write $P = D(i^*\lambda) \cup D(i^*\lambda)$. The map $g : P \rightarrow RP^\infty$, classifying the canonical line bundle, may be written

$$(1) \quad g = (f|N)p_0 \cup *$$

where p_0 is the projection of $D(i^*\lambda)$ and $*$ is homotopic to a trivial map (see the proof of [3, (26.1)]). Since $i^*\lambda$ is the normal bundle of N in M we may also assume f is of the form (1). Now form (H, h) from $(M \times I, f\pi_M)$ and $(P \times I, g\pi_P)$ by sewing copies of $(D(i^*\lambda) \times 1, (f|N)p_0)$.

$\partial(H, h) = (M, f) \cup (P, g) \cup (M, *)$, which proves $\partial\delta[\tilde{M}, \Delta] = [\tilde{M}, \Delta]$. $\square$

3. **Proof of Theorem 1.** The statement of the theorem can be sharpened:

THEOREM 1'. ρ maps torsion $\mathcal{O}_*$ isomorphically onto $\text{Im } d'$.

LEMMA 3. Let (M, T) be an orientable manifold with involution, and $(M, T) = \partial(Q, U)$ as unoriented manifolds with involution. If Q is in fact orientable, then

$$[M, T] = 2\beta + \omega[S^0, \Delta]$$

for some $\beta \in \mathcal{O}_*^+$ and $\omega \in \Omega_*$.

PROOF. Give Q an orientation and let $\partial Q = M'$ be its oriented boundary.

$$(2) \quad [M, T] = [M, T] - [M', T].$$

Of course $M' = M$ up to the sign of the orientation on each component. $2x = 0$ for $x \in \mathcal{O}_*^-$; also the involution $M \cup (-M) \rightarrow M \cup (-M)$, reversing components, bounds $(M \times [-1, 1], \text{Id} \times (-1))$.

(2) then reduces to the desired form. $\square$

LEMMA 4. If $x \in \mathcal{O}_n$ and $\rho x = 0$ then either $x \in 2\mathcal{O}_n$ or else $n = 4i$ and

$$(3) \quad x = 2\beta + \omega[S^0, \Delta]$$

for some $\beta \in \mathcal{O}_{4i}^+$ and $\omega \in \Omega_{4i}/\text{torsion}$.

Note. This is [4, Proposition 21] and, essentially, [2, Theorem 5.8]. Following is a short proof relying on [6].

PROOF. Write $x = y + z$, $y \in \mathcal{O}_*^+$, $z \in \mathcal{O}_*^-$. Since spy and spz lie in orthogonal summands we must have $\rho y = 0 = \rho z$.

Let (M, T) represent either y or z , and let (W, S, f) be a Wall manifold with involution and boundary $(M, T, *)$. By [6, (4.1)] we may assume $f(M) = \{a\}$ and f is transverse regular at $b \neq a$. If $b \in \text{Im } f$, cut W along $N = f^{-1}b$, as in [8, §1], obtaining an oriented cobordism of (M, T) with $(N \times S^0, (S|N) \times \text{Id})$. This bounds if S is orientation reversing. If $b \notin \text{Im } f$, then W is orientable and Lemma 3 applies. Finally, Rosenzweig showed [5] that $2\beta = 0$ if $n \neq 4i$ and that $\omega[S^0, \Delta] = 0$ if $2\omega = 0$. $\square$

Suppose $x \in \mathcal{O}_n$, $\rho x = 0 = 2x$. If $x \in 2\mathcal{O}_n$ then $x = 0$, for the only torsion is of order 2. If x has the form (3), then since $2x = 0$ we have $4\beta = 2\omega[S^0, \Delta]$. By the proof of [2, Corollary 5.5], index (ω) is even. Then [2, Theorem 4.5] proves $\omega[S^0, \Delta] \in 2\mathcal{O}_{4i}^+$. Thus $x = 0$ in this case also, and ρ is monic on the torsion. The reader can easily show that $\rho(\text{torsion}) = \text{Im } d'$, completing Theorem 1'. $\square$

4. **Calculation of ∂' .** To give meaning to Theorems 2 and 3, we compute the homomorphism ∂' . The structure of $\tilde{W}_*(\bigvee TBO(n))$ is determined by [3, §17] and [7, pp. 163–164].

LEMMA 5. $\tilde{W}_*(\bigvee TBO(n))$ is the W_* -polynomial algebra on generators $[D\lambda_i]$, one of each dimension $i \geq 1$, with λ_i a line bundle. If $i = 2m + 1$, we may take λ_i to be the canonical line bundle over RP^{2m} . If $i = 2m + 2$, we may take λ_i to be the canonical line bundle over $P(\xi \oplus 2m \rightarrow S^1)$, where $\xi \rightarrow S^1$ is the nontrivial line bundle.

The proof is perfectly straightforward, and all the needed cohomology data is listed in [6, (4.1)].

Recall that $\partial'[D\nu] = [Di^*\nu]$, if $\nu \rightarrow F$ and $i: N \subset F$ includes a submanifold dual to $w_1\nu + w_1F$. By [7, pp. 169–172], $D\nu$ is orientable if and only if $[D\nu] \in \text{Im } \partial'$. For future reference, if ν has even fiber dimension then by [6, (2.4)] it is also true that $P(i^*\nu)$ is dual in $P\nu$ to $w_1P\nu$. Then

$$2[P(i^*\nu)] = 0 \in \Omega_*,$$

by [8].

LEMMA 6. For all m , $\partial'[D\lambda_{2m+2}]=[D\lambda_{2m+1}]$ and $\partial'[D\lambda_{2m+1}]=0$. If $\partial_1: W_* \rightarrow W_*$ maps W to the submanifold dual to $w_1 W$ [8], then

$$\partial'(w\beta) = (\partial_1 w)\beta + w(\partial'\beta)$$

for all $w \in W_*$, $\beta \in \tilde{W}_*(\bigvee TBO(n))$.

REMARKS. The first statement is immediate and the second may be proved with characteristic numbers, exactly like [8, Lemma 3]. From the proof of [6, (4.1)], we see that $\partial's = sd'$. Since s is a W_* -monomorphism, the lemma also proves that $d'(w\beta) = (\partial_1 w)\beta + w(d'\beta)$ for $w \in W_*$, $\beta \in W_*^I$.

5. **Proof of Theorems 2 and 3.** If ν has odd fiber dimension, then by [6, (3.3)] we have $\delta\partial[D\nu] = [D\lambda]$, where $\lambda \rightarrow P\nu$ is the canonical line bundle. By Lemma 1, $P(\nu \oplus 1)$ is a Wall manifold. Let $t(\nu)$ be the involution on $P(\nu \oplus 1)$ induced by the linear map $1 \times (-1)$ in each fiber of $\nu \oplus 1$, and define

$$\sigma[D\nu] = [P(\nu \oplus 1), t(\nu)] \in W_*^I.$$

Then $s\sigma[D\nu] + \delta\partial[D\nu] = [D\nu]$, just as in [3, §28]. It follows that $\sigma[D\nu] = 0$ if and only if $[D\nu] = \delta\partial[D\nu] \in \tilde{W}_*(TBO(1))$. Also, since $\partial s = 0$, it follows that $s\rho(x) = \rho(x)$ for $x \in \mathcal{O}_*^-$. Since $P(\nu \oplus 1)$ is orientable if $D\nu$ is, and $t(\nu)$ is then orientation reversing, Theorem 2 is proved.

If $\nu \rightarrow F$ has even fiber dimension then we will assume $[D\nu] \in \text{Ker } p$. To justify this restriction, suppose $[M, T] \in \mathcal{O}_*^+$ and ν is the normal bundle to the fixed set of T . Then $P\nu$ bounds $(M - \text{Int } D\nu)/T$, so $p\rho[M, T] = 0$.

Suppose, then, that $[D\nu] \in \text{Ker } p$. Since $P\nu$ is a Wall manifold, Lemma 2 implies $\delta[D\nu] = [D(i^*\lambda \oplus 1)]$, where $i: N \subset P\nu$ includes a submanifold dual to λ . We define $\sigma: \text{Ker } p \rightarrow W_*^I$. Let $\tilde{N} = S(i^*\lambda)$, so $\tilde{N} \rightarrow N$ is a double cover and $\tilde{N}$ is imbedded in $S\nu$ with trivial normal bundle. $\tilde{N}$ is also imbedded in $S(i^*\lambda \oplus 1)$ with trivial normal bundle, so we sew $D\nu$ and $D(i^*\lambda \oplus 1)$ along the copies of $\tilde{N} \times I$ in the boundaries. The remaining boundary is then two copies of $P\nu$, which bounds some Q . Sewing on $2Q$ builds a Wall manifold $K(\nu)$, which admits an involution $t(\nu)$ that restricts to the antipodal involution on the disk bundles, and reverses the two copies of Q . Thus the fixed set of $t(\nu)$ is the disjoint union of F and N , and the normal bundles are ν and $i^*\lambda \oplus 1$, respectively.

Define $\sigma[D\nu] = [K(\nu), t(\nu)] \in W_*^I$; it is easy to see that this is a well-defined homomorphism. By construction,

$$s\sigma[D\nu] + \delta\partial[D\nu] = [D\nu].$$

Reasoning as before, $\sigma[D\nu] = 0$ if and only if $[D\nu] = \delta\partial[D\nu] \in \text{Im } q$, and $s\rho(x) = \rho(x)$ for $x \in \mathcal{O}_*^+$.

Finally, suppose $D\nu$ is oriented. Then so are $D(i^*\lambda \oplus 1)$ and $P\nu$. As noted in the last section, $P\nu$ represents a torsion class in Ω_* . Since $\Omega_* \rightarrow W_*$ is monic on torsion [8] we may assume Q is oriented. Thus $K(\nu)$ is oriented, and $t(\nu)$ is orientation preserving. This completes the proof of Theorem 3.

REMARKS. (1) We have shown that $\mathcal{O}_*^-$ is the $\mathbb{Z}_2$ -vector space on generators $[P(\nu \oplus 1), t(\nu)]$, for $[D\nu] \in \text{Im } \partial'$, $\dim \nu = 2m+1$, $m > 0$. The torsion of $\mathcal{O}_*^+$ is a $\mathbb{Z}_2$ -vector space on generators $[K(\nu), t(\nu)]$ for

$$[D\nu] \in \text{Im } \partial'' \cap \text{Ker } p',$$

but this is not very helpful, since we cannot actually construct $K(\nu)$ without knowing $Q = \partial P\nu$.

(2) If $[M, T] \in \mathcal{O}_*^-$, then clearly $[M]$ is a torsion class in Ω_* . Conversely, P. G. Anderson proved [1, Proposition 5] that every torsion class in Ω_* is represented by one of our examples $P(\nu \oplus 1)$.

REFERENCES

1. P. G. Anderson, *Cobordism classes of squares of orientable manifolds*, Ann. of Math. (2) **83** (1966), 47–53. MR **32** #4700.
2. P. E. Conner, *Orientation-preserving involutions*, Lectures on the action of a finite group, Lecture Notes in Math., no. 73, Springer-Verlag, Berlin, 1968, Chap. 2. MR **41** #2670.
3. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Heft 33, Academic Press, New York; Springer-Verlag, Berlin, 1964, MR **31** #750.
4. C. N. Lee and A. G. Wasserman, *Equivariant characteristic numbers*, Proc. Second Conf. Transformation Groups, part I, Lecture Notes in Math., vol. 298, Springer-Verlag, Berlin, 1972, pp. 191–216.
5. H. L. Rosenzweig, *Bordism of involutions on manifolds*, Illinois J. Math **16** (1972), 1–10.
6. R. J. Rowlett, *Wall manifolds with involution*, Trans. Amer. Math. Soc. **169** (1972), 153–162.
7. R. E. Stong, *Notes on cobordism theory*, Princeton Univ. Press, Princeton, N.J.; Univ. of Tokyo Press, Tokyo, 1968. MR **40** #2108.
8. C. T. C. Wall, *Determination of the cobordism ring*, Ann. of Math. (2) **72** (1960), 292–311. MR **22** #11403.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540