ON A PROPERTY OF RATIONAL FUNCTIONS. II

Q. I. RAHMAN

ABSTRACT. It is shown that if $r_n(z)$ is a rational function of degree n such that $r_n(0)=1$, $\lim_{|z| \rightarrow \infty} |r_n(z)|=0$ and all its poles lie in $|\zeta_1| \leq |z| \leq 1$ then $\max_{|z|=\rho < |\zeta_1|} |r_n(z)| \geq 1/(1-\rho^n)$.

If $r_n(z)$ is a rational function of n th degree

$$(1) \quad r_n(z) = u(z)/v(z), \quad u, v \text{ polynomials}$$

with

$$(2) \quad \deg u(z) < \deg v(z) = n, \quad r_n(0) \neq 0$$

then [1] for $1 < \rho \leq 2$ and arbitrary complex z_0 ,

$$(3) \quad \|r_n\|_{p, z_0, \rho} = \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |r_n(z_0 + \rho e^{i\varphi})|^p d\varphi \right)^{1/p}$$

can be estimated nontrivially from below in the maximal regularity circle around z_0 exclusively by p , $|r_n(z_0)|$, ρ , n and by the maximum distance of z_0 from the poles. Here we obtain the sharp lower bound for

$$(4) \quad \|r_n\|_{\infty, z_0, \rho} = \max_{-\pi \leq \varphi < \pi} |r_n(z_0 + \rho e^{i\varphi})|.$$

Without loss of generality we may suppose $z_0=0$, $r_n(0)=1$; further if the poles are $\zeta_1, \zeta_2, \dots, \zeta_n$ (repetition permitted), then

$$(5) \quad 0 < |\zeta_1| \leq |\zeta_2| \leq \dots \leq |\zeta_n| = 1,$$

and

$$(6) \quad 0 < \rho < |\zeta_1|.$$

THEOREM 1. *With the above normalizations we have*

$$(7) \quad \max_{|z|=\rho} |r_n(z)| \geq \frac{1}{1-\rho^n}.$$

The example $r_n^*(z)=1/(1-z^n)$ shows that (7) is sharp.

Received by the editors December 11, 1972.

AMS (MOS) subject classifications (1970). Primary 30A06, 30A40.

Key words and phrases. Rational functions, logarithmic derivative of a polynomial.

© American Mathematical Society 1973

PROOF OF THEOREM 1. Let $\max_{|z|=\rho} |r_n(z)| = M(\rho)$. For every real α the function $u(z) - e^{i\alpha} M(\rho) v(z)$ does not vanish in $|z| < \rho$. Hence if

$$u(z) = 1 + \sum_{k=1}^{n-1} a_k z^k, \quad v(z) = 1 + \sum_{k=1}^n b_k z^k,$$

then

$$u(\rho z) - e^{i\alpha} M(\rho) v(\rho z) \equiv (1 - e^{i\alpha} M(\rho)) + (a_1 - e^{i\alpha} M(\rho) b_1)(\rho z) + \cdots \\ + (a_{n-1} - e^{i\alpha} M(\rho) b_{n-1})(\rho z)^{n-1} - e^{i\alpha} M(\rho) b_n (\rho z)^n$$

does not vanish in $|z| < 1$. Consequently, the coefficient of z^n does not exceed in absolute value the constant term, i.e. $|1 - e^{i\alpha} M(\rho)| \geq |-e^{i\alpha} M(\rho) b_n \rho^n|$. Choosing α appropriately, we get $M(\rho)(1 - |b_n| \rho^n) \geq 1$ or $M(\rho) \geq 1/(1 - |b_n| \rho^n)$. But $|b_n| \geq 1$ since the zeros of $v(z)$ lie in $|z| \leq 1$. Hence $M(\rho) \geq 1/(1 - \rho^n)$.

The condition (2) is automatically satisfied in the important special case when $r_n(z)$ is the logarithmic derivative of a polynomial $\pi_n(z)$ of degree n with all zeros in the unit disk; hence Theorem 1 gives a lower bound for $\max_{|z|=\rho} |\pi'_n(z)/\pi_n(z)|$ depending on $|\pi'_n(0)/\pi_n(0)|$, ρ , n . However, we show that in this case there is a lower bound which is independent of $|\pi'_n(0)/\pi_n(0)|$. In fact, we prove

THEOREM 2. Let the zeros $\zeta_1, \zeta_2, \dots, \zeta_n$ of the polynomial $\pi_n(z)$ of degree n be such that $0 < |\zeta_1| \leq |\zeta_2| \leq \dots \leq |\zeta_n| \leq 1$. Then for $0 \leq \rho < |\zeta_1|$

$$(8) \quad \max_{|z|=\rho} \left| \frac{\pi'_n(z)}{\pi_n(z)} \right| \geq \frac{n\rho^{n-1}}{1 - \rho^n}.$$

The example $\pi_n^*(z) = 1 - z^n$ shows that (8) is sharp.

PROOF OF THEOREM 2. Let $\pi_n(z) = \sum_{k=0}^n c_k z^k$ and set

$$A = \max_{|z|=\rho} |\pi'_n(z)/\pi_n(z)|.$$

Then

$$A = \max_{|z|=1} \left| \frac{\pi'_n(\rho z)}{\pi_n(\rho z)} \right| = \max_{|z|=1} \left| \frac{c_1 + 2c_2\rho z + \cdots + nc_n\rho^{n-1}z^{n-1}}{c_0 + c_1\rho z + c_2\rho^2 z^2 + \cdots + c_n\rho^n z^n} \right| \\ = \max_{|z|=1} \left| \frac{n\bar{c}_n\rho^{n-1} + (n-1)\bar{c}_{n-1}\rho^{n-2}z + \cdots + 2\bar{c}_2\rho z^{n-2} + \bar{c}_1 z^{n-1}}{c_0 + c_1\rho z + c_2\rho^2 z^2 + \cdots + c_n\rho^n z^n} \right| \\ = n \left| \frac{\bar{c}_n}{c_0} \right| \rho^{n-1} \max_{|z|=1} \left| \frac{1 + \frac{n-1}{n} \frac{\bar{c}_{n-1}}{\bar{c}_n} \frac{z}{\rho} + \cdots + \frac{2}{n} \frac{\bar{c}_2}{\bar{c}_n} \left(\frac{z}{\rho}\right)^{n-2} + \frac{\bar{c}_1}{\bar{c}_n} \left(\frac{z}{\rho}\right)^{n-1}}{1 + \frac{c_1}{c_0} \rho z + \frac{c_2}{c_0} \rho^2 z^2 + \cdots + \frac{c_n}{c_0} \rho^n z^n} \right|$$

$$\begin{aligned}
 &= n \left| \frac{c_n}{c_0} \right| \rho^{n-1} \max_{|z|=\rho} \left| \frac{1 + \frac{n-1}{n} \frac{\bar{c}_{n-1}}{\bar{c}_n} \left(\frac{z}{\rho^2} \right) + \cdots + \frac{2}{n} \frac{\bar{c}_2}{\bar{c}_n} \left(\frac{z}{\rho^2} \right)^{n-2} + \frac{\bar{c}_1}{\bar{c}_n} \left(\frac{z}{\rho^2} \right)^{n-1}}{1 + \frac{c_1}{c_0} z + \frac{c_2}{c_0} z^2 + \cdots + \frac{c_n}{c_0} z^n} \right| \\
 &\cong n \left| \frac{c_n}{c_0} \right| \rho^{n-1} \frac{1}{1 - \rho^n}
 \end{aligned}$$

by Theorem 1. But $|c_n/c_0| \geq 1$ since all the zeros of $\pi_n(z)$ lie in $z \leq 1$. Hence $A \geq n\rho^{n-1}/(1 - \rho^n)$.

The problems considered in this paper were suggested to me by Professor Paul Turán for which I am very thankful to him.

REFERENCE

1. Q. I. Rahman and Paul Turán, *On a property of rational functions*, Ann. Univ. Sci. Budapest. Eötvös Sect. Math. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MONTREAL, MONTREAL, QUEBEC, CANADA