ISOMETRIC EMBEDDING OF A COMPACT RIEMANNIAN MANIFOLD INTO EUCLIDEAN SPACE¹

HOWARD JACOBOWITZ

ABSTRACT. An isometric immersion of an n -dimensional compact Riemannian manifold with sectional curvature always less than λ^{-2} into Euclidean space of dimension $2n-1$ can never be contained in a ball of radius λ . This generalizes and includes results of Tompkins and Chern and Kuiper.

Tompkins [4] proved that the flat n -dimensional torus could not be isometrically embedded into E , the Euclidean space of dimension $2n-1$. Chern and Kuiper [1] and Otsuki [3] generalized this and showed that a compact n -dimensional manifold whose sectional curvatures are everywhere nonpositive also cannot be isometrically embedded in E . Apparently these results also hold for immersions. The purpose of this note is to point out that a standard proof of this nonembeddability result essentially establishes a more general quantitative version.

THEOREM. *Let E be Euclidean space of dimension $2n-1$ and M a compact n -dimensional Riemannian manifold whose sectional curvatures are everywhere less than some constant λ^{-2} . Then no isometric immersion of M into E is contained in a ball of radius λ .*

To prove the theorem, we adapt the proof for nonpositive curvature given in Kobayashi and Nomizu [2, pp. 26-29]. Note the theorem includes the results mentioned above.

Let $f: M \rightarrow E$ be an isometric embedding. Identify the tangent space $T_x M$ for $x \in M$ with its realization as a linear subspace of E . Denote the length and inner product of vectors in E by $|X|$ and $\langle X, Y \rangle$. To prove the theorem, let us assume such an isometric immersion did exist. We can assume $|f(x)| \leq \lambda$ while $|f(x_0)| = \lambda$ for some x_0 and all x in M . Thus $\langle f(x_0), X \rangle = 0$ for all X in $T_{x_0} M$. Let $L(X, Y)$ denote the second fundamental form at x_0 of M in E . For X and Y in $T_{x_0} M$, $L(X, Y)$ is a vector in E orthogonal to M at x_0 . The sectional curvature of the two-plane spanned by

Received by the editors December 28, 1972.

AMS (MOS) subject classifications (1970). Primary 53C40.

Key words and phrases. Isometric embedding, sectional curvature, Tompkin counter-example.

¹ This work was supported in part by NSF Grant GP-18961.

© American Mathematical Society 1973

linearly independent vectors X and Y in $T_{x_0}M$ is given by

$$K(X, Y) = (|X|^2 |Y|^2 - \langle X, Y \rangle^2)^{-1} \\ \cdot (\langle L(X, X), L(Y, Y) \rangle - \langle L(X, Y), L(X, Y) \rangle).$$

For a proof of the theorem it suffices to find two vectors X and Y such that $K(X, Y) \geq \lambda^{-2}$.

Differentiating $\langle f(x), f(x) \rangle$ twice at the maximum $x=x_0$, one obtains $\langle L(X, X), \nu \rangle \leq -\lambda^{-1}|X|^2$ for the unit normal $\nu = \lambda^{-1}f(x_0)$ and all $X \in T_{x_0}M$. In particular $L(X, X)=0$ only for $X=0$. Now for any symmetric bilinear form $B: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ with $B(X, X)=0$ only for $X=0$, one can find linearly independent vectors X and Y such that $B(X, X)=B(Y, Y)$ and $B(X, Y)=0$. Indeed just solve $B(Z, Z)=0$ where Z is allowed to be complex. This observation is due to T. A. Springer; see [2]. Let X and Y be two such vectors for L . Since $\langle L(X, X), \nu \rangle \leq -\lambda^{-1}|X|^2$ while $L(X, X)=L(Y, Y)$ one has $\langle L(X, X), L(Y, Y) \rangle \geq \lambda^{-2}|X|^2|Y|^2$. This together with $L(X, Y)=0$ implies the inequality $K(X, Y) \geq \lambda^{-2}$ and so proves the theorem.

If for the manifold M we only knew that for each x , $T_x M$ contained a q -dimensional subspace T'_x with the property that the sectional curvature of each two plane in T'_x was greater than λ^{-2} , then we could conclude that no isometric immersion of M into E^{n+q-1} is contained in a ball of radius λ . This result, for nonpositive curvatures, is due to Chern and Kuiper [1].

REFERENCES

1. S. S. Chern and N. H. Kuiper, *Some theorems on the isometric embedding of compact Riemann manifolds in euclidean space*, Ann. of Math. (2) **56** (1952), 422–430. MR **14**, 408.
2. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*. Vol. II, Interscience Tracts in Pure and Appl. Math., no. 15, Interscience, New York, 1969. MR **38** #6501.
3. T. Otsuki, *On the existence of solutions of a system of quadratic equations and its geometrical application*, Proc. Japan Acad. **29** (1953), 99–100. MR **15**, 647.
4. C. Tompkins, *Isometric embedding of flat manifolds in Euclidean space*, Duke Math. J. **5** (1939), 58–61.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, HOUSTON, TEXAS 77001