RESIDUALLY F_p -GROUPS, FOR MANY PRIMES p , ARE ORDERABLE

A. H. RHEMTULLA

ABSTRACT. It is proved that if G is a residually finite p -group, for infinitely many primes p , then G can be linearly ordered.

Let F_p denote the class of finite p -groups. It is well known that free groups and free polynilpotent groups are residually F_p for all primes p . It is also known that such groups can be linearly ordered ([2, p. 49], [1, Theorem 4]). The purpose of this note is to prove the following general result on the orderability of residually F_p -groups.

THEOREM. *If G is locally a residually F_p -group for infinitely many primes p , then G can be linearly ordered.*

To prove the Theorem we shall need the following result.

LEMMA. *Let α_{ij} ($i=1, \dots, 2^n; j=1, \dots, n$) be nonnegative integers such that $\sum_{j=1}^n \alpha_{ij} > 0$ for all i . Let p be a prime such that $p \geq (2\alpha)^{2^n}$ where $\alpha = \max\{\alpha_{ij}\}$. Then in any solution set to the system of 2^n equations $\sum_{j=1}^n \varepsilon_{ij} \alpha_{ij} x_j \equiv 0 \pmod{p}$ associated with signs $\varepsilon_{ij} = \pm 1$, $x_j \equiv 0 \pmod{p}$ for some j .*

PROOF. We use induction on n . The result is trivially true when $n=1$. Assume the result holds for $n < m$ and let $n=m$. Let $d=2^{m-1}$ and for any integer r let $r'=d+r$. By renumbering the 2^m equations, if necessary, assume that $\varepsilon_{i1}=1$, $\varepsilon_{i'1}=-1$ and $\varepsilon_{ij}=\varepsilon_{i'j}$ for all $i \in \{1, \dots, d\}$ and $j > 1$. We produce a system of d equations in $m-1$ unknowns satisfying the hypotheses of the Lemma as follows. For any $r \in \{1, \dots, d\}$ we have

$$I \quad \alpha_{r1}x_1 + \sum_{j=2}^m \varepsilon_{rj} \alpha_{rj} x_j \equiv 0 \pmod{p},$$

Received by the editors March 5, 1973.

AMS (MOS) subject classifications (1970). Primary 06A55; Secondary 20E25.

Key words and phrases. Residually finite p -groups, ordered groups, free polynilpotent groups.

Research supported by the National Research Council of Canada Grant No. A-5299.

© American Mathematical Society 1973

and

$$\text{II} \quad -\alpha_{r'1}x_1 + \sum_{j=2}^m \varepsilon_{rj}\alpha_{r'j}x_j \equiv 0 \pmod{p}.$$

As the r th equation of the new system pick: I if $\alpha_{r1}=0$; II if $\alpha_{r1}\neq 0$ but $\alpha_{r'1}=0$; $\sum_{j=2}^m \varepsilon_{rj}(\alpha_{r1}\alpha_{r'j} + \alpha_{r'1}\alpha_{rj})x_j \equiv 0 \pmod{p}$ if $\alpha_{r1}\neq 0$ and $\alpha_{r'1}\neq 0$. Notice that for any r if $\sum_{j=2}^m \alpha_{rj}=0$ then $x_1 \equiv 0 \pmod{p}$ and the Lemma is proved. Thus we may assume that at least one coefficient of the r th equation of the new system is positive. Let β be the maximum of the coefficients of the new system of equations. Then $\beta \leq (2\alpha^2)$ so that $p \geq (2\alpha)^{2^m} = (4\alpha^2)^{2^{m-1}} \geq (2\beta)^{2^{m-1}}$. Thus the hypotheses of the Lemma are satisfied for this system and, by induction, $x_j \equiv 0 \pmod{p}$ for some j .

PROOF OF THE THEOREM. We use a characterization of ordered groups given by Fuchs [2, p. 34]. If x is an element of a group G and α a non-negative integer, we write $[x]^\alpha$ to denote a product of α conjugates of x in G .

Suppose, if possible, that the Theorem is false. Then there exist elements $x_i \neq e$ ($i=1, \dots, n$) in G such that

$$(1) \quad [x_1^{\varepsilon_{i1}}]^{\alpha_{i1}} \cdots [x_n^{\varepsilon_{in}}]^{\alpha_{in}} = e$$

for 2^n values of i associated with signs $\varepsilon_{ij} = \pm 1$. Moreover $\sum_{j=1}^n \alpha_{ij} > 0$ for all i . Assume that n is the least such integer. Let $\alpha = \max\{\alpha_{ij}\}$. We can assume that G is generated by the finite set of elements involved in the equations given by (1), so that G is residually F_p for infinitely many p . Choose a prime $p > (2\alpha)^{2^n}$ such that G is residually F_p for this prime. Let $X = \langle x_1^G, \dots, x_n^G \rangle$, the normal subgroup of G generated by $x_1, \dots, x_n$. There exists $K \triangleleft G$ maximal subject to $G/K \in F_p$, $X \not\leq K$.

Let θ be the homomorphism $G \rightarrow G/K = G^*$, and let $\theta(x_i) = a_i$, $\theta(X) = A$. Then $G^* \in F_p$, $A \neq \{e\}$, and, by our choice of K , A is a minimal normal subgroup of G^* . Thus A is cyclic of order p in the centre of G^* . By re-ordering the suffixes, if necessary, assume that $A = \langle a_1 \rangle$, $a_j = a_1^{\beta_j}$ where $\beta_1 = 1$, $0 < \beta_j < p$ for $j=2, \dots, m$ and $\beta_j = 0$ for $j > m$. The 2^n equations given by (1) now reduce to the following.

$$a_1^{(\sum_{j=1}^m \varepsilon_{ij} \alpha_{ij} \beta_j)} = e, \quad j = 1, \dots, m, i = 1, \dots, 2^n.$$

If $m < n$, then from the 2^n equations above choose 2^m equations associated with the 2^m choices of signs $\varepsilon_{ij} = \pm 1$ ($j=1, \dots, m$) in such a way that for each $i \in \{1, \dots, 2^m\}$ at least one $\alpha_{ij} \neq 0$ in this set of 2^m equations. This is possible by our choice of n . The above equations now reduce to

$$\sum_{j=1}^m \varepsilon_{ij} \alpha_{ij} \beta_j \equiv 0 \pmod{p}, \quad i = 1, \dots, 2^m.$$

The above equations satisfy the hypotheses of the Lemma and hence $\beta_j \equiv 0 \pmod{p}$ for some $j \in \{1, \dots, m\}$. This is a contradiction and the Theorem is proved.

REFERENCES

1. G. Baumslag, *Wreath products and extensions*, Math. Z. **81** (1963), 286–299. MR 27 #1503.
2. L. Fuchs, *Partially ordered algebraic systems*, Pergamon Press, New York; Addison-Wesley, Reading, Mass., 1963. MR 30 #2090.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA,
CANADA