

# AN EXTENSION OF A THEOREM OF EISENSTEIN

L. KUIPERS

**ABSTRACT.** In the present paper I obtain an extension of the so-called Eisenstein theorem which is proved by means of Riemann-Stieltjes integration.

**THEOREM.** *Let  $n$  be a positive integer  $\geq 2$ . Let  $p_1, p_2, \dots, p_n$  be odd integers and relatively prime in pairs. Then*

$$\sum_{i=1}^n \sum_{x=1}^{(p_i-1)/2} \prod_{k=1; k \neq i}^n \left[ \frac{p_k x}{p_i} \right] = \frac{1}{2^n} \prod_{k=1}^n (p_k - 1).$$

**PROOF.** The method of proof is Riemann-Stieltjes integration. Consider the integrals

$$\mathcal{I}_1 = \int_0^{1/2} [p_1 x] [p_2 x] \cdots [p_{n-1} x] d[p_n x],$$

$$\mathcal{I}_2 = \int_0^{1/2} [p_n x] d[p_1 x] [p_2 x] \cdots [p_{n-1} x].$$

First we observe that the greatest integer functions  $[p_1 x], [p_2 x], \dots, [p_n x]$  have no discontinuities in common on the interval  $0 < x \leq \frac{1}{2}$  in view of the condition on the integers  $p_1, p_2, \dots, p_n$ . Second, the discontinuities of  $[p_i x]$ ,  $i = 1, 2, \dots, n$ , on the interval  $0 < x \leq \frac{1}{2}$  are at  $x = k/p_i$ ,  $k = 1, 2, \dots, \frac{1}{2}(p_i - 1)$ , and the jump of  $[p_i x]$  at each of these discontinuities is equal to 1. Hence the value of  $\mathcal{I}_1$  is

$$\sum_{x=k/p_n} [p_1 x] [p_2 x] \cdots [p_{n-1} x], \quad k = 1, 2, \dots, \frac{1}{2}(p_n - 1)$$

or

$$\sum_{x=1}^{(p_n-1)/2} \left[ \frac{p_1 x}{p_n} \right] \left[ \frac{p_2 x}{p_n} \right] \cdots \left[ \frac{p_{n-1} x}{p_n} \right].$$

Consider the second integral. The discontinuities of  $[p_1 x] [p_2 x] \cdots [p_{n-1} x]$  are at  $x = k/p_i$ ,  $k = 1, 2, \dots, \frac{1}{2}(p_i - 1)$ ,  $i = 1, 2, \dots, n-1$ . Since there are no common discontinuities and the jumps of  $[p_i x]$  at the discontinuities  $k/p_i$  are equal to 1, the jump of  $[p_1 x] [p_2 x] \cdots [p_{n-1} x]$  at each

Received by the editors May 9, 1973.

AMS (MOS) subject classifications (1970). Primary 10A20, 10A25.

© American Mathematical Society 1974

$k/p_i$  is equal to

$$\left[ \frac{p_1 k}{p_i} \right] \left[ \frac{p_2 k}{p_i} \right] \dots \left[ \frac{p_{i-1} k}{p_i} \right] \left[ \frac{p_{i+1} k}{p_i} \right] \dots \left[ \frac{p_{n-1} k}{p_i} \right]$$

and so we obtain the value of the second integral

$$\mathcal{J}_2 = \sum_{i=1}^{n-1} \sum_{x=1}^{(p_i-1)/2} \prod_{k=1; k \neq i}^n \frac{p_k x}{p_i}.$$

This proves the theorem completely.

REMARK. For the history of the so-called Eisenstein theorem ( $n=2$  in our Theorem) and other generalizations we refer to Bruce C. Berndt's paper: *A generalization of a theorem of Gauss on  $[x]$* , presented at the Third Illinois Conference on Number Theory on April 7, 1973.

As is well known the so-called Eisenstein theorem which says: *If  $p$  and  $q$  are two odd distinct primes then*

$$\sum_{x=1}^{(q-1)/2} \left[ \frac{px}{q} \right] + \sum_{x=1}^{(p-1)/2} \left[ \frac{qx}{p} \right] = (1/4)(p-1)(q-1)$$

is very helpful in proving Gauss's quadratic reciprocity theorem.

Finally, thanks are due to the referee for pointing out to me a more general statement than first submitted.

DEPARTMENT OF MATHEMATICS, SOUTHERN ILLINOIS UNIVERSITY, CARBONDALE,  
ILLINOIS 62901