CORRECTIONS TO
"AN ANALYTICAL CRITERION FOR THE COMPLETENESS
OF RIEMANNIAN MANIFOLDS"

WILLIAM B. GORDON

A recent article in these *Proceedings* [4] was concerned with the following theorem.

Theorem. *In order that a smooth riemannian manifold M be complete, it is necessary and sufficient that M support a smooth proper function f whose gradient is bounded in modulus.*

The purpose of this present note is to call attention to the fact that our proof of the *necessity* part of the theorem contains a serious error, and that a correct proof of this part of the theorem is contained in an article by M. P. Gaffney [3]. (See also [1] and [2] for some earlier results which were tending in this direction.)

Our proof of *necessity* used the "fact" that an isometric embedding $j: M \rightarrow R^q$ is closed if the induced metric is complete. This is easily shown to be false; e.g., the embedding of R^1 into R^2 given by $t \rightarrow (e^t, \sin t)$ is not closed but the induced metric is complete. (On the other hand, the converse statement is true; i.e. a metric induced by a closed embedding is necessarily complete.)

A correct proof can be obtained as follows: Let p be a fixed point of M and let $f(x) = d(x, p)$, where d is the distance function corresponding to the given riemannian metric. Then the completeness of M implies that f is proper. (This is merely a restatement of that portion of the Hopf-Rinow theorem which asserts that a riemannian manifold is complete if and only if closed and bounded sets are compact.) The function f is not differentiable, but it is Lipshitz continuous with Lipshitz constant = 1; i.e., $|f(x) - f(y)| \leq d(x, y)$. Using this fact, Gaffney [3] shows, by means of various smoothing operations and partition of unity constructions, that for any positive ϵ there exists a smooth

Received by the editors April 24, 1973.

AMS (MOS) subject classifications (1970). Primary 53C20, 58A05; Secondary 70G05.

Copyright © 1974, American Mathematical Society

approximation $\hat{f}$ to f which satisfies (i) $|\hat{f}(x) - f(x)| < \epsilon$ and (ii) $|\hat{f}(y) - \hat{f}(x)| < (1 + \epsilon)d(x, y)$. The first relation implies that $\hat{f}$ is proper, and the second implies that $\|\nabla \hat{f}\| \leq 1 + \epsilon$.

BIBLIOGRAPHY

1. M. P. Gaffney, *The harmonic operator for exterior differential forms*, Proc. Nat. Acad. Sci. U.S.A. 37 (1951), 48–50. MR 13, 987.
2. ———, *A special Stoke's theorem for complete Riemannian manifolds*, Ann. of Math. (2) 60 (1954), 140–145. MR 15, 986.
3. ———, *The conservation property of the heat equation on Riemannian manifolds*, Comm. Pure Appl. Math. 12 (1959), 1–11. MR 21 #892.
4. W. B. Gordon, *An analytical criterion for the completeness of riemannian manifolds*, Proc. Amer. Math. Soc. 37 (1973), 221–225. MR 46 #6233.

MATHEMATICS RESEARCH CENTER, NAVAL RESEARCH LABORATORY, WASHINGTON, D. C. 20390