ESSENTIAL MAPS AND EMBEDDINGS OF ANNULI IN NONORIENTABLE M^3

R. J. DAIGLE AND C. D. FEUSTEL¹

ABSTRACT. In this note we give an example which shows that the existence of an essential map of an annulus in a nonorientable 3-manifold does not guarantee the existence of an essential embedding in that manifold.

I. Introduction. It has been reported by F. Waldhausen that the existence of an "essential" map of an annulus into a compact orientable 3-manifold guarantees the existence of an essential embedding of an annulus in that 3-manifold. It is natural to ask if the result above is true for nonorientable 3-manifolds. In this note we give an example to show that the answer to the question above is in general no.

Feustel has proved [2] that the existence of a certain type of essential map of an annulus into a compact (not necessarily orientable) 3-manifold guarantees the existence of an embedding having the same properties. Our example also shows that Feustel's conditions are necessary in case the 3-manifold is not orientable.

II. Notation. We shall let A denote an annulus, F a mobius band, S^1 the 1-sphere, and M a 3-manifold. All spaces will be simplicial complexes and all maps will be piecewise linear. We shall denote the boundary of a manifold N by ∂N and the components of ∂A by c_1 and c_2 . We let α be an arc embedded in A such that $c_j \cap \alpha$ is an endpoint of α for $j = 1, 2$. Such an arc is a *spanning arc* of A . A map $f: (A, \partial A) \rightarrow (M, \partial M)$ is *essential* if

- (1) $f_*: \pi_1(A) \rightarrow \pi_1(M)$ is monic,
- (2) the arc $f(\alpha)$ is not homotopic rel its boundary to an arc in ∂M .

We remark that condition (2) above is independent of our choice of the

Received by the editors May 8, 1973.

AMS (MOS) subject classifications (1970). Primary 55A25; Secondary 57C35.

Key words and phrases. Essential map, essential embedding.

¹ This author is partially supported by NSF Grant GP 15357.

spanning arc α . A connected surface S embedded in a manifold M is *incompressible* in M if S is not the 2-sphere and the natural map $i_*: \pi_1(S) \rightarrow \pi_1(M)$ induced by inclusion is monic.

We shall say that a simple loop λ_1 in the interior of $F \times [0, 1]$ is *knotted* in $F \times [0, 1]$ if the complement of the interior of a regular neighborhood R of λ_1 in $F \times [0, 1]$ is not homeomorphic to the product to a klein bottle with $[0, 1]$. Let λ be a 1-sided simple loop in the interior of F and t a point in $[0, 1]$. One can find a knotted loop λ_1 in $F \times [0, 1]$ by altering the loop $\lambda \times \{t\}$ inside a regular neighborhood of one of the points on $\lambda \times \{t\}$.

Let F_1 and F_2 be surfaces in M or ∂M . Then F_1 is *parallel* to F_2 if and only if there exists an embedding of $F_1 \times [0, 1]$ in M such that $F_1 = F_1 \times \{0\}$, and F_2 is the closure of $\partial(F_1 \times [0, 1]) - F_1 \times \{0\}$.

III. Construction of the manifold M . Let x be a point in S^1 and $M_1 = F \times S^1$. Let R be a regular neighborhood of $\lambda \times \{x\}$ in M_1 . Then R is homeomorphic to $F \times [0, 1]$. Let $\mathcal{D}$ be a disk embedded in R such that $\mathcal{D} \cap \partial R = \partial \mathcal{D}$ and $R - \mathcal{D}$ is connected. Let λ_1 be a knotted loop in R such that λ_1 meets $\mathcal{D}$ in a single point and crosses $\mathcal{D}$ at that point. Let R_1 be a regular neighborhood of λ_1 in R . Denote ∂R_1 by K_1 and ∂R by K . Let M_2^1 be the 3-manifold obtained by removing the interior of R_1 from M_1 . Let $h: M_2^1 \rightarrow M_2^2$ be a homeomorphism and M the union of M_2^1 and M_2^2 with the identification $y = h(y)$ for $y \in K_1$.

IV. There is an essential map of A into M . Let $p: (A, \partial A) \rightarrow (F, \partial F)$ be a covering map. Let x_0 be a point in S^1 such that R does not meet $F \times \{x_0\} \subset M_1$. Then we may define $f: (A, \partial A) \rightarrow (M_2^1, \partial M_2^1)$ by $f(s) = (p(s), x_0)$. We claim that f is essential. Clearly condition (1) is satisfied since the natural map $i_*: \pi_1(M_2^1) \rightarrow \pi_1(M)$ induced by inclusion is monic.

Let $(\hat{M}, \hat{p})$ be the orientable double cover of M . Now there is a map $\hat{f}: A \rightarrow \hat{M}$ such that $\hat{p}\hat{f} = f$. It can be seen that $\hat{f}(c_1)$ and $\hat{f}(c_2)$ lie on different components of $\partial \hat{M}$. It follows that $f(\alpha)$ is not homotopic rel its boundary to an arc in ∂M since such a homotopy could be lifted to $\hat{M}$.

V. There are no essential embeddings of A in M . Suppose that $g: (A, \partial A) \rightarrow (M, \partial M)$ is an essential embedding. We claim that we may assume that $g(A)$ does not meet K_1 . After a general position argument, we may assume that $g^{-1}(K_1)$ is a collection (possibly empty) of simple closed loops. Since K_1 and $g(A)$ are incompressible in M , any loop in $K_1 \cap g(A)$, nullhomotopic in M , is nullhomotopic both on $g(A)$ and on K_1 . It follows from

standard arguments that we may suppose that no loop in $K_1 \cap g(A)$ is null-homotopic in M . Similarly we may suppose that $K \cap g(A)$ and $h(K) \cap g(A)$ are collections of simple loops and that no loop in either collection is null-homotopic in M . We suppose that $g(A)$ meets K_1 .

Note that up to isotopy, the only simple essential loop on $T = \partial M_2^1 \cap \partial M$ which is homologous to a loop on K is $\partial F \times \{x_0\}$. We may assume that $g(A)$ meets T and that there is an annulus $A_1 \subset g(A)$ such that $A_1 \cap (T \cup K) = \partial A_1$. Thus $\partial A_1 \cap T$ is twice the generator of $\pi_1(R)$. Let $A_2 \subset g(A)$ be an annulus such that $A_2 \cap (K \cup K_1) = \partial A_2$ and A_2 meets both K_1 and K . Since $\partial A_1 \cap \partial A_2 \cap K$ is either empty or all of $\partial A_1 \cap K$ and ∂A_1 is orientable, standard arguments will show that λ_1 must have been unknotted. It follows that we may assume that $g(A)$ does not meet K_1 and that $g(\partial A) \subset \partial M_2^1$.

It is now clear that we may regard g as an essential embedding of A in M_2^1 or simply $M_1 - \lambda_1$. Let $(\hat{M}, \hat{p})$ be the orientable double cover of M_1 . Since $g(A) \subset M_1 - \lambda_1$ and $[g(c_1)] \in \hat{p}_* \pi_1(\hat{M}_1)$, we can find a map $\hat{g}: A \rightarrow \hat{M}_1$ such that $\hat{p}\hat{g} = g$ and $\hat{g}(A) \subset \hat{M}_1 - \hat{p}^{-1}\lambda_1$. Suppose that $\hat{g}(\partial A)$ lies on a single component of $\partial \hat{M}_1$. Let $\rho: \hat{M}_1 \rightarrow \hat{M}_1$ be the nontrivial covering translation of $\hat{M}_1$. Since $\hat{M}_1$ is homeomorphic to the product of a torus T with $[0, 1]$, it will follow from [3, Corollary 3.2] that $\hat{g}(A)$ and $\rho\hat{g}(A)$ are parallel to annuli in $\partial \hat{M}_1$. Since $\hat{p}^{-1}\lambda_1$ is connected and $\hat{g}(\partial A)$ and $\rho\hat{g}(\partial A)$ lie on distinct components of $\partial \hat{M}_1$, it will follow that either $\hat{g}(\alpha)$ or $\rho\hat{g}(\alpha)$ is homotopic rel its boundary to an arc in $\partial \hat{M}_1$ in the complement of $\hat{p}^{-1}\lambda_1$. One can project this homotopy to show that $g(A)$ was not essential.

Suppose that $\hat{g}$ carries c_1 and c_2 to distinct components of $\partial \hat{M}_1$. We observe that $\hat{g}(A) \cap \rho\hat{g}(A)$ is empty since g is an embedding. Let N_1 and N_2 be the closures of the components of the complement of $\hat{M}_1 - (\hat{g}(A) \cup \rho\hat{g}(A))$. Then we may suppose that $\hat{p}^{-1}\lambda_1 \subset N_1$ since $\hat{p}^{-1}(\lambda_1)$ is connected. We claim that $\rho N_1 = N_1$. Suppose that there is a point $z \in (N_1 - \rho N_1)$. Then $\rho z \in N_2$ and we can find a path β in N_1 from z to λ_1 which does not meet $\hat{g}(A) \cup \rho\hat{g}(A)$. But now $\rho\beta$ is a path which does not meet $\hat{g}(A) \cup \rho\hat{g}(A)$. It follows that $\rho\beta \subset N_2$ and $\hat{p}^{-1}(\lambda_1)$ meets N_2 . This is impossible. Since $\rho(N_1 - \rho N_1) = \rho N_1 - N_1$, $N_1 = \rho N_1$. Similarly $\rho N_2 = N_2$.

It is now clear that $g(A)$ separates M_1 into two components whose closures are $X_1 = pN_1$ and $X_2 = pN_2$. We obtain the commutative diagram shown in Figure 1 where all homomorphisms are induced by inclusion. Since $\pi_1(M_1) = z \oplus z$, $\pi_1(M_1)$ is not a nontrivial free product with amalgamation. Thus i_1 or i_2 must be onto. Since $\lambda_1 \subset X_1$, i_2 is onto. It follows from

Figure 1

[1, Theorem (1.1)] that X_2 is homeomorphic to $A \times [0, 1]$ and that $g(\alpha)$ is homotopic in X_2 rel its boundary to an arc in ∂M_1 .

Thus g was not essential.

REFERENCES

1. E. M. Brown and R. H. Crowell, *Deformation retractions of 3-manifolds into their boundaries*, *Ann. of Math. (2)* 82 (1965), 445–458. MR 31 #6236.
2. C. D. Feustel, *On embedding essential annuli in M^3* , *Canad. J. Math.* (to appear).
3. F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, *Ann. of Math. (2)* 87 (1968), 56–88. MR 36 #7146.

DEPARTMENT OF MATHEMATICS, VIRGINIA POLYTECHNIC INSTITUTE AND
STATE UNIVERSITY, BLACKSBURG, VIRGINIA 24061