SYLOWIZERS IN LOCALLY FINITE GROUPS

M. J. TOMKINSON

ABSTRACT. A result of W. Gaschütz for finite soluble groups is extended to two classes of locally finite, locally soluble groups.

The concept of Sylowizers has been introduced by W. Gaschütz [2]. If R is a π -subgroup of the group G , then a π -Sylowizer of R in G is a subgroup S of G maximal with respect to containing R as a Sylow π -subgroup. [π denotes a set of primes and a Sylow π -subgroup is simply a maximal π -subgroup.] A straightforward Zorn's lemma argument shows that π -Sylowizers of any π -subgroup R must always exist. Gaschütz proved the following conjugacy theorem:

Let G be a finite soluble group and R a normal subgroup of some Sylow π -subgroup P of G . Then the π -Sylowizers of R in G are conjugate in G .

It is our aim in this note to extend this result to the class $\mathfrak{G}$ of periodic locally soluble FC -groups and the class $\mathfrak{U}$ defined by:

$G \in \mathfrak{U}$ if and only if G is locally finite and for each $H \leq G$ and for each set of primes π , the Sylow π -subgroups of H are conjugate in H . [The necessary results about $\mathfrak{G}$ and $\mathfrak{U}$ may be found in [5] and [1] respectively.]

The proof for finite groups involves the usual consideration of a counterexample of minimal order. This method cannot be employed for infinite groups and we make a more direct construction of the π -Sylowizers of R , although this construction is based on the ideas used in Gaschütz's proof.

Although we only prove the extension of the theorem for $\mathfrak{G}$ and $\mathfrak{U}$, the construction of the π -Sylowizers is carried out in a much wider class of groups. We define $\mathfrak{X}_\pi$ to be the class of upper π -separable locally finite groups G such that PK/K is a Sylow π -subgroup of H/K whenever $K \triangleleft H \leq G$ and P is a Sylow π -subgroup of H .

A $\mathfrak{G}$ -group is clearly upper π -separable and also its Sylow π -subgroups have the necessary homomorphism property [5, 4.1 (iii)]. Also $\mathfrak{U}$ -groups are

Received by the editors April 23, 1973 and, in revised form, August 7, 1973.

AMS (MOS) subject classifications (1970). Primary 20E25.

Key words and phrases. Locally finite groups, FC -groups, $\mathfrak{U}$ -groups, Sylow subgroups.

Copyright © 1974, American Mathematical Society

upper π -separable for all π [3, Theorem E] and the second property is given in [1, 2.1 (ii)]. Thus $\mathfrak{X}_\pi$ is a QS-closed class containing both $\mathfrak{G}$ and $\mathfrak{U}$.

In the construction of the Sylowizers of R we shall require the following elementary result stated in [2] for finite groups.

Lemma 1. *Let R be a π -subgroup of the $\mathfrak{X}_\pi$ -group G and $N \triangleleft G$.*

(a) *If $N \leq R$ then S is a π -Sylowizer of R in G if and only if $N \leq S$ and S/N is a π -Sylowizer of R/N in G/N .*

(b) *If N is a π' -group then S is a π -Sylowizer of R in G if and only if $N \leq S$ and S/N is a π -Sylowizer of RN/N in G/N .*

This lemma shows immediately in the finite case that if G is a minimal counterexample to the theorem then $O_{\pi'}(G) = 1$ and there are no normal subgroups of G contained in R . We show that in any $\mathfrak{X}_\pi$ -group we can consider a section G_ρ/R_ρ and the π -subgroup RR_ρ/R_ρ with these properties.

We define four chains of subgroups in G depending only on the π -subgroup R . These chains are defined inductively as follows:

$$R_0 = 1, \quad G_0 = G, \quad Q_0 = O_{\pi'}(G) \text{ and } P_0 = O_{\pi'}(G).$$

$$R_{\alpha+1} = P_\alpha \cap RQ_\alpha, \quad G_{\alpha+1} = N_{G_\alpha}(R_{\alpha+1}), \quad Q_{\alpha+1}/R_{\alpha+1} = O_{\pi'}(G_{\alpha+1}/R_{\alpha+1}) \text{ and } P_{\alpha+1}/R_{\alpha+1} = O_{\pi'}(G_{\alpha+1}/R_{\alpha+1}).$$

Clearly

$$R_{\alpha+1} \geq R_\alpha, \quad G_{\alpha+1} \leq G_\alpha, \quad Q_{\alpha+1} \geq R_{\alpha+1} \geq Q_\alpha \text{ and } P_{\alpha+1} \geq P_\alpha.$$

If α is a limit ordinal, we define

$$R_\alpha = \bigcup_{\beta < \alpha} R_\beta, \quad G_\alpha = \bigcap_{\beta < \alpha} G_\beta, \quad Q_\alpha = \bigcup_{\beta < \alpha} Q_\beta \text{ and } P_\alpha = \bigcup_{\beta < \alpha} P_\beta.$$

It is clear that, for each ordinal α , R_α , Q_α and P_α are normal subgroups of G_α and $R_\alpha \leq Q_\alpha \leq R_{\alpha+1} \leq P_\alpha$.

Let ρ be the first ordinal such that $R_\rho = R_{\rho+1}$. (Such an ordinal must exist, for otherwise $|R_\alpha|$ would be greater than or equal to the cardinality of α , for all α , and this is clearly not possible if the cardinality of α is greater than $|G|$.) Then $G_{\rho+1} = N_{G_\rho}(R_{\rho+1}) = G_\rho$, $Q_{\rho+1}/R_\rho = O_{\pi'}(G_\rho/R_\rho)$ and so $Q_{\rho+1} = Q_\rho$ and similarly $P_{\rho+1} = P_\rho$. Thus all four chains become stationary at the ordinal $\rho = \rho(R)$.

Lemma 2. *G_ρ contains $N_G(R)$ and R is a Sylow π -subgroup of RR_ρ .*

Proof. By induction on ρ . The result is clearly true if $\rho = 0$ and so we may assume that $N_G(R) \leq G_\alpha$ and R is a Sylow π -subgroup of RR_α for each $\alpha < \rho$.

If ρ is a limit ordinal, then the result follows easily from $G_\rho = \bigcap_{\alpha < \rho} G_\alpha$ and $R_\rho = \bigcup_{\alpha < \rho} R_\alpha$. Therefore we may assume that $\rho = \alpha + 1$.

Since $N_G(R) \leq G_\alpha$, $N_G(R)$ normalizes P_α and Q_α and hence normalizes $R_\rho = P_\alpha \cap RQ_\alpha$. Thus $N_G(R) \leq G_\rho$.

$RR_\rho = R(P_\alpha \cap RQ_\alpha) = R(P_\alpha \cap R)Q_\alpha = RQ_\alpha$. Q_α/R_α is a π' -group and R is a Sylow π -subgroup of RR_α . It follows easily that R is a Sylow π -subgroup of $RQ_\alpha = RR_\rho$.

Lemma 3. *Let R be a π -subgroup of the $\mathfrak{X}_\pi$ -group G and S a π -Sylowizer of R in G . If $K \leq S \leq H$ and $K \triangleleft H$, then S/K is a π -Sylowizer of RK/K in H/K .*

Proof. Since R is a Sylow π -subgroup of S , it follows that RK/K is a Sylow π -subgroup of S/K . To show that S/K is maximal in H/K with this property, suppose that RK/K is a Sylow π -subgroup of T/K and $S \leq T \leq H$. Let R^* be a Sylow π -subgroup of T containing R . Then R^*K/K is a π -subgroup of T/K containing RK/K . Thus $R^*K = RK$ and $R \leq R^* \leq RK \leq S$. But R is a Sylow π -subgroup of S and so $R = R^*$ is a Sylow π -subgroup of T . By the maximality of S , we have $T = S$, as required.

We are now able to show that we need only consider the section G_ρ/R_ρ .

Theorem 1. *Let R be a π -subgroup of the $\mathfrak{X}_\pi$ -group G . Then, with the notation above, S is a π -Sylowizer of R in G if and only if $R_\rho \leq S \leq G_\rho$ and S/R_ρ is a π -Sylowizer of RR_ρ/R_ρ in G_ρ/R_ρ .*

Proof. Suppose first that S is a π -Sylowizer of R in G . By Lemma 3, it is sufficient to show that $R_\rho \leq S \leq G_\rho$. We prove this by induction on ρ . It is clearly true if $\rho = 0$ and so we may assume that $R_\alpha \leq S \leq G_\alpha$ for all ordinals $\alpha < \rho$. If ρ is a limit ordinal, then the result follows easily from $G_\rho = \bigcap_{\alpha < \rho} G_\alpha$ and $R_\rho = \bigcup_{\alpha < \rho} R_\alpha$. Therefore we may assume that $\rho = \alpha + 1$.

$R_\alpha \leq S \leq G_\alpha$ and so, by Lemma 3, S/R_α is a π -Sylowizer of RR_α/R_α in G_α/R_α . Q_α/R_α is a normal π' -subgroup of G_α/R_α and so $Q_\alpha \leq S$ (Lemma 1 (ii)). Thus $S \geq RQ_\alpha \geq R_{\alpha+1} = R_\rho$.

RQ_α/Q_α is a Sylow π -subgroup of S/Q_α and so the normal π -subgroup $(S \cap P_\alpha)/Q_\alpha$ of S/Q_α is contained in RQ_α/Q_α . Hence $S \cap P_\alpha = RQ_\alpha \cap P_\alpha = R_\rho$ and so $R_\rho \triangleleft S$, i.e. $S \leq G_\rho$.

Conversely, let S/R_ρ be a π -Sylowizer of RR_ρ/R_ρ in G_ρ/R_ρ . Since R is a Sylow π -subgroup of RR_ρ (Lemma 2), it follows that R is a Sylow π -subgroup of S . Thus S is contained in a π -Sylowizer T of R in G . By the first part of the proof, $T \leq G_\rho$ and T/R_ρ is a π -Sylowizer of RR_ρ/R_ρ in G_ρ/R_ρ . Hence $T = S$, as required.

Theorem 2. *Let R be a normal subgroup of the Sylow π -subgroup P of*

the $\mathfrak{X}_\pi$ -group G . Then S is a π -Sylowizer of R in G if and only if S/R_ρ is a Sylow π' -subgroup of G_ρ/R_ρ .

Proof. Since $Q_\rho = R_\rho = Q_{\rho+1}$, G_ρ/R_ρ has no normal π' -subgroups and so $P_\rho/R_\rho = O_\pi(G_\rho/R_\rho)$. Since $N_G(R) \leq G_\rho$ (Lemma 2), we have $P \leq G_\rho$. PR_ρ/R_ρ is a Sylow π -subgroup of G_ρ/R_ρ and so contains the normal π -subgroup P_ρ/R_ρ . P_ρ and RR_ρ are both normal in PR_ρ and so $[P_\rho, RR_\rho] \leq P_\rho \cap RR_\rho = R_\rho$, i.e. RR_ρ centralizes P_ρ/R_ρ . But $C_{G_\rho}(P_\rho/R_\rho) \leq P_\rho$ [3, Lemma 3.1] and so $RR_\rho \leq P_\rho$ and $R_\rho = R_{\rho+1} = RR_\rho \geq R$. Therefore S/R_ρ is a π -Sylowizer of the trivial subgroup in G_ρ/R_ρ and so is a Sylow π' -subgroup of G_ρ/R_ρ .

Since the Sylow π' -subgroups of a $\mathfrak{G}$ -group are locally conjugate and the Sylow π' -subgroups of a $\mathfrak{U}$ -group are conjugate we immediately have the following corollaries.

Corollary 1. *If R is a normal subgroup of a Sylow π -subgroup of a $\mathfrak{G}$ -group G , then the π -Sylowizers of R in G are locally conjugate in G .*

Corollary 2. *If R is a normal subgroup of a Sylow π -subgroup of a $\mathfrak{U}$ -group G , then the π -Sylowizers of R in G are conjugate in G .*

REFERENCES

1. A. D. Gardiner, B. Hartley and M. J. Tomkinson, *Saturated formations and Sylow structure in locally finite groups*, J. Algebra 17 (1971), 177–211. MR 42 #7778.
2. W. Gaschütz, *Sylowisatoren*, Math. Z. 122 (1971), 319–320.
3. B. Hartley, *Sylow theory in locally finite groups*, Compositio Math. 25 (1972), 263–280.
4. D. J. S. Robinson, *Finiteness conditions and generalized soluble groups*. I, Springer-Verlag, Berlin and New York, 1972.
5. M. J. Tomkinson, *Formations of locally soluble FC-groups*, Proc. London Math. Soc. (3) 19 (1969), 675–708. MR 41 #5501.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GLASGOW, G12 8QW SCOTLAND