A NOTE ON EXTREME ELEMENTS IN $A_0(K, E)$

M. SHARIR¹

ABSTRACT. We give a short and simple proof to a theorem of Fakhoury, characterizing extreme elements in the unit ball of $A_0(K, E)$.

Let V be a Banach space whose dual is an L^1 -space. Denote by K the unit ball of V^* equipped with the w^* -topology. Let E be a Banach space. $S(E)$ will denote its closed unit ball and $\text{ext } S(E)$ the set of extreme points in $S(E)$. $A_0(K, E)$ will be the Banach space of all the symmetric affine functions from K into E , continuous in the w^* -topology on K , and in the norm topology on E . Fakhoury has shown in [1] that if E has certain properties, then $f \in A_0(K, E)$ is extreme in $S(A_0(K, E))$ if and only if $f(\text{ext } K) \subset \text{ext } S(E)$. This result bears immediate characterization of extreme compact operators (if $E = F^*$, then $A_0(K, E)$ is precisely the space of compact operators from F into V), and generalizes similar results of Lazar [2]. By observing a simple property of spaces having the 3.2.I.P. (cf. [4] for a proper definition), and by using a selection theorem of Lazar and Lindenstrauss [3], we are able to prove Fakhoury's result in a very simple and direct way.

Lemma. *Let E be a Banach space having the 3.2.I.P. and let $x, y \in S(E)$, $a \in E$, such that $x \pm a \in S(E)$. Then there exists an element $b \in E$, such that $y \pm b \in S(E)$, and $\|b - a\| \leq \|y - x\|$.*

Proof. Define three closed balls in E :

$$S_1 = \{b \in E; \|b - y\| \leq 1\},$$

$$S_2 = \{b \in E; \|b + y\| \leq 1\},$$

$$S_3 = \{b \in E; \|b - a\| \leq \|y - x\|\}.$$

These balls intersect in pairs, for $0 \in S_1 \cap S_2$, $y + a - x \in S_1 \cap S_3$ and

---

Received by the editors August 15, 1973.

AMS (MOS) subject classifications (1970). Primary 47D20, 46E15, 46E30, 47B05.

¹ This paper is a part of the author's Ph.D. Thesis, prepared at Tel Aviv University, under the supervision of Dr. A. J. Lazar.

Copyright © 1974, American Mathematical Society

$-y + a + x \in S_2 \cap S_3$. Hence they have a nonempty intersection, and any $b \in S_1 \cap S_2 \cap S_3$ has all the desired properties.

Theorem. *Let the scalars be real, and let E be a Banach space with the 3.2.I.P. and let K be as above. Then an element ϕ of $A_0(K, E)$ is extreme in the unit ball of this space if and only if $\phi(\text{ext } K) \subset \text{ext } S(E)$.*

Proof. Since one direction is immediate, assume ϕ to be extreme. We make use of a selection theorem of Lazar and Lindenstrauss [3]. Define a set-valued mapping $\Sigma: K \rightarrow 2^{S(E)}$ by

$$\Sigma(\mu) = \{x \in S(E); \|\phi(\mu) \pm x\| \leq 1\}, \quad \mu \in K.$$

Now, for each $\mu \in K$, $\Sigma(\mu)$ is a norm-closed convex nonvoid subset of $S(E)$. Also $\Sigma(-\mu) = \Sigma(\mu) = -\Sigma(\mu)$ for each $\mu \in K$ (thus Σ is symmetric), and for each $\mu_1, \mu_2 \in K$, $0 \leq \alpha \leq 1$, we have

$$\alpha\Sigma(\mu_1) + (1 - \alpha)\Sigma(\mu_2) \subset \Sigma(\alpha\mu_1 + (1 - \alpha)\mu_2)$$

(thus Σ is convex). Σ is also lower semicontinuous (in the sense of Michael [5]) with the w^* -topology on K and the norm-topology on $S(E)$: Let $\mu \in K$, $x \in \Sigma(\mu)$, and $\mu_\alpha \xrightarrow{w^*} \mu$ in K . We have to show the existence of $x_\alpha \in \Sigma(\mu_\alpha)$, for each α , such that $x_\alpha \rightarrow x$ in norm. But this is immediate since $\phi(\mu_\alpha) \rightarrow \phi(\mu)$ in norm, and, for each α , the Lemma provides us with a $x_\alpha \in E$ such that $\|\phi(\mu_\alpha) \pm x_\alpha\| \leq 1$ (hence $x_\alpha \in \Sigma(\mu_\alpha)$) and $\|x_\alpha - x\| \leq \|\phi(\mu_\alpha) - \phi(\mu)\|$. Hence the lower semicontinuity of Σ is obvious. Now, assume that $\phi(\text{ext } K) \not\subset \text{ext } S(E)$. Hence, there is a $\mu_0 \in \text{ext } K$ and a nonzero element x_0 of $\Sigma(\mu_0)$. Now, $f(\alpha\mu_0) = \alpha x_0$, $\alpha \in [-1, 1]$, is a w^* -continuous affine symmetric selection of Σ restricted to $\text{conv}(\{\mu_0\} \cup -\{\mu_0\})$, for which $\{\mu_0\}$ is obviously essentially closed (cf. [3] for proper definitions). Hence, there is a w^* -continuous affine symmetric selection $\psi \in A_0(K, E)$ of Σ , such that $\psi(\mu_0) = x_0 \neq 0$. Hence, for each $\mu \in K$ one has $\|\phi(\mu) \pm \psi(\mu)\| \leq 1$ and therefore $\|\phi \pm \psi\| \leq 1$. Since ϕ is extreme we must have $\psi = 0$, a contradiction which completes the proof.

Remark. The estimate $\|x_\alpha - x\| \leq \|\phi(\mu_\alpha) - \phi(\mu)\|$, and therefore also the requirement that E will have the 3.2.I.P., is apparently too strong for the proof of the Theorem and weaker assumptions will do as well.

REFERENCES

1. H. Fakhoury, *Préduaux de L-espaces et éléments extrémaux*, C. R. Acad. Sci. Paris Sér. A-B 272 (1971), A1703-A1706. MR 43 #6700.

2. A. J. Lazar, *Affine functions on simplexes and extreme operators*, Israel J. Math. 5 (1967), 31–43. MR 35 #2128.
3. A. J. Lazar and J. Lindenstrauss, *Banach spaces whose duals are L_1 spaces and their representing matrices*, Acta Math. 126 (1971), 165–193. MR 45 #862.
4. J. Lindenstrauss, *Extension of compact operators*, Mem. Amer. Math. Soc. No. 48 (1964). MR 31 #3828.
5. E. Michael, *Continuous selections. I*, Ann. of Math. (2) 63 (1956), 361–382. MR 17, 990.

DEPARTMENT OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV,
ISRAEL