SPLITTING GROUPS BY INTEGERS

W. HAMAKER AND S. STEIN

ABSTRACT. A question concerning tiling Euclidean space by crosses raised this algebraic question: Let G be a finite abelian group and S a set of integers. When do there exist elements $g_1, g_2, \dots, g_n$ in G such that each nonzero element of G is uniquely expressible in the form sg_i for some s in S and some g_i ? The question is answered for a broad (but far from complete) range of S and G .

Let G be a finite abelian group and $S = \{s_1, s_2, \dots, s_k\}$ a set of k distinct integers. If there are elements $g_1, g_2, \dots, g_n$ in G such that each nonzero element of G is uniquely expressible in the form $s_i g_j$, $1 \leq i \leq k$, $1 \leq j \leq n$, we will say that S *splits* G . The set $\{g_1, \dots, g_n\}$ is a *splitting set*. The cases $S = \{1, 2, \dots, k\}$ and $S = \{\pm 1, \pm 2, \dots, \pm k\}$ arise in the case of tiling Euclidean space by certain starbodies (see [2] and [3]). In [1] it was shown that $S = \{1, 3, 27\}$ splits no finite abelian group. If each $s_i \in S$ is relatively prime to $|G|$, the order of G , we call the splitting *nonsingular*. Throughout we will assume that all groups have at least two elements.

Theorem 1. *Let S split the groups A and B such that the splitting of A is nonsingular. Let $0 \rightarrow A \xrightarrow{\alpha} G \xrightarrow{\beta} B \rightarrow 0$ be an exact sequence (kernel β = image α , α one-one, β onto). Then S splits G .*

Proof. Let $a_1, a_2, \dots, a_p$ be a splitting set in A , and $b_1, b_2, \dots, b_q$ be a splitting set in B . For each n , $1 \leq n \leq q$, select $g_n \in G$ such that $\beta(g_n) = b_n$. We assert that the set

$$(1) \quad \{\alpha(a_j): 1 \leq j \leq p\} \cup \{\alpha(a) + g_n: a \in A - \{0\}, 1 \leq n \leq q\}$$

is a splitting set for G .

To begin, let $U = \{a_1, \dots, a_p\}$ and $V = \{g_1, \dots, g_n\}$. Then we wish to show first that $G - \{0\} = S(\alpha(U) \cup (\alpha(U) + V))$. Noting that $SV \cup \{0\}$ is a complete set of coset representatives for $G \bmod \alpha(A)$, we have

Received by the editors November 29, 1973.

AMS (MOS) subject classifications (1970). Primary 20K05; Secondary 50B30.

Key words and phrases. Exact sequence, abelian group, splitting.

Copyright © 1974, American Mathematical Society

$$G - \{0\} = (\alpha(A) - \{0\}) \cup (\alpha(A) + SV) = S(\alpha(U)) \cup (\alpha A + SV).$$

Now, since S splits A in a nonsingular manner, $SA = A$, hence $S(\alpha A) = \alpha A$. Also, for any sets X and Y in G such that $SX = X$, we have $X + SY = SX + SY$. Thus

$$\begin{aligned} G - \{0\} &= S(\alpha(U)) \cup (\alpha A + SV) = S(\alpha(U)) \cup S(\alpha A + V) \\ &= S(\alpha(U) \cup (\alpha A + V)), \end{aligned}$$

as desired.

The uniqueness of the representation follows from the fact that the set (1) contains $(|A| - 1)/k + (|B| - 1)/k + (|A| - 1)(|B| - 1)/k$ or $(|A| |B| - 1)/k = (|G| - 1)/k$ elements.

The following two corollaries are of special interest.

Corollary 1. *Let S split the groups A and B in such a way that the splitting of A is nonsingular. Then S splits the product $A \times B$.*

In the next corollary $C(q)$ denotes the cyclic group of order q .

Corollary 2. *Let S split $C(q)$ in a nonsingular manner. Then S splits $C(q^n)$ for each positive integer n .*

Proof. There is an exact sequence $0 \rightarrow C(q) \rightarrow C(q^{n+1}) \rightarrow C(q^n) \rightarrow 0$. Theorem 1, combined with an induction on n , establishes the Corollary.

The two corollaries, together with the fundamental theorem of abelian groups, yield the following theorem.

Theorem 2. *Let p be an odd prime integer. Let $S = \{1, 2, \dots, p-1\}$ or $\{\pm 1, \pm 2, \dots, \pm(p-1)/2\}$. Then S splits any abelian group whose order is a power of p .*

[As [2] or [3] show, Theorem 2 implies, for example, that a $(p-1)/2$ cross tiles Euclidean n space if $n(p-1) + 1$ is a power of p , say p^b , in at least as many geometrically inequivalent ways as there are nonisomorphic abelian groups of order p^b .]

The next theorem is sort of a converse to Theorem 1.

Theorem 3. *Let $0 \rightarrow A \xrightarrow{\alpha} G \xrightarrow{\beta} B \rightarrow 0$ be an exact sequence of groups. Assume that S splits G . If each s_i in S is relatively prime to $|B|$, then S splits A . If each s_i in S is relatively prime to $|A|$, then S splits B .*

Proof. We begin by proving the first assertion. Let $T = \{g_1, g_2, \dots, g_n\}$ be the splitting set of G . We assert that $\alpha(A) \cap T$ is a splitting set for $\alpha(A)$. (Hence $\alpha^{-1}(\alpha(A) \cap T)$ would be a splitting set for A .)

We establish first that

$$\alpha(A) - \{0\} = S(\alpha(A) \cap T),$$

or, equivalently,

$$\alpha(A) \cap ST = S(\alpha(A) \cap T).$$

Clearly, $\alpha(A) \cap ST \supseteq S(\alpha(A) \cap T)$. To show that $S(\alpha(A) \cap T) \subseteq \alpha(A) \cap ST$, note that for $s \in S$ and $t \in T$

$$st \in \alpha(A) \Rightarrow s\beta(t) = \beta(st) = 0 \Rightarrow \beta(t) = 0,$$

since $(s, |B|) = 1$. Hence $t \in \alpha(A)$ and $\alpha(A) \cap ST \subseteq S(\alpha(A) \cap T)$.

Thus every element of $\alpha(A) - \{0\}$ is representable in the form sg where $s \in S$ and $g \in \alpha(A) \cap T$. Showing that this representation is unique is straightforward.

The second assertion in the theorem is an immediate consequence of the fact that a homomorphic image of G is isomorphic to a subgroup of G . This reduces the second case to the first. This concludes the proof.

The assumption in Theorem 3 that $(s_i, |B|) = 1$ cannot be removed. To see this, consider $S = \{1, 2, 3\}$ and an exact sequence $0 \rightarrow C(2) \rightarrow C(4) \rightarrow C(2) \rightarrow 0$.

Theorem 3, combined with Corollaries 1 and 2, yield the following reduction of the problem of determining all nonsingular splittings.

Theorem 4. *Let G be a finite abelian group and $S = \{s_1, s_2, \dots, s_k\}$ a set of integers with each s_i relatively prime to $|G|$. Then S splits G if and only S splits $C(p)$ for each prime p that divides $|G|$.*

REFERENCES

1. W. Hamaker, *Factoring groups and tiling space*, *Aequationes Math.* **9** (1973), 145–149.
2. S. K. Stein, *Factoring by subsets*, *Pacific J. Math.* **22** (1967), 523–541. MR 36 #2517.
3. ———, *A symmetric star body that tiles but not as a lattice*, *Proc. Amer. Math. Soc.* **36** (1972), 543–548.

DEPARTMENT OF MATHEMATICS, DREXEL UNIVERSITY, PHILADELPHIA, PENNSYLVANIA 19104

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA 95616