AN EXTENSION OF RUNGE'S THEOREM

L. A. RUBEL¹

ABSTRACT. Every uniformly continuously differentiable function on the compact set X in the complex plane is the uniform limit of rational functions with poles off X .

The usual hypotheses of Runge's theorem demand that the function f , to be approximated, have an analytic extension to a neighborhood of the set X on which the approximation is to take place. Our result only places conditions on f on X itself, and implies the usual form of Runge's theorem.

Definition. On a compact set X in the complex plane $\mathbb{C}$, a continuous complex-valued function f is said to be continuously uniformly differentiable (written $f \in D^{UC}(X)$) if there is a continuous function f' on X s.t. for every $\epsilon > 0$ there exists a $\delta > 0$ s.t. $\forall z \in X$ and $\forall w, w' \in X$, $w \neq w'$, with $|w - z| < \delta$, $|w' - z| < \delta$, we have

$$(1) \quad \left| \frac{f(w') - f(w)}{w' - w} - f'(z) \right| < \epsilon.$$

Let $R(X)$ denote the uniform closure on X of the rational functions with poles off X .

Theorem. $D^{UC}(X) \subseteq R(X)$.

Proof. We first observe that from (1), it follows by the Whitney extension theorem [W, p. 65] that f has a C^1 extension (still called f) to all of $\mathbb{C}$. Next, it follows from (1) by an elementary argument that if z_0 is a point of X for which there exist two sequences $\{z_n\}$ and $\{z'_n\}$ of points of X that converge to z_0 along two different lines through z_0 , then the Cauchy-Riemann conditions hold at z_0 . Now we remark that z_0 satisfies the above condition if it is a point of density of X with respect to planar Lebesgue measure $dx dy$. For if we let $E_n = \{\theta: z_0 \pm re^{i\theta} \notin X, 0 < r < 2^{-n}\}$, then all E_n must have measure zero at a point of density z_0 . But almost every point of

Received by the editors April 22, 1974.

AMS (MOS) subject classifications (1970). Primary 30A82, 41A20.

¹ This research was supported by a grant from the National Science Foundation.

X is a point of density of X . Thus, we have proved that if $f \in D^{UC}(X)$ it has a C^1 extension to $\mathbb{C}$ that satisfies the Cauchy-Riemann equations at almost all points of X . Finally, by a slight variation of the proof of [B, Corollary 3.22, p. 160], we conclude that $f \in R(X)$. Indeed, if we take $\mu \perp R(X)$ and let $\hat{\mu}$ be its Cauchy transform we have

$$\int f d\mu = -\frac{1}{\pi} \iint_X \frac{\partial f}{\partial \bar{z}} \hat{\mu}(z) dx dy.$$

But $\partial f / \partial \bar{z} = 0$ a.e. on X and $\hat{\mu}(z) = 0$ off X so that $\mu \perp f$ and, hence, $f \in R(X)$ by the Hahn-Banach theorem.

REFERENCES

- [B] Andrew Browder, *Introduction to function algebras*, Benjamin, New York, 1969. MR 39 #7431.
 [W] Hassler Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. 36 (1934), 63–89.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, URBANA, ILLINOIS 61801