HELSON SETS WHICH DISOBEY SPECTRAL SYNTHESIS

SADAHIRO SAEKI

ABSTRACT. In this paper it is shown that every nondiscrete LCA group contains a compact independent Helson set which disobeys spectral synthesis.

In [3] T. W. Körner has constructed an independent compact H_1 -set of type M . This result solves negatively the long-standing problem whether or not every Helson set obeys spectral synthesis. Körner's construction of such a set is, however, very complicated, and R. Kaufman [2] has simplified it (see also [6]). In this paper, we modify Kaufman's method to prove that every nondiscrete metrizable LCA group contains an independent compact Helson set of type M . Consequently it is shown that every nondiscrete LCA group contains a Helson set which disobeys spectral synthesis.

Let G be a LCA group with dual $\hat{G}$. We denote by $A(G)$ and $PM(G)$ the Fourier algebra on G and the conjugate space of $A(G)$, respectively. Each element of $PM(G)$ is called a pseudo-measure on G . For $f \in A(G)$ and $P \in PM(G)$, we define

$$\langle f, P \rangle = (\hat{f} * \hat{P})(1) = \int_{\hat{G}} \hat{f}(\chi) \hat{P}(\chi^{-1}) d\chi,$$

where $\hat{f}$ and $\hat{P}$ denote the functions in $L^1(\hat{G})$ and in $L^\infty(\hat{G})$ whose (inverse) Fourier transforms are f and P , respectively. If $\mu \in M(\hat{G})$ and $\check{\mu}(x) = \int_{\hat{G}} \chi(x) d\mu(\chi)$ ($x \in G$), then we denote by $\check{\mu}P$ the pseudo-measure on G defined by the requirement $(\check{\mu}P)^\wedge = \mu * \hat{P}$. It is well known that if $P \in PM(G)$ has compact support, then $\hat{P}$ can be chosen from the space $C(\hat{G})$; we will always do this. For such a P , $\langle \check{\mu}, P \rangle = (\mu * \hat{P})(1)$ ($\mu \in M(\hat{G})$) is well defined, and we have $\langle \chi, \gamma P \rangle = \hat{P}(\chi^{-1} \gamma^{-1})$ ($\chi, \gamma \in \hat{G}$). A pseudo-function on G is any pseudo-measure whose Fourier transform is a continuous function on $\hat{G}$ which vanishes at infinity. The space of all pseudo-functions is denoted by $PF(G)$.

Received by the editors July 25, 1973.

AMS (MOS) subject classifications (1970). Primary 43A45; Secondary 43A20.

Key words and phrases. LCA group, Helson set, set of type M , H_1 -set, K_q -set, Fourier algebra, spectral synthesis, pseudo-function.

Let now E be a closed set in G , and $q = q(G)$ the supremum of the natural numbers n such that every neighborhood of 0_G contains an element with order $\geq n$. E is called an M -set (or a set of type M) if $PF(E) \neq 0$ (that is, if E carries a nonzero pseudo-function); strongly independent if E is independent in the usual sense and $\text{ord}(e) = q$ for all $e \in E$; and an H_1 -set if $\|\mu\|_M = \|\mu\|_{PM}$ for all $\mu \in M(E)$. In the case $q < \infty$, E is called a K_q -set if every function $f \in C(E)$ with $f^q = 1$ is the restriction of a character in $\hat{G}$; and a weak K_q -set if to each $\mu \in M^+(E)$ and $\epsilon > 0$ there corresponds a K_q -set $K \subset E$ such that $\mu(E \setminus K) < \epsilon$.

Our main result now follows (cf. [3, p. 105]).

Theorem 1. *Every nondiscrete LCA group contains a strongly independent compact set which is either an H_1 -set or a weak K_q -set but disobeys spectral synthesis. If, in addition, the group under consideration is metrizable, such a set can be chosen as a set of type M .*

The proof becomes clear after some lemmas and theorems are established.

Lemma 1. *To each $0 < \epsilon < 1$ there corresponds a natural number N_ϵ with the following property: for any neighborhood V of 0 of a compact abelian group H and any natural number $N \geq N_\epsilon$, there exists an $F \in A(H^N)$ such that*

- (i) $\text{supp } F \subset V_\epsilon^N$, and
- (ii) $|\hat{F}(k)| < \epsilon |\hat{F}(0)| \quad \forall k \in \hat{H}^N \setminus \{0\}$.

Here V_ϵ^N is the set of all $(x_j)_{j=1}^N \in H^N$ such that $x_j \in V$ for at least $(1 - \epsilon)N$ indices $j = 1, 2, \dots, N$, and $\hat{H}^N$ denotes the additively written dual of H^N .

The proof is essentially identical with that of [2, Lemma 2], so we omit it.

Lemma 2. *Let G be a LCA I-group, $\hat{K}$ a compact subset of $\hat{G}$, and $M, N \in \mathbb{N}$ (the natural numbers). Then there exist N characters $\chi_1, \dots, \chi_N \in \hat{G}$ such that the sets $\chi_1^{k_1} \cdots \chi_N^{k_N} \hat{K}$ ($k_j \in \{0, \pm 1, \dots, \pm M\}$, $1 \leq j \leq N$) are pairwise disjoint.*

Proof. By the structure theorem [1, (9.8)], G contains an open subgroup G_0 which is topologically isomorphic to $R^a \times H$ for some $a \in \{0, 1, 2, \dots\}$ and some compact abelian group H . Since there exists a continuous homomorphism of $\hat{G}$ onto $\hat{G}_0$, we may assume that $G = G_0 = R^a \times H$.

Suppose that there exists a $\gamma \in \hat{G}$ such that

$$(1) \quad \lim_n \gamma^n = \infty.$$

Take $p \in \mathbb{N}$ so large that $n > p$ $\gamma^n \notin \hat{K}^{-1}\hat{K}$, and choose N natural numbers $q_1, \dots, q_N$ so that $|k_1 q_1 + \dots + k_N q_N| > p$ for all choices of $k_j \in \{0, \pm 1, \pm 2, \dots, \pm 2M\}$, $1 \leq j \leq N$, such that $(k_1, \dots, k_N) \neq (0, \dots, 0)$. Then the elements $\chi_j = \gamma^{q_j}$, $1 \leq j \leq N$, have the required property.

Assume now that no $\gamma \in \hat{G}$ satisfies (1). Then $a = 0$, $G = H$ is compact and $\hat{G}$ is a torsion group. Therefore every finitely-generated subgroup of $\hat{G}$ is finite. But $\hat{G}$ is not of bounded order because G is an I-group (see [4, 2.5.4]). Thus, setting $\chi_0 = 1$, we can find $\chi_1, \dots, \chi_N \in \hat{G}$ so that

$$\text{ord}(\chi_j) > 2M \cdot \text{Card}[G_p(\hat{K} \cup \{\chi_0, \dots, \chi_{j-1}\})]$$

for all $j = 1, 2, \dots, N$. As is easily seen, the χ_j , $1 \leq j \leq N$, have the required property.

Lemma 3. *Let G be a LCA I-group, and τ a pseudo-function on G whose support E is compact. Let also $0 < \epsilon < 1$, and g any function in $C(E)$ whose range is a finite subset of $T = \{z: |z| = 1\}$. Then there exist $\tau' \in PF(G)$ and $\chi_j \in \hat{G}$, $1 \leq j \leq N$, such that*

- (a) $\|\tau' - \tau\|_{PM} < \epsilon$,
- (b) $\text{supp } \tau' \subset \text{supp } \tau$,
- (c) $\|g - (1/N) \sum_{j=1}^N \chi_j\|_{C(\text{supp } \tau')} < \epsilon$.

Proof. We may assume that g is defined and continuous on some compact neighborhood U of E , and that $g(U)$ is a finite subset of T . Thus

$$(1) \quad C = \sup \{\|g^k\|_{A(U)}: k \in \mathbb{Z}\} < \infty,$$

where

$$\|g^k\|_{A(U)} = \inf \{\|f\|_{A(G)}: f \in A(G), f = g^k \text{ on } U\}.$$

Put $V = \{z \in T: |z - 1| < \epsilon\}$, so that

$$(2) \quad V_\epsilon^N = \{(z_j)_1^N \in T^N: \text{Card}\{j: |z_j - 1| < \epsilon\} \geq (1 - \epsilon)N\}$$

for all $N \in \mathbb{N}$. We apply Lemma 1 to find an $N \in \mathbb{N}$ and an $F \in A(T^N)$ such that

$$(3) \quad \text{supp } F \subset V_\epsilon^N,$$

$$(4) \quad \hat{F}(0) = 1 \quad \text{and} \quad |\hat{F}(k)| < \epsilon \quad \forall k \in \mathbb{Z}^N \setminus \{0\}.$$

Choose a finite set $L \subset \mathbf{Z}^N$ so that

$$(5) \quad \sum_{k \notin L} |\hat{F}(k)| \cdot C \|\tau\|_{PM} < \epsilon,$$

and put

$$(6) \quad \hat{K} = \bigcup_{k \in L} \{\chi \in \hat{G}: |(g^{(k)}\tau)^\wedge(\chi)| \geq \epsilon/\text{Card } L\},$$

where $(k) = \sum_j k_j$ for $k = (k_1, \dots, k_N) \in \mathbf{Z}^N$. Since $g^{(k)}\tau \in PF(G)$ and L is finite, $\hat{K}$ is compact. It follows from Lemma 2 that there are N characters $\chi_1, \dots, \chi_N \in \hat{G}$ such that the sets

$$(7) \quad \chi_1^{-k_1} \dots \chi_N^{-k_N} \hat{K} \quad (k \in L)$$

are pairwise disjoint. Note that the series

$$\sum_{k \in \mathbf{Z}^N} \hat{F}(k) g^{(k)} \bar{\chi}_1^{k_1} \dots \bar{\chi}_N^{k_N}$$

converges to $F(g\bar{\chi}_1, \dots, g\bar{\chi}_N) \in A(U)$ in the norm of $A(U)$ by (1). Setting

$$(8) \quad \tau' = F(g\bar{\chi}_1, \dots, g\bar{\chi}_N)\tau = \sum_{k \in \mathbf{Z}^k} \hat{F}(k) g^{(k)} \bar{\chi}_1^{k_1} \dots \bar{\chi}_N^{k_N} \tau,$$

we claim that τ' and $\chi_1, \dots, \chi_N$ have the required properties if ϵ is replaced by $\epsilon/(C\|\tau\|_{PM} + 3)$.

To prove part (a), put

$$\nu = \sum_{0 \neq k \in L} \hat{F}(k) g^{(k)} \bar{\chi}_1^{k_1} \dots \bar{\chi}_N^{k_N} \tau.$$

Then we have

$$\begin{aligned} |\hat{\nu}(\chi)| &\leq \sum_{0 \neq k \in L} |\hat{F}(k)| \cdot |(g^{(k)}\tau)^\wedge(\chi_1^{k_1} \dots \chi_N^{k_N} \chi)| \\ &\leq \epsilon \sum_{k \in L} |(g^{(k)}\tau)^\wedge(\chi_1^{k_1} \dots \chi_N^{k_N} \chi)| \end{aligned}$$

for all $\chi \in \hat{G}$ by (4). If one of the summands in the last sum is $\geq \epsilon/\text{Card } L$, then the other summands are $< \epsilon/\text{Card } L$ by (6) and (7). Therefore, $|\hat{\nu}(\chi)| \leq \epsilon(C\|\tau\|_{PM} + 1)$ by (1); hence $\|\nu\|_{PM} \leq \epsilon(C\|\tau\|_{PM} + 1)$. It follows from (8), (4), and (5) that

$$\begin{aligned}
 \|\tau' - \tau\|_{PM} &= \left\| \sum_{k \neq 0} \hat{F}(k) g^{(k)} \overline{\chi_1}^{k_1} \cdots \overline{\chi_N}^{k_N} \tau \right\|_{PM} \\
 &\leq \|\nu\|_{PM} + \sum_{k \notin L} |\hat{F}(k)| \cdot \|g^{(k)} \tau\|_{PM} \\
 &< \epsilon(C\|\tau\|_{PM} + 1) + \epsilon = \epsilon(C\|\tau\|_{PM} + 2).
 \end{aligned}$$

Part (b) is trivial from (8).

To verify (c), note that $\text{supp } \tau'$ is contained in the closed set $\{x \in E: \text{Card}\{j: |g(x) - \chi_j(x)| \leq \epsilon\} \geq (1 - \epsilon)N\}$, which is easily seen from (2), (3), and (8). Thus $x \in \text{supp } \tau'$ implies

$$\left| g(x) - N^{-1} \sum_{j=1}^N \chi_j(x) \right| \leq N^{-1} \sum_{j=1}^N |g(x) - \chi_j(x)| \leq \epsilon + N^{-1} \cdot 2\epsilon N = 3\epsilon.$$

This completes the proof.

Theorem 2. *Let G be a LCA I-group and τ a pseudo-function on G whose support E is totally disconnected and metrizable. Then for each $\epsilon > 0$ there exists a $\nu \in PF(E)$, with $\|\nu - \tau\|_{PM} < \epsilon$, whose support is a strongly independent H_1 -set.*

Proof. Since $\hat{\tau}$ is uniformly continuous, we may assume that E is compact. Since E is metrizable, $C(E)$ is separable, and so $\{f \in C(E): |f| = 1\}$ contains a countable dense set $\{g_n\}$. We may assume $\text{Card}[g_n(E)] < \infty$ for all n , because E is totally disconnected.

Let $\epsilon > 0$ be given, and set $\tau_0 = \tau$. Suppose that $\tau_0, \dots, \tau_{n-1} \in PF(E)$ are constructed for some natural number n . We apply Lemma 3 to find a $\tau_n \in PF(E)$ which satisfies (a), (b), and (c) in Lemma 3 with τ, ϵ, g replaced by $\tau_{n-1}, \epsilon/2^n, g_n$, respectively. The sequence $\{\tau_n\}_1^\infty$ obtained in this way converges to some $\nu \in PF(E)$ and we have

$$\|\nu - \tau\|_{PM} \leq \sum_{n=1}^{\infty} \|\tau_n - \tau_{n-1}\|_{PM} < \epsilon.$$

Clearly $\text{supp } \nu \subset \bigcap_n \text{supp } \tau_n$. Thus, for each $n \in \mathbb{N}$, there are finitely many characters $\chi_1, \dots, \chi_N \in \hat{G}$ such that

$$\left\| g_n - \frac{1}{N} \sum_{j=1}^N \chi_j \right\|_{C(\text{supp } \nu)} < \frac{\epsilon}{2^n}.$$

This implies that $\text{supp } \nu$ is an H_1 -set which is independent over Z .

Lemma 2'. *Let G be a nondiscrete LCA group with $q = q(G) < \infty$, $E \subset G$ and $\hat{K} \subset \hat{G}$ compact, and $N \in \mathbb{N}$. Then there are N characters $\chi_1, \dots, \chi_N \in \hat{G}$ with the following two properties:*

- (i) *The sets $\chi_1^{k_1} \cdots \chi_N^{k_N} \hat{K}$ ($k_j \in \{0, 1, \dots, q-1\}$, $1 \leq j \leq N$) are pairwise disjoint;*
- (ii) *$\chi_j^q = 1$ on some neighborhood of E ($1 \leq j \leq N$).*

Proof. We may assume that G is compactly generated. Thus $G = R^a \times Z^b \times H$ for some nonnegative integers a and b and some compact abelian group H (see [1, (9.8)]). Since $q < \infty$, $a = 0$ and H is of bounded order. Let $\hat{K}_0$ be the natural projection of $\hat{K}$ into $\hat{H}$. Since H is a compact group with $q(H) = q(G) = q < \infty$, we can find N characters $\gamma_1, \dots, \gamma_N \in \hat{H}$ with order q so that the sets $\gamma_1^{k_1} \cdots \gamma_N^{k_N} \hat{K}_0$ ($k_j \in \{0, 1, \dots, q-1\}$, $1 \leq j \leq N$) are pairwise disjoint. (Note that $\hat{H}$ is a weak direct product of finite cyclic groups [4, B8].) Setting $\chi_j = 1 \otimes \gamma_j \in T^b \times \hat{H} = \hat{G}$, we see that the elements $\chi_1, \dots, \chi_N$ have the required property.

Lemma 3'. *Let G be a nondiscrete LCA group with $q < \infty$, and τ a pseudo-function on G with compact support E . Let also $0 < \epsilon < 1$, and g any function in $C(E)$ with $g^q = 1$. Then there exists a $\tau' \in PF(G)$ such that*

- (a) $\|\tau' - \tau\|_{PM} < \epsilon$;
- (b) $\text{supp } \tau' \subset \text{supp } \tau$;
- (c) $\|g - (1/N) \sum_{j=1}^N \chi_j\|_{C(\text{supp } \tau')} < \epsilon$

for some characters $\chi_1, \dots, \chi_N \in \hat{G}$ with $\chi_j^q = 1$ on E .

The proof is very similar to that of Lemma 3. The needed modifications are as follows. Replace T and V by $T_q = \{z \in T: z^q = 1\}$ and $\{1\} \subset T_q$, respectively; realize the dual of T_q^N as $\{0, 1, \dots, q-1\}^N$ in the usual way; put $L = \{0, 1, \dots, q-1\}^N$; and use Lemma 2' instead of Lemma 2.

The following theorem can be proved by applying Lemma 3' just as Theorem 2 was proved. We omit the proof.

Theorem 3. *Let G be a nondiscrete LCA group with $q = q(G) < \infty$, and τ a pseudo-function on G whose support E is metrizable. Then, given $\epsilon > 0$, we can find a pseudo-function $\nu \in PF(E)$, with $\|\nu - \tau\|_{PM} < \epsilon$, whose support K has the following property: to each $\delta > 0$ and $g \in C(K)$ with $g^q = 1$, there correspond finitely many characters $\chi_1, \dots, \chi_N \in \hat{G}$ such that $\chi_j^q = 1$ on K ($1 \leq j \leq N$) and $\|g - (1/N) \sum_{j=1}^N \chi_j\|_{C(K)} < \delta$.*

It is now a routine matter to derive Theorem 1 from Theorems 2 and 3 because every LCA group G contains a closed metrizable group H with $q(H) = q(G)$, and every Helson set of type M disobeys spectral synthesis (cf. [4, 5.6.10]).

Remarks. (a) For a characterization of Helson sets of (spectral) synthesis, we refer to [5].

(b) The totally disconnectedness assumption on E in Theorem 2 is unnecessary. Let E be a closed metrizable subset of G , and $\tau \in PF(E)$. Then there exists a sequence $\{\tau_n\}$ in $PF(E)$ such that $\text{supp } \tau_n$ is totally disconnected and $\|\tau_n - \tau\|_{PM} = o(1)$. We omit the proof.

REFERENCES

1. E. Hewitt and K. A. Ross, *Abstract harmonic analysis*. Vol. I: *Structure of topological groups. Integration theory, group representations*, Die Grundlehren der math. Wissenschaften, Band 115, Academic Press, New York; Springer-Verlag, Berlin, 1963. MR 28 #158.
2. R. Kaufman, *M-sets and distributions*, *Astérisque Math. France* 5 (1973), 225–230.
3. T. W. Kömer, *A pseudo-function on a Helson set*, *Astérisque Math. France* 5 (1973), 3–224.
4. W. Rudin, *Fourier analysis on groups*, Interscience Tracts in Pure and Appl. Math., no. 12, Interscience, New York, 1962. MR 27 #2808.
5. S. Saeki, *A characterization of SH-sets*, *Proc. Amer. Math. Soc.* 30 (1971), 497–503. MR 44 #731.
6. T. W. Körner, *A pseudo-function on a Helson set. II*, *Astérisque Math. France* 5 (1973), 231–239.

DEPARTMENT OF MATHEMATICS, KANSAS STATE UNIVERSITY, MANHATTAN, KANSAS 66506

Current address: Department of Mathematics, Tokyo Metropolitan University, Setagaya, Tokyo, Japan