TRANSFORMATION GROUPS RESEMBLING THE ADJOINT REPRESENTATION

R. W. SULLIVAN¹

ABSTRACT. If G is a compact, connected Lie group, the isotropy subgroups of the adjoint representation of G are connected and the dimension of the fixed point set of a maximal torus of G is equal to the rank of G . Results similar to these are given when G acts differentiably on an integral cohomology sphere and has the adjoint representation as weak linear model. This is done by analyzing an induced action of the Weyl group of G .

1. Let G be a compact, connected Lie group. G acts on itself by conjugation, fixing the identity element and thereby giving rise to an orthogonal representation of G on the tangent space at the identity. This representation, called the *adjoint representation*, is completely described by its restriction to a maximal torus T of G , and this restriction is in turn completely described by the roots of G which are considered as vectors in the Lie algebra of T . The Lie algebra of T is contained in the Lie algebra of G ; since it is fixed by T it is acted on by $NT/T = W$, where NT is the normalizer of T in G . The finite group W , called the *Weyl group* of G , permutes the roots of G so that if α is a root, so is $\sigma\alpha$ for any σ in W . W is generated by the orthogonal reflections through the hyperplanes perpendicular to the (non-zero) roots. If S_α is the reflection through the plane perpendicular to α and σ is an element of W , $\sigma S_\alpha \sigma^{-1}$ is reflection through the root $\sigma\alpha$. Every reflection in W is of the form S_β for some nonzero root β [2, p. 74]. $W(G)$ will also be used to denote the Weyl group of G .

Let $F(T)$ denote the fixed point set of T ; G_x the subgroup of elements of G which fix x (called the *isotropy subgroup*); G_x^0 the connected component of G_x which contains the identity. If g is an element of G which fixes x but does not lie in G_x^0 , then $g^{-1}Tg = k^{-1}Tk$ for some k in G_x^0 since all maximal tori of G_x^0 are conjugate, so gk^{-1} is an element of $W(G)_x$ when x is fixed by T . If α is a nonzero root and H_α is the plane fixed by S_α , the centralizer in G of the corank one subtorus of T corresponding to H_α

Received by the editors January 21, 1974.

AMS (MOS) subject classifications (1970). Primary 57E15.

Key words and phrases. Weyl group, adjoint representation, cohomology sphere.

¹ Partially supported by National Science Foundation Grant GU 3171.

Copyright © 1975, American Mathematical Society

is denoted by G_α . Since G_α is connected [3, pp. 26–31] and $W(G_\alpha)$ is generated by S_α , any point in $F(T)$ fixed by S_α is also fixed by G_α . Since $W(G)_x$ is generated by reflections which fix x , we have

Proposition 1. (a) If $x \in F(T)$ is fixed by S_α , $G_\alpha \subset G_x$.

(b) The isotropy subgroups of the adjoint representation are connected.

(c) For $x \in F(T)$, $W(G)_x = W(G_x)$.

2. We want to generalize these observations to differentiable actions on cohomology spheres. If M is a differentiable manifold, a *diffeomorphism reflection* on M is a diffeomorphism r such that (a) r reverses orientation and $r^2 = 1$; (b) the complement of the fixed point set of r is disconnected. We shall refer to diffeomorphism reflections simply as *reflections*. The fixed point set of a reflection on an integral cohomology sphere HS^n is a Z_2 cohomology sphere by Smith theory. However, a standard argument [6, p. 35] shows that in fact it is an integral cohomology sphere HS^{n-1} .

If W is a group which is the Weyl group of a semisimple Lie group G , the *rank* of W is defined as the rank of G (i.e. the dimension of a maximal torus of G). For such a group W we consider differentiable actions (W, ϕ, HS^k) . Such an action is called *effective* if no element of W fixes every element of HS^k . Such an action is called *normal* if (a) each linear reflection in W is a reflection on HS^k ; (b) no product of two distinct linear reflections in W acts trivially. Considering ϕ as a homomorphism of W into the group of diffeomorphisms of HS^k , we say that a reflection on HS^k is *proper* if it is the image under ϕ of a linear reflection in W .

Theorem 2. If (W, ϕ, HS^k) ($k \geq 1$) is a normal action, then it is effective and each reflection is proper.

Proof. The proof is by induction on the rank of W . If $\text{rank } W = 1$, the theorem follows trivially from the definition. Let $\text{rank } W = n > 1$ and suppose that the theorem is true for $\text{rank } W$ less than n . If S is a proper reflection there is a point x in HS^k which is fixed by S but which is not fixed by any other proper reflection since the proper reflections have distinct fixed point sets (which meet transversally). If t is any other element of W fixing such an x , then $t^{-1}St = S$. Hence any element of W acting trivially must lie in the center of W . If w is such an element, the eigenvalues of w are ± 1 ; but if w takes $+1$ as an eigenvalue, then w lies in a Weyl subgroup of W of less rank by Proposition 1(c). So we may assume that $w = -1$.

Lemma. The element -1 in W can be written as the product of n mutually commuting distinct linear reflections where $n = \text{rank } W$.

Proof of Lemma. It is sufficient to prove this when W is the Weyl group of a simple Lie group. If $W = W(E_6)$, $W(A_q)$, $q \neq 1$, or $W(D_p)$, p odd, then $-I$ is not an element of W [2, p. 284]. For the remaining classical groups it is obvious, and since $W(B_4)$ is contained in $W(F_4)$ and $W(D_8)$ is contained in $W(E_8)$, we are left with G_2 and E_7 . We use the notation of [2, pp. 250–275]. For G_2 reflect through roots in the set $\{\epsilon_1, \epsilon_2\}$. For E_7 use roots in the set $\{\epsilon_1 - \epsilon_2, \epsilon_1 + \epsilon_2, \epsilon_3 - \epsilon_4, \epsilon_3 + \epsilon_4, \epsilon_5 - \epsilon_6, \epsilon_5 + \epsilon_6, \epsilon_7 - \epsilon_8\}$.

Returning to the proof of the theorem, we have $w = R_1 R_2 \cdots R_n$ acting trivially, $R_i R_j = R_j R_i$. Since each R_i reverses orientation, n must be even. But each R_i acts on the fixed point set of R_n , $F(R_n)$, and must reverse its orientation (for $i \neq n$) since it does so locally near a fixed point. Hence the case n even is also impossible.

If s in W is an improper reflection we have essentially the same situation. Since $s^2 = I$, either s lies in a Weyl subgroup of smaller rank or $s = -I$, so we may write $s = R_1 R_2 \cdots R_n$ in accordance with the Lemma above. Each R_i acts on $F(s)$ and reverses the orientation (or interchanges the two fixed points when $k = 1$). Since the product acts trivially on $F(s)$ we are done.

3. Let (G, ϕ, HS^n) be a differentiable action of a compact connected Lie group G on an integral cohomology sphere HS^n . Let $S(\phi)$ be the cw -system of ϕ [8]. $S(\phi)$ is defined as a collection of linear functionals on the Lie algebra of T which vanish on the Lie algebras of the connected co-rank one subtori of T which have fixed point sets of greater dimension than that of T . $S(\phi)$ includes the zero functional with multiplicity $dF(T) + 1$, where $dF(T)$ denotes the dimension of the fixed point set of T ; $S'(\phi)$ denotes the collection of nonzero elements in $S(\phi)$. We assume for the remainder of this section that $S'(\phi) = S'(\text{Ad}_G)$; the elements of this set then vanish on the hyperplanes perpendicular to the roots of G discussed in §1. The fixed point set of T is an integral cohomology sphere acted upon by $W(G)$. The fact that this is a normal action is proved in [8, 1.11] and will not be repeated here. It is an easy consequence of a theorem of W.-Y. Hsiang which states that Proposition 1(a) holds for differentiable actions (cf. [5, Proposition 3, p. 349]).

Since $W(G)_x$ is generated by the reflections which fix x [4, Theorem 1] and since these are now known to be proper reflections, it follows, just as in the proof of Proposition 1, that the isotropy subgroups are connected and that $W(G)_x = W(G_x)$ for x in $F(T)$. That T is a principal isotropy subgroup of ϕ now follows from [7, Theorem 3.3]. Since the $W(G)$ action is now

known to be effective, it follows from Theorem 3 of [4] that $dF(T) \geq \text{rank } G - 1$. Thus we have

Theorem 3. *Let (G, ϕ, HS^n) be a differentiable action of a compact connected Lie group with $S'(\phi) = S'(\text{Ad}_G)$. Then:*

- (a) $dF(T) \geq \text{rank}(G) - 1$;
- (b) *All isotropy subgroups are connected*;
- (c) *For x in $F(T)$, $W(G)_x = W(G_x)$.*
- (d) $S(\phi) = S(\text{Ad}_G + r \text{ trivial copies})$, where $r = dF(T) - \text{rank } G + 1 = dF(G) + 1$.

Remark. Part of the proof of Theorem 3 of [4] has not yet appeared, namely the proof of Theorem 2'. In lieu of this, one can produce a weaker inequality in (a) above as follows: $W(G)$ contains subgroups isomorphic to $Z_2 \times \cdots \times Z_2 = (Z_2)^k$, where each copy of Z_2 is generated by a reflection in $W(G)$. If $-I$ is an element of $W(G)$, the Lemma shows that $k = \text{rank } G$; otherwise k is at least $\frac{1}{2} \text{rank } G$. One can apply the formula of Borel [1, Chapter XIII, Theorem 2.3] with Z_2 coefficients to these subgroups to show that $dF(T)$ is at least $\frac{1}{2} \text{rank } G$.

REFERENCES

1. A. Borel, et al., *Seminar on transformation groups*, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J., 1960. MR 22 #7129.
2. N. Bourbaki, *Éléments de mathématiques*. Fasc. XXXIV. *Groupes et algèbres de Lie*. Chaps. 4, 5, 6, Actualités Sci. Indust., no. 1337, Hermann, Paris, 1968. MR 39 #1590.
3. G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
4. W.-C. Hsiang and W.-Y. Hsiang, *A fixed theorem for finite diffeomorphism groups generated by reflections*, The Steenrod Algebra and its Applications (Proc. Conf. to Celebrate N. E. Steenrod's Sixtieth Birthday, Battelle Memorial Inst., Columbus, Ohio, 1970), Lecture Notes in Math., vol. 168, Springer-Verlag, Berlin, 1970, pp. 90–106. MR 42 #8519.
5. W.-Y. Hsiang, *On the splitting principle and the geometric weight system of topological transformation groups*. I, Proc. Second Conf. on Compact Transformation Groups, part 1, Lecture Notes in Math., vol. 298, Springer-Verlag, Berlin and New York, 1972.
6. N. E. Steenrod, *Cohomology operations*, Ann. of Math. Studies, no. 50, Princeton Univ. Press, Princeton, N. J., 1962. MR 26 #3056.
7. R. Sullivan, *Differentiable actions of classical groups on spheres*, Topology 9 (1970), 155–167. MR 41 #6244.
8. ———, *Linear models for compact groups acting on spheres*, Topology 13 (1974), 77–87.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, ALBANY,
NEW YORK 12222