A NOTE ON A COROLLARY OF SARD'S THEOREM

JOHN C. WELLS

ABSTRACT. A corollary of Sard's theorem is the following:

Corollary. Let $f: K \rightarrow R^n$ be a smooth (i.e. $f \in C^k$, $k \geq 1$) map defined on a compact subset K of R^n . Let $C = \{y | f^{-1}(y) \text{ is infinite}\}$. Then the Lebesgue measure of C is zero.

The purpose of this note is to show that a similar version of this theorem holds for Lipschitz functions.

Theorem. Let $f: K \rightarrow R^n$ with f Lipschitz, K a measurable subset of R^n and $m(K)$ the Lebesgue measure of K , less than ∞ . Let $C = \{y | f^{-1}(y) \text{ is infinite}\}$. Then $m(C) = 0$.

The proof of the theorem will require two lemmas.

Lemma 1. Let $f: K \rightarrow R^m$ with K a compact subset of R^n and f a continuous function. Then there exists $K' \subset K$ with $f(K') = f(K)$, $f|_{K'}$ one to one and K' a Borel set.

Proof. Let x_i denote the i th coordinate of x . Define for $j = 1, 2, 3, \dots$, $A_j = \{x | x \in K \text{ and there exists } y \in K \text{ with } y_1 \leq x_1 - 1/j \text{ and } f(y) = f(x) \text{ or } y \in K \text{ with } y_1 = x_1, y_2 \leq x_2 - 1/j \text{ and } f(y) = f(x) \text{ or } \dots \text{ or } y \in K \text{ with } y_1 = x_1, \dots, y_{n-1} = x_{n-1}, y_n \leq x_n - 1/j \text{ and } f(y) = f(x)\}$. Since A_j is compact for each j , $K' = K - \bigcup_j A_j$ is Borel. If $x \in f(K)$ observe that y defined by

$$y_1 = \inf \{z_1 | f(z) = f(x)\}, \dots, y_n = \inf \{z_n | z_1 = y_1, \dots, z_{n-1} = y_{n-1}, f(z) = f(x)\}$$

is not contained in A_j for any j and also $f(y) = f(x)$. Thus $f(K') = f(K)$. Clearly $f|_{K'}$ is one to one.

Lemma 2. Let $f: K \rightarrow R^m$ be a continuous map from K a measurable subset of R^n . Then there exist K', K'' with $K'' \subset K' \subset K$, K'' and K' Borel sets, $f|_{K''}$ one to one, $m(K - K') = 0$ and $f(K'') = f(K')$.

Proof. There exists an F_σ set K' , with $K' \subset K$ and $m(K - K') = 0$, and we can write $K' = \bigcup_n K_n$ where K_n is compact for each n . By Lemma 1

Received by the editors April 3, 1974 and, in revised form, June 26, 1974.
 AMS (MOS) subject classifications (1970). Primary 26A63, 28A75.

Copyright © 1975. American Mathematical Society

there are Borel sets $K'_n \subset K_n$ with $f(K'_n) = f(K_n)$ and $f|_{K'_n}$ one to one for each n . Define recursively $D_1 = K'_1, \dots, D_k = D_{k-1} \cup (K'_k - f^{-1}(f(D_{k-1})))$. Then $K'' = \bigcup_k D_k$ is Borel, $f(K'') = f(K')$ and $f|_{K''}$ is one to one.

Proof of Theorem. Suppose $m(C) = a > 0$. Let $A_1 = f^{-1}(C)$ and find by Lemma 2 $N''_1 \subset N'_1 \subset A_1$ with N''_1 and N'_1 Borel, $m(A_1 - N'_1) = 0$, $f|_{N''_1}$ one to one and $f(N''_1) = f(N'_1)$. Since f is Lipschitz, $m(f(A_1 - N'_1)) = 0$. Therefore $m(f(N''_1)) = a$. Now $f(A_1 - N''_1) = C$ by the definition of C . Thus for $k = 2, 3, \dots$ by letting $A_k = A_{k-1} - N''_{k-1}$ we can repeat the above argument to find $N''_k \subset N'_k \subset A_k$ with $f|_{N''_k}$ one to one, $f(N''_k) = f(N'_k)$ and $m(f(N''_k)) = a$. Suppose that L is the Lipschitz constant for f ; then $m(N''_k) \geq a/L$ for each k . But since $N''_k \cap N''_{k'} = \emptyset$ for $k \neq k'$, this implies $m(K) = \infty$, a contradiction.

Remarks. The requirement that f be Lipschitz in the theorem cannot be weakened to a requirement of continuity. To show this let f be the first coordinate of a continuous map of the unit interval I onto the unit square I^2 (i.e. a space-filling curve). Then $f^{-1}(y)$ is infinite for each $y \in I$.

On the other hand, for $n = 1$ absolute continuity will suffice as the following argument shows. Suppose f is a real valued a.c. function on I . Then f is differentiable except on a set A_1 of measure zero. $m(f(A_1)) = 0$ since f is a.c. Let $A_2 = \{x \mid f'(x) = 0\}$ and let η be an arbitrary positive number. For each $x \in A_2$ there are arbitrarily small intervals containing x such that $|f(y) - f(x)| \leq |y - x|$ when y is in the intervals. Thus the collection $\mathcal{F}$ of all such intervals with x varying over A_2 is a Vitale covering of A_2 , so there is a disjoint collection $\{I_n\} \subset \mathcal{F}'$ with $m(A_2 - \bigcup_n I_n) = 0$. But then

$$m(f(A_2)) \leq m\left(f\left(A_2 - \bigcup_n I_n\right)\right) + m\left(f\left(\bigcup_n I_n\right)\right) \leq 0 + 2\eta \cdot \sum_n \text{length}(I_n) \leq 2\eta.$$

Since η is arbitrary, $m(f(A_2)) = 0$. Finally, if $y \in f(I) - f(A_1) - f(A_2)$ then $f^{-1}(y)$ must consist of isolated points and hence be finite. Thus $m\{y \mid f^{-1}(y) \text{ is infinite}\} = 0$.

REFERENCES

1. John Milnor, *Topology from the differential viewpoint*, Univ. Press of Virginia, Charlottesville, Va., 1965. MR 37 #2239.
2. A. Sard, *The measure of the critical values of differential maps*, Bull. Amer. Math. Soc. 48 (1942), 883-890. MR 4, 153.

DEPARTMENT OF MATHEMATICS, CALIFORNIA STATE UNIVERSITY, NORTHRIDGE, CALIFORNIA 91324