A LOCAL CHARACTERIZATION OF DARBOUX $\mathcal{B}$ FUNCTIONS

RICHARD G. GIBSON

ABSTRACT. A. M. Bruckner and J. B. Bruckner gave the definition of Darboux $\mathcal{B}$ functions and proved a theorem which is a local characterization of real-valued Darboux $\mathcal{B}$ functions. The purpose of this paper is to generalize this theorem. To this end, the definition of a function being Darboux $\mathcal{B}$ at a point is given which has a metric continuum as its range. Hence, the theorem that a function is Darboux $\mathcal{B}$ if and only if it is Darboux $\mathcal{B}$ at each point.

1. Introduction. A function having domain and range a subset of the real line is said to be a *Darboux function* if the image of every connected subset of its domain is a connected set. A detail account of Darboux functions can be found in the survey article by Bruckner and Ceder [2].

A. M. Bruckner and J. B. Bruckner [1] gave the definition of Darboux $\mathcal{B}$ functions which are defined on a euclidean space and having a separable metric space as the range, and proved a theorem which is a local characterization of real-valued Darboux $\mathcal{B}$ functions. Then they made the statement that it would be of interest to know whether such a theorem exists in case the range is not a subset of the real line. The purpose of this paper is to give an answer to this query. The local condition given by Bruckner and Bruckner [1] was in terms of limit superior and limit inferior. That local condition will be replaced by one where the range of the function is a metric continuum. How much less one can assume about the range without losing the results is an open question.

§2 contains some of the needed definitions and a definition of Darboux $\mathcal{B}$ at a point which replaces the local condition given in [1]. §3 contains the main theorem which is preceded by a lemma.

2. Preliminaries. In this paper X will be a euclidean space, Y a metric continuum, and $\mathcal{B}$ a topological base for X such that each member of $\mathcal{B}$ is connected. A function f defined on X and having range Y is said to be

Received by the editors February 16, 1974.

AMS (MOS) subject classifications (1970). Primary 54C10; Secondary 26A15.

Key words and phrases. Darboux functions, Darboux $\mathcal{B}$ functions, Darboux $\mathcal{B}$ at a point, property L.

Copyright © 1975, American Mathematical Society

Darboux $\mathcal{B}$ provided that if U is in $\mathcal{B}$, then $f(\text{cl}(U))$ is connected in Y . A base $\mathcal{B}$ for X is said to satisfy *condition* (*) provided any translation of an element of $\mathcal{B}$ is in $\mathcal{B}$. The base $\mathcal{B}$ satisfies *condition* (**) provided that for each $x \in X$ and $U \in \mathcal{B}, x \in \text{cl}(U)$, there exists $V \in \mathcal{B}$ such that $x \in \text{cl}(V)$ and $\text{cl}(V) - (x) \subset U$.

Let f be a function defined on X and having range Y . The function f is said to have property L at the point x in X provided that, if $U \in \mathcal{B}$ and x is a boundary point of U , then $f(x)$ is a limit point of $f(U)$. If x is a point in X , then a point y in Y is said to be a *limit point of f at x* provided that there exists a sequence $(x_n)_{n=1}^{\infty}$ of points in X which converges to x such that the sequence $(f(x_n))_{n=1}^{\infty}$ converges to y .

Definition. Let X be a euclidean space and let $\mathcal{B}$ be a base of connected sets for X which satisfies conditions (*) and (**). Let f be a function defined on X and having range Y . The function f is said to be *Darboux* $\mathcal{B}$ at the point p in X provided that the following statements are true.

- (1) f has property L at the point p and $f(p)$ is a limit point of f at p .
- (2) If $U \in \mathcal{B}, p \in U$, and a and b are limit points of f at p relative to U , then for each closed subset C of Y which separates a from b there exists $x \in U$ such that $f(x)$ is in C .

3. **A local characterization.** A lemma will be proved before proving the main theorem.

Lemma. Let f be a function defined on X and having range Y and let $\mathcal{B}$ be a basis for X which satisfies conditions (*) and (**). Let $U \in \mathcal{B}$ and let $f(A)$ and $f(B)$ be two mutually separated sets such that the union of A and B is $\text{cl}(U)$. If f has property L at each point in X and P is the boundary of A relative to $\text{cl}(U)$, then the following statements are true.

- (1) P is the boundary of B relative to $\text{cl}(U)$.
- (2) P is closed in X .
- (3) $P \cap U$ is nonempty, dense-in-itself, and of type G_δ .
- (4) $A \cap P'$ and $B \cap P'$ are both dense in $P' = P \cup U$.

Proof. It is obvious that P is the boundary of B relative to $\text{cl}(U)$ and P is closed in X .

To prove that $P \cap U$ is nonempty, note that P is nonempty; for, if P were empty, then A and B would form a separation of the connected set $\text{cl}(U)$. Suppose $P \cap U$ is empty. Then $P \subset \text{cl}(U) - U$, and $U \subset A$ or $U \subset B$. Assume $U \subset A$. Choose any x in B . Then x is in $\text{cl}(U) - U$ and x is a boundary point of U . Thus $f(x)$ is a limit point of $f(U)$. So $f(x)$ is a limit point of

$f(A)$ which is in $f(B)$. This contradicts the fact that $f(A)$ and $f(B)$ are mutually separated. Therefore $P \cap U$ is nonempty.

Now suppose $P \cap U$ is not dense-in-itself. Then there is a point x in $P \cap U$, say x is in $P \cap A$, and a V in $\mathfrak{B}$, $V \subset U$, such that $P \cap V = (x)$. If X is of dimension greater than or equal to 2, then $V - (x)$ is connected. So $V - (x) \subset A$ or $V - (x) \subset B$. Since x is in P , $V - (x)$ is not a subset of A . So $V - (x) \subset B$. By condition (**), there is a $W \in \mathfrak{B}$ such that x is in $\text{cl}(W)$ and $\text{cl}(W) - (x) \subset V - (x) \subset B$. Since x is a boundary point of W , $f(x)$ is a limit point of $f(W)$. So $f(x)$ is a limit point of $f(B)$ which is in $f(A)$. This is a contradiction. Thus $P \cap U$ is dense-in-itself if X is of dimension greater than or equal to 2. If the dimension of X is 1, the result is easily proved.

Since P is closed and U is open, $P \cap U$ is of type G_δ .

To prove that $A \cap P'$ is dense in P' , suppose that $A \cap P'$ is not dense in P' . Let $V \subset U$ be an open sphere centered at a point $p \in P' \cap B$ such that $V \cap A \cap P'$ is empty. Then $A \cap V$ is open and nonempty. Let x be a point of $A \cap V$ such that $d(x, X - V)$ is greater than $2d(x, p)$ where d is the euclidean metric of X . Choose $W \in \mathfrak{B}$ such that $x \in W \subset A \cap V$ and $d(\text{cl}(W), X - V)$ is greater than $d(\text{cl}(W), p)$. Let q be a point of B nearest x . Then q is in V . Let W' be a translation of W such that $q \in \text{cl}(W') - W'$ and $W' \subset A$. By condition (*), $W' \in \mathfrak{B}$. Now choose $W'' \in \mathfrak{B}$ such that $q \in \text{cl}(W'')$ and $\text{cl}(W'') - (q) \subset W'$. Now $q \in B$ and $\text{cl}(W'') - (q) \subset A$. Since q is a boundary point of W'' , we have a contradiction by the hypothesis of the theorem. Thus $A \cap P'$ is dense in P' . In a similar way, it can be shown that $B \cap P'$ is dense in P' .

Theorem. Let X be a euclidean space and let $\mathfrak{B}$ be a base of connected sets for X which satisfies conditions (*) and (**). Let f be a function defined on X and having a metric continuum Y as its range. Then f is Darboux $\mathfrak{B}$ if and only if f is Darboux $\mathfrak{B}$ at each point in X .

Proof. Suppose f is not Darboux $\mathfrak{B}$. Then there exists $U \in \mathfrak{B}$ such that $f(\text{cl}(U))$ is not connected. Let $f(A)$ and $f(B)$ be two mutually separated sets such that the union of A and B is $\text{cl}(U)$.

Let $Y(A)$ and $Y(B)$ be two disjoint open subsets of Y such that $f(A) \subset Y(A)$ and $f(B) \subset Y(B)$. Let C be the complement of $Y(A) \cup Y(B)$. Then C separates $Y(A)$ and $Y(B)$. Let P be the boundary of A in $\text{cl}(U)$ and let $P' = P \cap U$. By the lemma both the sets $A \cap P'$ and $B \cap P'$ are dense in P' .

We now show that if p is in $A \cap U$ and W is an open set containing C , then there exists an open set $V \in \mathfrak{B}$ containing p such that if $x \in \text{cl}(V)$,

then $f(x) \in Y(A) \cup W$. Suppose the condition fails for some $p \in A \cap U$ and $W \supset C$. Let $V_1, V_2, \dots$ be a sequence of sets each containing p such that $V_k \in \mathcal{B}$ for each $k = 1, 2, \dots$ and $\bigcap_{k=1}^{\infty} V_k = \{p\}$. Then for each k there exists $x_k \in V_k$ such that $f(x_k)$ is not in $Y(A) \cup W$. Since $Y(A) \cup W$ is open and Y is compact, $Y - Y(A) \cup W$ is compact. Thus there exists a subsequence of $(f(x_k))_{k=1}^{\infty}$ which converges to some point $y \in Y - Y(A) \cup W$, and hence y is a limit point of f at p which is in $Y(B)$. Since $f(p) \in Y(A)$ and $f(p)$ is a limit point of f at p , there exists $x \in U$ such that $f(x) \in C$. This is a contradiction. Similarly, if $p \in B \cap U$ and W is an open set containing C , then there exists $V \in \mathcal{B}$ containing p such that if $x \in \text{cl}(V)$, then $f(x) \in Y(B) \cup W$.

Let $W_1, W_2, \dots$ be a sequence of open subsets of Y such that $\bigcap_{k=1}^{\infty} W_k = C$. Let $p_1 \in A \cap P'$. Choose $V_1 \in \mathcal{B}$ such that $p_1 \in V_1$, the diameter of $V_1 < 1$, $\text{cl}(V_1) \subset U$, and if $x \in \text{cl}(V_1)$, then $f(x) \in Y(A) \cup W_1$. Since $B \cap P'$ is dense in P' , there is a p_2 in $B \cap P' \cap V_1$. Let $V_2 \in \mathcal{B}$ such that $p_2 \in V_2$, $V_2 \subset V_1$, the diameter of $V_2 < 1/2$, and if $x \in \text{cl}(V_2)$, then $f(x) \in Y(B) \cup W_2$. Continuing in this manner, we obtain a sequence of points $(p_k)_{k=1}^{\infty}$ and a sequence of open sets $(V_k)_{k=1}^{\infty}$ such that for each k , the following conditions are satisfied.

1. $p_k \in V_k \cap P'$, $V_k \in \mathcal{B}$, $V_{k+1} \subset V_k$, the diameter of $V_k < 1/k$.
2. If $x \in \text{cl}(V_k)$ and k is odd, then $f(x) \in Y(A) \cup W_k$.
3. If $x \in \text{cl}(V_k)$ and k is even, then $f(x) \in Y(B) \cup W_k$.

Note that the set $\bigcap_{k=1}^{\infty} \text{cl}(V_k)$ consists of a single point, say p .

Since $\text{cl}(V_1) \subset U$, $p \in U$. Since $p \in \text{cl}(V_k)$ for each $k = 1, 2, \dots$, it follows that $f(p)$ is in each W_k . Thus $f(p) \in \bigcap_{k=1}^{\infty} W_k$, and so $f(p) \in C$. This is a contradiction, and hence f is Darboux $\mathcal{B}$.

We now assume f is Darboux $\mathcal{B}$ and prove that f is Darboux $\mathcal{B}$ at each point of X . The fact that f has property L of the definition of Darboux $\mathcal{B}$ at each point is obvious. We now prove that $f(p)$ is a limit point of f at p . Let $(W_k)_{k=1}^{\infty}$ be a descending sequence of open subsets of Y each containing $f(p)$ such that $\bigcap_{k=1}^{\infty} W_k = \{f(p)\}$. Let $(V_k)_{k=1}^{\infty}$ be a descending sequence of sets in $\mathcal{B}$ each containing p such that $\bigcap_{k=1}^{\infty} V_k = \{p\}$. Since f is Darboux $\mathcal{B}$ and $p \in V_k$, $f(V_k)$ is a connected subset of Y which contains $f(p)$ for each $k = 1, 2, \dots$. Now there exists $x_k \in V_k$ such that $x_k \neq p$ and $f(x_k) \in W_k$ for each $k = 1, 2, \dots$. The sequence $(x_k)_{k=1}^{\infty}$ converges to p and the sequence $(f(x_k))_{k=1}^{\infty}$ converges to $f(p)$. Therefore $f(p)$ is a limit point of f at p .

We now prove part (2) of the definition of Darboux $\mathcal{B}$ at a point. Let U

$\in \mathfrak{B}$, $p \in U$, and let a and b be limit points of f at p relative to U . Let C be a closed subset of Y which separates a from b . Thus there exists two mutually separated open sets V and W such that $Y - C = V \cup W$, $a \in V$ and $b \in W$. Since a and b are limit points of f at p , there exist $y, z \in U$ such that $f(y) \in V$ and $f(z) \in W$. Thus C separates $f(y)$ and $f(z)$. Since $f(U)$ is connected, there exists $x \in U$ such that $f(x) \in C$, and we are finished.

REFERENCES

1. A. M. Bruckner and J. B. Bruckner, *Darboux transformations*, Trans. Amer. Math. Soc. 128 (1967), 103–111. MR 36 #1586.
2. A. M. Bruckner and J. G. Ceder, *Darboux continuity*, Jber. Deutsch. Math.-Verein. 67 (1964/65), Abt. 1, 93–117. MR 32 #4217.
3. James Dugundji, *Topology*, Allyn and Bacon, Boston, Mass., 1967. MR 33 #1824.

DEPARTMENT OF MATHEMATICS, ALABAMA STATE UNIVERSITY, MONTGOMERY,
ALABAMA 36101