SOLUTION OF INTEGRAL EQUATIONS BY PRODUCT INTEGRATION

JON C. HELTON

ABSTRACT. Suppose f , h and G are functions with values in a normed complete ring. With suitable restrictions on these functions, it is established that

$$f(x) = h(x) + \int_a^x f(u)G(u, v)$$

for $a \leq x \leq b$ only if $\int_a^x h(u)G(u, v) \Pi^x(1 + G)$ exists and is $f(x) - h(x)$ for $a \leq x \leq b$, and that

$$f(x) = h(x) + \int_a^x G(u, v)f(u)$$

for $a \leq x \leq b$ only if $\int_a^x \Pi^v(1 + \mathcal{G})G(u, v)h(u)$ exists and is $f(x) - h(x)$ for $a \leq x \leq b$, where $\mathcal{G}(s, r) = G(r, s)$.

Lower case letters are used to represent functions from R to N , and capital letters are used to represent functions from $R \times R$ to N , where R denotes the set of real numbers and N denotes a ring which has a multiplicative identity element represented by 1 and a norm $|\cdot|$ with respect to which N is complete and $|1| = 1$. For a subdivision $\{x_i\}_{i=0}^n$ of an interval $[a, b]$, we use G_i and f_i to denote $G(x_{i-1}, x_i)$ and $f(x_i)$, respectively.

The statement that $G \in OB^\circ$ on $[a, b]$ means there exist a subdivision D of $[a, b]$ and a number B such that, if $\{x_i\}_{i=0}^n$ is a refinement of D , then $\sum_{i=1}^n |G_i| < B$.

The statement that $\int_a^b G$ exists means there exists an element L of N such that, if $\epsilon > 0$, then there exists a subdivision D of $[a, b]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D , then $|L - \sum_{i=1}^n G_i| < \epsilon$. Further, $G \in OA^\circ$ on $[a, b]$ only if $\int_a^b G$ exists and $\int_a^b |G - \int G| = 0$.

The statement that $\int_a^b \Pi^b(1 + G)$ exists means there exists an element L

Presented to the Society, March 4, 1974; received by the editors January 17, 1974.

AMS (MOS) subject classifications (1970). Primary 45N05; Secondary 26A39, 26A42.

Key words and phrases. Sum integral, product integral, subdivision-refinement integral, integral equation, interval function, normed complete ring.

Copyright © 1975, American Mathematical Society

of N such that, if $\epsilon > 0$, then there exists a subdivision D of $[a, b]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D , then $|L - \prod_{i=1}^n (1 + G_i)| < \epsilon$. Further, $G \in OM^\circ$ on $[a, b]$ only if ${}_x\Pi^\gamma(1 + G)$ exists for $a \leq x \leq y \leq b$ and $\int_a^b |1 + G - \Pi(1 + G)| = 0$. The existence of integrals is defined similarly on intervals $[p, q]$, where $q < p$.

The function h is quasi-continuous on $[a, b]$ only if $\lim_{x \rightarrow p^-} h(x)$ exists for $a < p \leq b$ and $\lim_{x \rightarrow p^+} h(x)$ exists for $a \leq p < b$. Further, $G \in OL^\circ$ on $[a, b]$ only if $\lim_{x \rightarrow p^-} G(x, p)$ and $\lim_{x, y \rightarrow p^-} G(x, y)$ exist for $a < p \leq b$, and $\lim_{x \rightarrow p^+} G(p, x)$ and $\lim_{x, y \rightarrow p^+} G(x, y)$ exist for $a \leq p < b$. See B. W. Helton [2] and J. S. Mac Nerney [6] for additional background.

The theorems in this paper were suggested by a result of J. S. Mac Nerney [5, Theorem 3.4, p. 361]. However, their proofs are constructed by using techniques similar to those employed by B. W. Helton [2, §5, pp. 307–314]. There, the integral equations

$$f(x) = h(x) + \int_a^x f(u)G(u, v) \quad \text{and} \quad f(x) = h(x) + \int_a^x G(u, v)f(u)$$

were solved with the restriction that h have bounded variation. We now solve these integral equations with the restriction that h be quasi-continuous. The reader is also referred to related results by D. B. Hinton [4, Theorems 4.1, 4.2, 4.3, 4.4, pp. 325–327] and C. W. Bitzer [1, Theorems 4.1, 4.2, 5.9, pp. 447–451].

We now state three lemmas that are used in the development of the results in this paper.

Lemma 1. *If G is a function from $R \times R$ to N and $G \in OB^\circ$ on $[a, b]$, then the following statements are equivalent:*

- (1) $G \in OA^\circ$ on $[a, b]$, and
- (2) $G \in OM^\circ$ on $[a, b]$ [2, Theorem 3.4, p. 301].

Lemma 2. *If H and G are functions from $R \times R$ to N , $H \in OL^\circ$ on $[a, b]$ and G is in OA° and OB° on $[a, b]$, then GH and HG are in OA° and OM° on $[a, b]$ [3, Theorem 2, p. 494].*

Lemma 3. *If F and G are functions from $R \times R$ to N belonging to OB° on $[a, b]$, $F \in OA^\circ$ on $[a, b]$ and each of ${}_x\Pi^\gamma(1 + G)$ and $\int_x^y F(u, v) {}_v\Pi^\gamma(1 + G)$ exists for $a \leq x < y \leq b$, then*

$$\int_a^b \left| F(x, y) - \int_x^y F(u, v) {}_v\Pi^\gamma(1 + G) \right|$$

exists and is zero [2, Lemma, p. 307].

Theorem 1. If f and h are functions from R to N , G is a function from $R \times R$ to N , h is quasi-continuous on $[a, b]$ and $G \in OB^\circ$ on $[a, b]$, then the following statements are equivalent:

- (1) f is bounded on $[a, b]$, $G \in OA^\circ$ on $[a, b]$, $f(u)G(u, v) \in OA^\circ$ on $[a, b]$ and $f(x) = h(x) + \int_a^x f(u)G(u, v)$ for $a \leq x \leq b$, and
- (2) $G \in OM^\circ$ on $[a, b]$ and $\int_a^x h(u)G(u, v) \prod_v^\times (1 + G)$ exists and is $f(x) - h(x)$ for $a \leq x \leq b$.

Proof. (1) $\rightarrow$ (2). Since G is in OA° and OB° on $[a, b]$, it follows from Lemma 1 that $G \in OM^\circ$ on $[a, b]$. Suppose $a \leq x \leq b$. If $a = x$, then (2) follows immediately. Therefore, suppose $a < x$. The existence of $I(a, x)$ follows from Lemma 2, where

$$I(r, s) = \int_r^s h(u)G(u, v) \prod_v^\times (1 + G)$$

for $a \leq r \leq s \leq b$. We now show that $I(a, x)$ is equal to $f(x) - h(x)$. Let $\epsilon > 0$.

There exists a subdivision D_1 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_1 , then

$$\left| I(a, x) - \sum_{i=1}^n h_{i-1} G_i \prod_{x_i}^\times (1 + G) \right| < \epsilon/3.$$

There exist a subdivision D_2 of $[a, x]$ and a number B such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_2 , then

$$(1) \prod_{i=1}^n (1 + |G_i|) < B, \text{ and}$$

$$(2) \sum_{i=1}^n |h_{i-1} G_i| < B.$$

Since G is in OM° and OB° on $[a, x]$, there exists a subdivision D_3 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_3 and $1 \leq i \leq n$, then

$$\left| \prod_{x_i}^\times (1 + G) - \prod_{k=i+1}^n (1 + G_k) \right| < \epsilon(3B)^{-1}.$$

Since $f(u)G(u, v) \in OA^\circ$ on $[a, x]$, there exists a subdivision D_4 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_4 , then

$$\sum_{i=1}^n \left| f_{i-1} G_i - \int_{x_{i-1}}^{x_i} f(u)G(u, v) \right| < \epsilon(3B)^{-1}.$$

Let D denote the subdivision $\bigcup_{i=1}^4 D_i$ of $[a, x]$. Suppose $\{x_i\}_{i=0}^n$ is a refinement of D . For $1 \leq i \leq n$, we have that

$$f_i - f_{i-1} = h_i - h_{i-1} + f_{i-1}G_i + c_i,$$

where

$$c_i = \int_{x_{i-1}}^{x_i} f(u)G(u, v) - f_{i-1}G_i.$$

Thus,

$$f_i = h_i - h_{i-1} + f_{i-1}(1 + G_i) + c_i.$$

Now, by using iteration for $i = 1, 2, \dots, n$, we have that

$$f_n = h_n + \sum_{i=1}^n h_{i-1}G_i \prod_{k=i+1}^n (1 + G_k) + \sum_{i=1}^n c_i \prod_{k=i+1}^n (1 + G_k).$$

Hence,

$$\begin{aligned} & |I(a, x) - \{f(x) - h(x)\}| \\ &= \left| h(x) + I(a, x) - h_n \right. \\ &\quad \left. - \sum_{i=1}^n h_{i-1}G_i \prod_{k=i+1}^n (1 + G_k) - \sum_{i=1}^n c_i \prod_{k=i+1}^n (1 + G_k) \right| \\ &\leq \left| \sum_{i=1}^n h_{i-1}G_i \prod_{k=i+1}^n (1 + G_k) - \sum_{i=1}^n h_{i-1}G_i \prod_{k=i+1}^n (1 + G_k) \right| \\ &\quad + \left| I(a, x) - \sum_{i=1}^n h_{i-1}G_i \prod_{k=i+1}^n (1 + G_k) \right| + \sum_{i=1}^n |c_i| \prod_{k=i+1}^n (1 + G_k) \\ &< \sum_{i=1}^n |h_{i-1}G_i| \prod_{k=i+1}^n (1 + G_k) + \epsilon/3 + B[\epsilon(3B)^{-1}] \\ &< B[\epsilon(3B)^{-1}] + 2\epsilon/3 = \epsilon. \end{aligned}$$

Therefore, (1) implies (2).

Proof. (2) $\rightarrow$ (1). Since h is bounded on $[a, b]$ and $G \in OB^\circ$ on $[a, b]$, it follows that f is bounded on $[a, b]$. Since G is in OM° and OB° on $[a, b]$, it follows from Lemma 1 that $G \in OA^\circ$ on $[a, b]$. Further, it follows by employing Lemma 2 that $f(u)G(u, v) \in OA^\circ$ on $[a, b]$. Suppose $a \leq x \leq b$. If $a = x$, then (2) follows immediately. Therefore, suppose $a < x$. Let $\epsilon > 0$.

There exists a subdivision D_1 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_1 , then

$$\left| \int_a^x f(u)G(u, v) - \sum_{i=1}^n f_{i-1}G_i \right| < \frac{\epsilon}{3}.$$

There exists a number B such that, if $a \leq t \leq x$, then $|I(a, t)| < B$, where $I(a, t)$ is defined as in the first part of the proof. Now, since $G \in OM^\circ$ on $[a, x]$, there exists a subdivision D_2 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_2 , then

$$\sum_{i=1}^n \left| 1 + G_i - \prod_{x_{i-1}}^{x_i} (1 + G) \right| < \epsilon (3B)^{-1}.$$

It follows by applying Lemma 3 that there exists a subdivision D_3 of $[a, x]$ such that, if $\{x_i\}_{i=0}^n$ is a refinement of D_3 , then

$$\sum_{i=1}^n |h_{i-1} G_i - I(x_{i-1}, x_i)| < \frac{\epsilon}{3}.$$

Let D denote the subdivision $\bigcup_{i=1}^3 D_i$ of $[a, x]$. Suppose $\{x_i\}_{i=0}^n$ is a refinement of D . Thus,

$$\begin{aligned} & \left| h(x) + \int_a^x f(u)G(u, v) - f(x) \right| \\ & \leq \left| h(x) + \sum_{i=1}^n f_{i-1} G_i - h(x) - I(a, x) \right| + \left| \int_a^x f(u)G(u, v) - \sum_{i=1}^n f_{i-1} G_i \right| \\ & < \left| \sum_{i=1}^n \{h_{i-1} + I(a, x_{i-1})\} G_i - I(a, x) \right| + \epsilon/3 \\ & = \left| \sum_{i=1}^n \{h_{i-1} G_i + [I(a, x_{i-1})][1 + G_i] - I(a, x_{i-1})\} - I(a, x) \right| + \epsilon/3 \\ & \leq \left| \sum_{i=1}^n \left\{ h_{i-1} G_i + [I(a, x_{i-1})] \left[\prod_{x_{i-1}}^{x_i} (1 + G) \right] - I(a, x_{i-1}) \right\} - I(a, x) \right| \\ & \quad + \sum_{i=1}^n \left| I(a, x_{i-1}) \right| \left| 1 + G_i - \prod_{x_{i-1}}^{x_i} (1 + G) \right| + \epsilon/3 \\ & < \left| \sum_{i=1}^n \{I(a, x_i) - I(a, x_{i-1})\} - I(a, x) \right| \\ & \quad + \sum_{i=1}^n |h_{i-1} G_i - I(x_{i-1}, x_i)| + B[\epsilon (3B)^{-1}] + \epsilon/3 \\ & < 0 + \epsilon/3 + 2\epsilon/3 = \epsilon. \end{aligned}$$

Therefore, (2) implies (1).

Theorem 2. If f and h are functions from R to N , G is a function from $R \times R$ to N , $\mathcal{G}(y, x) = G(x, y)$ for $a \leq x < y \leq b$, h is quasi-continuous on $[a, b]$ and $G \in OB^\circ$ on $[a, b]$, then the following statements are equivalent:

- (1) f is bounded on $[a, b]$, $G \in OA^\circ$ on $[a, b]$, $G(u, v)f(u) \in OA^\circ$ on $[a, b]$ and $f(x) = h(x) + \int_a^x G(u, v)f(u)$ for $a \leq x \leq b$, and
- (2) $\mathcal{G} \in OM^\circ$ on $[b, a]$ and $\int_a^x \Pi^v (1 + \mathcal{G})G(u, v)h(u)$ exists and is $f(x) - h(x)$ for $a \leq x \leq b$.

The proof of Theorem 2 is similar to the proof of Theorem 1, and therefore, we omit it.

REFERENCES

1. C. W. Bitzer, *Stieltjes-Volterra integral equations*, Illinois J. Math. **14** (1970), 434–451.
2. B. W. Helton, *Integral equations and product integrals*, Pacific J. Math. **16** (1966), 297–322. MR **32** #6167.
3. ———, *A product integral representation for a Gronwall inequality*, Proc. Amer. Math. Soc. **23** (1969), 493–500. MR **40** #1562.
4. D. B. Hinton, *A Stieltjes-Volterra integral equation theory*, Canad. J. Math. **18** (1966), 314–331. MR **32** #6169.
5. J. S. Mac Nerney, *Stieltjes integrals in linear spaces*, Ann. of Math. (2) **61** (1955), 354–367. MR **16**, 716.
6. ———, *Integral equations and semigroups*, Illinois J. Math. **7** (1963), 148–173. MR **26** #1726.

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85281