ON A CONJECTURE OF GROSS

HUI-HSIUNG KUO¹

ABSTRACT. Gross' conjecture about the generalized Laplacian is proved as a consequence of the main theorem.

1. **Gross' conjecture.** The generalized Laplacian of a Borel measurable function f in an abstract Wiener space (H, B) is defined by

$$\Delta f(x) = 2 \lim_{r \downarrow 0} \{E[r_x^{(r)}]\}^{-1} \{E[f(x + W(r_x^{(r)}))] - f(x)\},$$

where W is a Wiener process in B starting at the origin and $\tau_x^{(r)}$ is the first exit time for $x + W$ from the open ball of radius r in B with center x . Let A be a bounded operator from B to B^* and $u(x) = \frac{1}{2} \langle Ax, x \rangle$. Under the assumption that the B norm $\|\cdot\|$ is twice continuously B -differentiable away from the origin and that the second Fréchet B -derivative is bounded on the annulus $1 \leq \|x\| \leq 2$, Gross [2, p. 148] showed that $\Delta u(0) = \text{trace}(A|H)$. Then he conjectured that the conclusion remains true without any differentiability assumption on the B norm or any assumption concerning the existence of smooth functions on B with bounded support. The purpose of this note is to prove this conjecture. In order to use the results in [3], we assume that there exists a sequence of finite dimensional projections with range in B^* converging strongly to the identity both in B and in H .

2. **Main theorem.** We use the same notation as in [3].

Theorem 1. *Let f be a twice continuously H -differentiable function in B such that $D^2f(x) \in \mathfrak{B}_1(H, H)$, the Banach space of trace class operators of H , for all x in B and D^2f is continuous from B to $\mathfrak{B}_1(H, H)$. Let τ be a stopping time (w.r.t. W) such that $\tau < \infty$ a.s. and $\int_1^\infty E[1_{t \leq \tau} |Df(x_0 + W(t))|^2] dt < \infty$. Then*

$$E[f(x_0 + W(\tau))] = f(x_0) + E\left[\int_0^\tau \frac{1}{2} \text{trace } D^2f(x_0 + W(t)) dt\right].$$

Received by the editors February 7, 1974 and, in revised form, May 28, 1974.
AMS (MOS) subject classifications (1970). Primary 60H05, 60J45.

¹Research supported by NSF Grant GP-38010.

Proof. From Ito's formula [3, Theorem 4.1]

$$\begin{aligned} f(x_0 + W(\tau)) &= f(x_0) + \int_0^\tau (Df(x_0 + W(t)), dW(t)) \\ &\quad + \int_0^\tau \frac{1}{2} \text{trace } D^2 f(x_0 + W(t)) dt. \end{aligned}$$

Upon taking expectations, we obviously obtain the conclusion provided $E[\int_0^\tau (Df(x_0 + W(t)), dW(t))] = 0$. This follows from

Lemma 1. *Let τ be a stopping time and $\sigma(t)$ an H -valued nonanticipating process. Assume $\tau < \infty$ a.s. and $\int_1^\infty E[1_{t \leq \tau} |\sigma(t)|^2] dt < \infty$. Then $E[\int_0^\tau (\sigma(t), dW(t))] = 0$.*

Proof. The proof of [4, §2.3(4)] can be easily modified to show that

$$\int_0^\tau (\sigma(t), dW(t)) = \int_0^\infty (1_{t \leq \tau} \sigma(t), dW(t)).$$

But

$$\begin{aligned} E \left[\int_n^m (1_{t \leq \tau} \sigma(t), dW(t)) \right]^2 &= E \left[\int_n^m |1_{t \leq \tau} \sigma(t)|^2 dt \right] \\ &= \int_n^m E[1_{t \leq \tau} |\sigma(t)|^2] dt \rightarrow 0 \quad \text{as } n, m \rightarrow \infty. \end{aligned}$$

This gives the conclusion since $E[\int_0^n (\sigma(t), dW(t))] = 0$ for any $n \geq 0$.

Corollary 1. *Let A be a bounded operator from B to B^* and $u(x) = \frac{1}{2} \langle Ax, x \rangle$. Then $\Delta u(x) = \text{trace}(A|H)$ for all x in B .*

Proof. It is easy to see that $\|(A|H)^*(b)\|_{B^*} \leq \|A\|_{B, B^*} \|b\|$. Hence $(A|H)^*$ extends uniquely to a bounded operator from B to B^* . Obviously, $Du(x) = \frac{1}{2} \{Ax + (A|H)^*x\}$ and $D^2u(x) = \frac{1}{2} \{(A|H) + (A|H)^*\}$. It is well known that $(A|H)$ is a trace class operator of H , hence so is $(A|H)^*$. Let $a = \|A\|_{B, H}$ and $b = \|(A|H)^*\|_{B, H}$. Then

$$\begin{aligned} E[1_{t \leq \tau_x(r)} |Du(x + W(t))|^2] &\leq (a^2 + b^2) E[1_{t \leq \tau_x(r)} \|x + W(t)\|^2] \\ &\leq (a^2 + b^2) r^2 E[1_{t < \tau_x(r)}] = (a^2 + b^2) r^2 P(\tau_x^{(r)} \geq t). \end{aligned}$$

But $\{\omega; \tau_x^{(r)}(\omega) \geq t\} \subset \{\omega; \|W(t)\| \leq r\}$. Hence,

$$\begin{aligned} E[1_{t \leq \tau_x^{(r)}} |Du(x + W(t))|^2] &\leq (a^2 + b^2)r^2 P(\|W(t)\| \leq r) \\ &= (a^2 + b^2)r^2 p_t(S_r), \end{aligned}$$

where $S_r = \{y; \|y\| \leq r\}$ and p_t is the Wiener measure.

Now, $\int_1^\infty p_t(S_r) dt \leq \int_0^\infty p_t(S_r) dt = G(S_r)$, which is finite by [2, Remark 3.5, p. 147]. Therefore, u satisfies the assumption in Theorem 1 for all x in B and we have

$$E[u(x + W(\tau_x^{(r)}))] - u(x) = \frac{1}{2} E \left[\int_0^{\tau_x^{(r)}} \text{trace}(A|H) dt \right] = \frac{1}{2} \text{trace}(A|H) E[\tau_x^{(r)}].$$

It follows obviously that $\Delta u(x) = \text{trace}(A|H)$ for all x in B .

Theorem 2. *Let g be a twice continuously H -differentiable function in an open subset U of B such that D^2g is continuous from U to $\mathcal{B}_1(H, H)$. Let V be an open subset of U with positive B -distance from the complement of U . Let τ be a stopping time such that*

- (a) *almost surely, $\tau < \infty$ and $x_0 + W(t) \in V$ for all $0 \leq t \leq \tau$, and*
- (b) *$\int_1^\infty E[1_{t \leq \tau} |Dg(x_0 + W(t))|^2] dt < \infty$.*

Then

$$E[g(x_0 + W(\tau))] = g(x_0) + E \left[\int_0^\tau \frac{1}{2} \text{trace } D^2g(x_0 + W(t)) dt \right].$$

Proof. By [1, Lemma 5.4], there exists a twice H -differentiable function ϕ from B to $[0, 1]$ such that (i) $\phi \equiv 1$ in V , (ii) the support of ϕ has positive B -distance from the complement of U and (iii) $D^2\phi$ is continuous from B to $\mathcal{B}_1(H, H)$. This theorem follows by applying Theorem 1 to the function $f(x) = \phi(x)g(x)$.

Corollary 2. *Let g be a twice continuously H -differentiable function in an open subset U of B such that D^2g is continuous from U to $\mathcal{B}_1(H, H)$. Then*

$$\Delta g(x) = \text{trace } D^2g(x), \quad x \in U.$$

Proof. Let $x \in U$. Take V to be a small ball around x . Then

$$\int_1^\infty E[1_{t \leq \tau_x^{(r)}} |Dg(x_0 + W(t))|^2] dt < \infty$$

in Theorem 2 is satisfied by the same argument as in the proof of Corollary 1. Hence

$$E[g(x + W(\tau_x^{(r)}))] - g(x) = E\left[\int_0^{\tau_x^{(r)}} \frac{1}{2} \text{trace } D^2 g(x + W(t)) dt\right].$$

The desired conclusion follows from the continuity of $D^2 g$.

3. General theorem. Without much extra effort, Theorem 1 can be generalized to diffusion processes. Let $X(t)$ be the solution of the following stochastic integral equation:

$$X(t) = x + \int_0^t A(X(s)) dW(s) + \int_0^t \sigma(X(s)) ds,$$

where A and σ are given as in [3, Theorem 5.1].

Theorem 3. Let f be a twice continuously H -differentiable function in B such that $D^2 f(x) \in \mathcal{B}_1(H, H)$ for all x in B and $D^2 f$ is continuous from B to $\mathcal{B}_1(H, H)$. Let τ be a stopping time such that $\tau < \infty$ a.s. and $\int_1^\infty E[1_{t \leq \tau} |A^*(X_t) Df(X_t)|^2] dt < \infty$. Then

$$E[f(X(\tau))] = f(x) + E\left[\int_0^\tau \left\{ \langle Df(X(t)), \sigma(X(t)) \rangle + \frac{1}{2} \text{trace } A^*(X(t)) D^2 f(X(t)) A(X(t)) \right\} dt\right].$$

REFERENCES

1. C. M. Elson, *An extension of Weyl's lemma to infinite dimensions*, Trans. Amer. Math. Soc. 194 (1974), 301–324.
2. L. Gross, *Potential theory of Hilbert space*, J. Functional Analysis 1 (1967), 123–181. MR 37 #3331.
3. H.-H. Kuo, *Stochastic integrals in abstract Wiener space*, Pacific J. Math. 41 (1972), 469–483. MR 46 #5561.
4. H. P. McKean, Jr., *Stochastic integrals*, Probability and Math. Statist., no. 5, Academic Press, New York and London, 1969. MR 40 #947.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VIRGINIA 22903