CONVERGENT NETS OF PARABOLIC AND GENERALIZED SUPERPARABOLIC FUNCTIONS

NEIL A. EKLUND

ABSTRACT. The well-known convergence properties of families of harmonic functions are generalized to functions which satisfy $Lu = 0$ where L is the weak parabolic operator in divergence form. Properties of superharmonic functions are obtained for generalized superparabolic functions. These results are obtained on any bounded domain in E^{n+1} .

Consider the parabolic operator in divergence form given by

$$Lu = u_t - \{a_{ij}(x, t)u_{,i} + d_j(x, t)u_{,j} - b_j(x, t)u_{,j} - c(x, t)u.$$

In two preceding papers by the author [4], [5], existence, representation, and a maximum principle were obtained for solutions of $Lu = 0$ in the cylindrical domain $Q = \Omega \times (0, T)$ for $\Omega \subset E^n$, and generalized superparabolic functions in Q were introduced. In this article the author will consider convergent nets of parabolic and superparabolic functions on a bounded domain U assumed to be in $E^n \times (0, T)$. Since some of the properties obtained in this article will depend on results for superparabolic functions, it is necessary to restate these results for the domain U . The numbering of definitions and theorems in this article continues from those in [5].

Definition 7. Let $z = (x, t)$, $w = (y, s) \in U$. $z \prec w$ in U if there is a polygonal path $\{C_{z,w}(\alpha), 0 \leq \alpha \leq 1\}$ such that

- (1) $C_{z,w}(0) = \{z\}$, $C_{z,w}(1) = \{w\}$,
- (2) if $C_{z,w}(\alpha) = \{(\xi_\alpha, \tau_\alpha)\}$, then $\alpha < \beta$ implies $\tau_\alpha \leq \tau_\beta$,
- (3) $\{C_{z,w}(\alpha); 0 \leq \alpha \leq 1\} \subset U$.

Note that Definitions 4, 5, and 6 can be generalized to the domain U since all properties are in terms of standard rectangles in the given domain. Accordingly, let $\mathcal{L}_U$, $\mathcal{L}'_U$, $\mathcal{L}''_U$, and $\mathcal{S}''_U$ denote the corresponding spaces. The

Received by the editors April 3, 1974.

AMS (MOS) subject classifications (1970). Primary 35K10; Secondary 31C99.

Key words and phrases. Parabolic operator, sequences, generalized superparabolic functions, parabolic minorants.

theorems and corollaries stated below correspond to the analogous results obtained for Q in [5].

Theorem 4'. *If $u \in \mathcal{S}_U'''$ and if there is a $z_0 \in U$ such that $0 \geq u(z_0) = \inf_U u$, then $u(z) = u(z_0)$ for all $z \prec z_0$ in U .*

The following corollary is the analogue of that preceding Theorem 7 in [5].

Corollary. *If $u \in \mathcal{S}_U'$, then*

(1) $u(z_0) < \infty$ implies $u(z) < \infty$ for all $z \prec z_0$ in U ;

(2) $u(z_0) = \infty$ implies $u(z) = \infty$ for all $z_0 \prec z$ in U .

Theorem 10'. *Let $u \in \mathcal{S}_U'$ and let R be a standard rectangle in U . If $u < \infty$ on R , then $L(u; z, R)$ exists and the function*

$$v(z) = \begin{cases} u(z), & z \in U - R, \\ L(u; z, R), & z \in R, \end{cases}$$

satisfies $u \geq v$, $Lv = 0$ on R , and $v \in \mathcal{S}_U'$.

Lemma 2. *If $\{u_\alpha; \alpha \in A\}$ is a net of functions parabolic on U which converges uniformly on U to u , then u is parabolic on U .*

Proof. For any standard rectangle R , $\bar{R} \subset U$, and any $z \in R$, $u_\alpha(z) = L(u_\alpha; z, R)$. Since $u_\alpha \rightarrow u$ uniformly on R , it follows that $u(z) = L(u; z, R)$. Since R in U was arbitrary, u is parabolic on U .

Theorem 11. *If $\mathcal{F}$ is a family of parabolic functions on U uniformly bounded on a neighborhood of each point of U and if K is compact in U , then the family $\mathcal{F}$ is uniformly equicontinuous on K and each net $\{u_\alpha; \alpha \in A\}$ in $\mathcal{F}$ has a subnet which converges uniformly on K . If $\{u_\alpha; \alpha \in A\}$ is a convergent net of uniformly bounded parabolic functions on U , then $u \equiv \lim_A u_\alpha$ is parabolic on U .*

Proof. For each $z \in K$, let N_z denote the neighborhood of z on which the family $\mathcal{F}$ is uniformly bounded. Then $\bigcup \{N_z; z \in K\}$ is an open cover for K and, by the Heine-Borel theorem, there is a finite subcover N . Let $N = \bigcup_{j=1}^k \{N_{z_j}; z_j \in K\}$. Then the family $\mathcal{F}$ is uniformly bounded on N (and hence on K). Each member of $\mathcal{F}$ is parabolic on U and, hence, is Hölder continuous on U . Since K is compact in U , the Hölder coefficients and exponents can be made uniform for the family $\mathcal{F}$ and it follows that $\mathcal{F}$ is an equicontinuous family on K . It follows from Arzela's theorem that each net $\{u_\alpha; \alpha \in A\}$ in $\mathcal{F}$ has a uniformly convergent subnet on K .

Let $\{u_\alpha; \alpha \in K\}$ be a convergent net of uniformly bounded parabolic

functions on U . Then $u \equiv \lim_A u_\alpha$ exists. Let $\{K_j\}$ be a sequence of compact sets in U such that $K_j \uparrow U$. It follows from what has been shown that for K_1 there is a subnet $\{u_\alpha\}_1$ of $\{u_\alpha; \alpha \in A\}$ which converges to a parabolic function on K_1 . Since $u_\alpha \rightarrow u$ on U , u is parabolic on K_1 . Using what has been shown on K_2 , there is a subnet $\{u_\alpha\}_2$ of $\{u_\alpha\}_1$ which converges to a parabolic function, u , on K_2 . Continuing this process, the diagonalization process can be used to see that u is parabolic on all compact subsets of U and, hence, on U itself.

Definition 8. Let u be defined on U . Let $U_P = \{z \in U; u \text{ locally parabolic at } z\}$ and $U_\pm = \{z \in U; u(z) = \pm \infty\}$. u is said to have property $P_\pm$ if

(1) $w \in U_P$ implies $z \in U_P$ for all $z \in U$ with $z \prec w$.

(2) $w \in U_\pm$ implies $z \in U_\pm$ for all $z \in U$ with $z \succ w$.

(3) $U - U_P - U_+ - U_-$ is a finite union of sets of the form $\Omega_j \times (t_j)$.

Theorem 12. If u_j is an increasing (or decreasing) sequence of parabolic functions on U , then $u(z) = \lim u_j(z)$ satisfies property P_+ (or P_-).

Proof. Assume $u_j \uparrow u$. If for some $w \in U$, $u(w) < \infty$, then for any standard rectangle R with $\bar{R} \subset U$ and $w \in R$, the monotone convergence theorem implies that

$$(1) \quad L(u; w, R) = \lim L(u_j; w, R) = \lim u_j(w) = u(w) < \infty$$

and, hence, $u(z) < \infty$ on $\partial_P R$ with $z \prec w$. Therefore, U_P satisfies property (1) of Definition 8. Since $u \geq u_1 \geq -\infty$ by the parabolicity of u_1 , $U_- = \emptyset$. To see that U_+ satisfies property (2), suppose $u(w) = +\infty$ and suppose for some $z \in U$ with $w \prec z$ that $u(z) < \infty$. Then by our first argument $u(w) < \infty$ and a contradiction is obtained. Therefore property (2) is satisfied. Since $u_j \uparrow u$, for any $z \in U$ either $u(z) < \infty$ or $u(z) = \infty$ must hold. Hence, property (3) is vacuously satisfied.

Definition 9. A family $\mathcal{F}$ of functions defined on U is *right-directed* if for each pair $u, v \in \mathcal{F}$ there is a function $w \in \mathcal{F}$ such that $u \leq w$ and $v \leq w$. $\mathcal{F}$ is *left-directed* if the above inequalities are reversed.

Lemma 3. If $\{u_i; i \in I\}$ is a right-directed (left-directed) family of functions parabolic on U , then $u \equiv \sup_I u_i$ ($u \equiv \inf_I u_i$) satisfies property P_+ (P_-) in U .

Proof. If $\{u_i; i \in I\}$ contains only one function then we are done. If

$\{u_i; i \in I\}$ contains two or more functions, then the right-directedness can be used to obtain a sequence $\{u_j\}$ in $\{u_i; i \in I\}$ such that $u_j \uparrow u \equiv \sup_I u_i$. The desired result follows from Theorem 12.

Definition 10. A family $\mathcal{F}$ in $\mathcal{S}'_U$ is *saturated* if

- (1) $u, v \in \mathcal{F} \Rightarrow u \wedge v \in \mathcal{F}$,
- (2) $u \in \mathcal{F} \Rightarrow u^* \in \mathcal{F}$ where, for some standard rectangle R in U ,

$$(2) \quad u^*(z) = \begin{cases} u(z) & \text{in } U - R, \\ L(u; z, R) & \text{in } R. \end{cases}$$

Theorem 13. If $\mathcal{F}$ is a saturated family in $\mathcal{S}'_U$, then $v \equiv \inf_{\mathcal{F}} u$ satisfies property P_- in U .

Proof. Let R be an arbitrary standard rectangle. It will be shown that v satisfies property P_- on R and the desired result will follow from the arbitrariness of R in U .

For each $u \in \mathcal{F}$, let u^* be defined as in (2). By Theorem 10', $u^* \leq u$ on R . Moreover, since $\mathcal{F}$ is saturated in $\mathcal{S}'_U$, $u \in \mathcal{F}$ implies $u^* \in \mathcal{F}$. Therefore,

$$v = \inf \{u; u \in \mathcal{F}\} = \inf \{u^*; u \in \mathcal{F}\}.$$

Let $\mathcal{F}^* = \{u^*; u \in \mathcal{F}\}$. If it can be shown that $\mathcal{F}^*$ is a left-directed family, the desired result will follow from Lemma 3. Let $u, w \in \mathcal{F}$. Then $u \wedge w \in \mathcal{F}$ and hence u^*, w^* , and $(u \wedge w)^* \in \mathcal{F}$. However $u \wedge w \leq u$ implies $(u \wedge w)^* \leq u^*$ and similarly $(u \wedge w)^* \leq w^*$. That is, for any two elements $u^*, w^* \in \mathcal{F}^*$, there is an element, namely $(u \wedge w)^*$, in $\mathcal{F}$ such that $(u \wedge w)^* \leq u^*$ and $(u \wedge w)^* \leq w^*$. Hence, $\mathcal{F}^*$ is left-directed and the proof is complete.

The preceding results for families of parabolic and generalized super-parabolic functions will next be used to find the greatest parabolic minorant of a given function u on U . Let W be an open set in U and let $u \in \mathcal{S}'_U$. Let $\{R_j\}$ be a sequence of standard rectangles satisfying

- (1) $\bar{R}_j \subset W$ for all j ,
- (2) $W = \bigcup_{j=1}^{\infty} R_j$,
- (3) each rectangle R_j occurs infinitely often in the sequence $\{R_j\}$.

Define

$$u_1(z) = \begin{cases} u(z), & z \in U - R_1, \\ L(u; z, R_1), & z \in R_1, \end{cases}$$

and inductively

$$u_n(z) = \begin{cases} u_{n-1}(z), & z \in U - R_n, \\ L(u_{n-1}; z, R_n), & z \in R_n. \end{cases}$$

Then by Theorem 10' $u_n \leq u_{n-1}$ on U , u_n is parabolic on R_n , and $u_n \in \mathcal{S}_U^?$. Since $\{u_n\}$ is a decreasing sequence, $u_\infty(z) \equiv \lim_{n \rightarrow \infty} u_n(z)$ satisfies property P_- on U .

Definition 11. u_∞ is called the *reduction of u over W relative to U* .

It appears as if for a given open set W in U the corresponding reduction u_∞ will depend on the defining sequence $\{R_j\}$. It will follow from Theorem 15 that u_∞ is independent of the choice of $\{R_j\}$ if there is a parabolic function v on W with $v \leq u$.

Theorem 14. If $u \in \mathcal{S}_U^?$, W is open in U , and u_∞ is the reduction of u over W relative to U corresponding to the sequence $\{R_j\}$, then u_∞ satisfies property P_- on each component of W , and $u_\infty = u$ on $U - W$.

Proof. Without loss of generality assume W is connected. For each j , there is a sequence $\{j_k\}$ such that $R_{j_k} = R_j$ for all k . Thus, on R_j ,

$$u_\infty(z) = \lim_{k \rightarrow \infty} u_{j_k}(z).$$

Since u_{j_k} is parabolic on R_j for all k , it follows from Theorem 12 that u_∞ satisfies property P_- on R_j . Using this procedure on each different rectangle R_j in the defining sequence, it follows that u_∞ satisfies property P_- on W . Since $u_j \equiv u$ on $U - W$ for all j , $u_\infty = u$ on $U - W$.

Lemma 4. If $u \in \mathcal{S}_U^?$, W is open in U , v is parabolic on W , and $v \leq u$ on W , then $v \leq u_\infty \leq u$ on W where u_∞ is the reduction of u over W in U .

Proof. Let $\{R_j\}$ be the sequence of standard rectangles which define the reduction of u over W , u_∞ . Then

$$v(z) = L(v; z, R_1) \leq L(u; z, R_1) = u_1(z) \leq u(z) \quad \text{on } R_1,$$

and

$$v(z) \leq u(z) = u_1(z) \quad \text{on } W - R_1.$$

Therefore, $v \leq u_1 \leq u$ on W . Proceeding inductively, the desired result is obtained.

Definition 12. If $u \in \mathcal{S}_U^?$, v is parabolic on U , and $v \leq u$ on U , then v is called a *parabolic minorant* of u . v is the *greatest parabolic minorant* of u if v is a parabolic minorant of u and any other parabolic minorant of u is $\leq v$.

Theorem 15. *If $v \in \mathcal{S}'_U$, W is open in U , and u has a parabolic minorant v on W , then u has a unique greatest parabolic minorant on W ; namely, u_∞ .*

Proof. It follows from Lemma 4 that $v \leq u_\infty \leq u$ on W . But v parabolic on W implies $v > -\infty$ on W . Therefore, since u_∞ satisfies property P_- on W , u_∞ is parabolic on W . If there were two such u_∞ , say $u_{\infty,1}$ and $u_{\infty,2}$, then $u_{\infty,1} \leq u_{\infty,2} \leq u$ on W since $u_{\infty,1}$ is a parabolic minorant and $u_{\infty,2} \leq u_{\infty,1} \leq u$ on W since $u_{\infty,2}$ is a parabolic minorant. Therefore, u_∞ is unique and it is the greatest parabolic minorant of u .

Theorem 16. *If $u, v \in \mathcal{S}'_U$ have harmonic minorants, then $(u + v)_\infty = u_\infty + v_\infty$.*

Proof. Since u and v have harmonic minorants, $u + v$ does also and, hence, u_∞ , v_∞ , and $(u + v)_\infty$ are independent of the defining sequence $\{R_j\}$. For such a sequence and $j \geq 1$

$$\begin{aligned} (u + v)_j(x) &= \begin{cases} (u + v)_{j-1}(z) & \text{on } U - R_j, \\ L((u + v)_{j-1}; z, R_j) & \text{on } R_j, \end{cases} \\ &= \begin{cases} u_{j-1}(z) & \text{on } U - R_j \\ L(u_{j-1}; z, R_j) & \text{on } R_j \end{cases} + \begin{cases} v_{j-1}(z) & \text{on } U - R_j \\ L(v_{j-1}; z, R_j) & \text{on } R_j \end{cases} \\ &= u_j(z) + v_j(z). \end{aligned}$$

Since $u_j \downarrow u_\infty$ and $v_j \downarrow v_\infty$, it follows that $(u + v)_j \downarrow (u + v)_\infty$ and $(u + v)_\infty = u_\infty + v_\infty$.

REFERENCES

1. D. G. Aronson, *Non-negative solutions of linear parabolic equations*, Ann. Scuola Norm. Sup. Pisa (3) 22 (1968), 607–694.
2. D. G. Aronson and James Serrin, *Local behavior of solutions of quasilinear parabolic equations*, Arch. Rational Mech. Anal. 25 (1967), 81–122. MR 39 #5952.
3. Neil Eklund, *Boundary behavior of solutions of parabolic equations with discontinuous coefficients*, Bull. Amer. Math. Soc. 77 (1971), 788–792. MR 45 #2311.
4. ———, *Existence and representation of solutions of parabolic equations*, Proc. Amer. Math. Soc. 47 (1975), 137–142.
5. ———, *Generalized super-solutions of parabolic equations* (submitted).
6. Avner Friedman, *Partial differential equations of parabolic type*, Prentice-Hall, Englewood Cliffs, N. J., 1964. MR 31 #6062.

7. O. A. Ladyženskaja and N. N. Ural'ceva, *Linear and quasi-linear equations of elliptic type*, "Nauka", Moscow, 1964; English transl., Academic Press, New York, 1968. MR 35 #1955; 39 #5941.

8. Walter Rudin, *Real and complex analysis*, McGraw-Hill, New York, 1966. MR 35 #1420.

9. Neil Trudinger, *Pointwise estimates and quasilinear parabolic equations*, Comm. Pure Appl. Math. 21 (1968), 205–226. MR 37 #1758.

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37235

Current address: Department of Mathematics, Centre College, Danville, Kentucky 40422