SOMETIMES ONLY SQUARE MATRICES CAN BE DIAGONALIZED

LAWRENCE S. LEVY

ABSTRACT. It is proved that every SQUARE matrix over a serial ring is equivalent to some diagonal matrix, even though there are rectangular matrices over these rings which cannot be diagonalized.

Two matrices A and B over a ring R are *equivalent* if $B = PAQ$ for invertible matrices P and Q over R .

Following R. B. Warfield, we will call a module *serial* if its submodules are totally ordered by inclusion; and we will call a ring R *serial* if R is a direct sum of serial left modules, and also a direct sum of serial right modules. When R is artinian, these become the *generalized uniserial* rings of Nakayama [E-G].

Note that, in serial rings, finitely generated ideals do *not* have to be principal; for example, let R be any full ring of lower triangular matrices over a field. Moreover, it is well known (and easily proved) that if $aR + bR$ is a nonprincipal right ideal in *any* ring, the 1×2 matrix $[a \ b]$ is not equivalent to a diagonal matrix. Thus serial rings do not have the property that every rectangular matrix can be diagonalized.

1. Background. Serial rings are clearly *semiperfect* ($R/\text{rad}R$ is semisimple artinian and idempotents can be lifted modulo $\text{rad}R$).

(1.1) *Every indecomposable projective module over a semiperfect ring R is $\cong eR$ for some primitive idempotent e of R . (See [M].)*

A double-headed arrow $\rightarrow$ will denote an "onto" homomorphism.

By a *projective cover* of a module X we mean a *minimal epimorphism* $\phi: P \rightarrow X$ with P projective. ("Minimality" means that no submodule other than P itself is mapped *onto* X ; equivalently, $\ker \phi$ is superfluous in P .)

(1.2) *(Uniqueness of the projective cover). Let $f: P \rightarrow X$ be a module epimorphism with P projective, and suppose that X has a projective cover $\phi: P_1 \rightarrow X$. Then $P = P' \oplus P''$ where $f(P'') = 0$ and $f: P' \rightarrow X$ is a projective cover. Moreover, there is an isomorphism $\theta: P' \rightarrow P_1$ such that $(f|P') = \phi\theta$. (See [B, 2.3].)*

(1.3) *Over a semiperfect ring, every finitely generated module has a projective cover [B, 2.1].*

Received by the editors July 17, 1974.

AMS (MOS) subject classifications (1970). Primary 16A64, 16A46, 15A21, 15A33; Secondary 16A50.

Key words and phrases. Matrix, matrix equivalence, diagonal matrices, generalized uniserial ring, serial ring, projective module.

(1.4) If $\phi_i: P_i \twoheadrightarrow X_i$ is a projective cover of X_i over any ring ($i = 1, \dots, n$), then

$$\phi = \bigoplus_{i=1}^n \phi_i: \bigoplus_{i=1}^n P_i \twoheadrightarrow \bigoplus_{i=1}^n X_i$$

is a projective cover of $\bigoplus X_i$.

Proof. Since X_i is finitely generated, minimality of ϕ_i shows that P_i is finitely generated, too. Hence we easily establish: $\ker \phi_i$ is superfluous in $P_i \iff$

$$\ker \phi_i \subseteq \text{rad } P_i = \bigcap (\text{all maximal submodules of } P_i).$$

(This is more or less [BO, p. 64, Corollary 3].) Then use the fact that $\text{rad}(\bigoplus P_i) = \bigoplus \text{rad } P_i$ [BO, p. 64, Corollary 2].

2. Theorem. Let $\alpha: P \rightarrow Q$ be a homomorphism of finitely generated, projective right modules P and Q over a serial ring R . Then there exist decompositions $P = P_1 \oplus \dots \oplus P_s$ and $Q = Q_1 \oplus \dots \oplus Q_t$ with each P_i and Q_i indecomposable, such that

$$(1) \quad \alpha(P_i) \subseteq Q_i \quad \text{for } i \leq w \quad \text{and} \quad \alpha(P_i) = 0 \quad \text{for } i > w$$

where $w = \min\{s, t\}$.

Proof. This theorem is a slight extension of Warfield's theorem [W, 3.3], which states that if X is a finitely generated submodule of a finitely generated projective module Q over a serial ring R , then there is a decomposition

$$(2) \quad Q = Q_1 \oplus \dots \oplus Q_t \quad (\text{each } Q_i \text{ indecomposable})$$

such that

$$(3) \quad X = X_1 \oplus \dots \oplus X_t \quad (\text{each } X_i = X \cap Q_i).$$

Choose the numbering so that X_i is nonzero for $i \leq w$ and $X_i = 0$ for $i > w$.

Now let P be the module in our theorem, and $X = \alpha(P)$. Each X_i in (3) is finitely generated, because it is a homomorphic image of the finitely generated module X , and therefore has a projective cover $\phi_i: P_i \twoheadrightarrow X_i$. Moreover, X_i is serial ($X_i \subseteq$ the serial module Q_i) and therefore P_i is indecomposable. (If $P_i = A \oplus B$, then $X_i = \phi_i(A) + \phi_i(B)$. Since X_i is serial we can suppose $\phi_i(A) \supseteq \phi_i(B)$. Then $X_i = \phi_i(A)$ and by "minimality" of the projective cover, $A = P_i$.)

By (1.4),

$$\bigoplus_{i=1}^w \phi_i: \bigoplus_{i=1}^w P_i \twoheadrightarrow \bigoplus_{i=1}^w X_i$$

is a projective cover of X ; and by uniqueness of the projective cover (1.2), we can suppose that $\bigoplus_i P_i$ is a direct summand of P (each ϕ_i then becomes the restriction of α to P_i) with the complimentary summand being mapped by α to 0.

Thus the theorem is proved except that the w obtained above might be $< \min\{s, t\}$. But, when $i > w$ we have obtained, $\alpha(P_i) = \{0\} \subseteq Q_i$ provided, of course, that $i \leq \min\{s, t\}$. Thus we can change to $w = \min\{s, t\}$. Q.E.D.

3. Main Theorem. *Every square matrix over a serial ring is equivalent to some diagonal matrix.*

Proof. Since R is serial, it has a decomposition

$$(1) \quad R = e_1 R \oplus \dots \oplus e_d R$$

with $\{e_i\}$ an orthogonal set of primitive idempotents.

Let $\alpha: R^{(n)} \rightarrow R^{(n)}$ be left multiplication by the given $n \times n$ matrix A , elements of $R^{(n)}$ being written as columns. It will suffice to write $R^{(n)} = R'_1 \oplus \dots \oplus R'_n = R_1 \oplus \dots \oplus R_n$, with each R'_i and $R_i \cong R_R$, and each $\alpha(R'_i) \subseteq R_i$.

The first step is to write $R^{(n)}$ as a direct sum of indecomposable modules P_k and Q_k in such a way that α takes the form

$$(2) \quad \alpha: R^{(n)} = \bigoplus_{k=1}^{nd} P_k \rightarrow R^{(n)} = \bigoplus_{k=1}^{nd} Q_k \quad \text{with } \alpha(P_k) \subseteq Q_k$$

where d is the integer defined in (1). To do this, merely apply Theorem 2, with $P = Q = R^{(n)}$. To see that k runs from 1 through nd , as claimed in (2), note that the ring of endomorphisms of $e_i R$ is $e_i R e_i = \text{local}$ since R is semi-perfect [L, p. 76, Proposition 2], and therefore the Azumaya-Krull-Schmidt theorem applies to decompositions of $R^{(n)}$ (see [L, p. 78, Corollary]).

We will have our submodules R'_i and R_i if we can express the set $\{1, 2, \dots, nd\}$ as the union of n disjoint subsets $S(1), \dots, S(n)$ such that, for each b ,

$$(3) \quad \bigoplus_{k \in S(b)} P_k \cong R \quad \text{and} \quad \bigoplus_{k \in S(b)} Q_k \cong R.$$

To do this, recall that

$$\bigoplus_{k=1}^{nd} P_k = R^{(n)} = \left(\bigoplus_{i=1}^d e_i R \right)^{(n)}$$

By the Krull-Schmidt theorem, there are at least n indices k such that $P_k \cong e_1 R$. (There will be more than n such indices k if, say, $e_1 R \cong e_2 R$.) Choose n of these indices, and write $P_k \cong e_1 R$ to indicate that P_k is one of the "chosen" P 's which are $\cong e_1 R$. Then choose n different P_k 's

" $\cong$ " e_2R , and so on, until we partition $\{P_k\}$ into d disjoint n -element subsets, each associated with a distinct e_iR (and thus we exhaust the e_iR , too). Similarly designate n of the Q_k 's " $\cong$ " each e_iR .

We can keep track of all this by introducing a set of ordered triples $(k, i \rightarrow j)$, one for each $k \in \{1, 2, \dots, nd\}$, to indicate that P_k " $\cong$ " e_iR and Q_k " $\cong$ " e_jR . The arrow serves as a reminder that we are thinking of $\alpha: P_k \rightarrow Q_k$ as a map: $e_iR \rightarrow e_jR$.

Next, let M be the $d \times d$ matrix of nonnegative integers whose (i, j) entry is

$$(4) \quad m_{ij} = \text{the number of triples of the form } (k, i \rightarrow j) \quad (k \text{ varying}).$$

This matrix M has two properties which interest us. For each i , the sum of the entries $\sum_{j=1}^d m_{ij}$ in row i equals the number of P_k which are " $\cong$ " e_iR ; that is, n . Similarly the sum of the entries in each column equals n .

Now, a theorem of Birkhoff states that if a square matrix M of nonnegative real numbers has all of its row sums and column sums equal to the same number, then M is a sum $u_1M_1 + u_2M_2 + \dots$ of nonnegative multiples of permutation matrices M_b . (A $d \times d$ permutation matrix is a square matrix which can be obtained by permuting the rows of the $d \times d$ identity matrix.) In fact, one can see by looking through the proof of Birkhoff's theorem given in [H, p. 52], if the entries of M are all nonnegative integers, we can choose the multipliers u_b to be nonnegative integers, too. Writing this multiplication as repeated addition, we see that our matrix M defined in (4) is a sum of permutation matrices. Checking the row sums then shows that M is the sum of exactly n permutation matrices

$$(5) \quad M = M_1 + \dots + M_n \quad (M_b = \text{permutation matrix}).$$

Now we build the sets $S(b)$. For each ordered pair (i, j) such that $m_{ij} \neq 0$, the number of k 's which occur in a triple $(k, i \rightarrow j)$ equals m_{ij} , and this also equals the number of matrices M_b in (5) whose (i, j) entry equals 1. Therefore we can choose a 1-1 correspondence between the numbers k in the set $\{1, 2, \dots, nd\}$ and the nonzero entries of the matrices $\{M_b\}$, such that $k \leftrightarrow (i, j)_b =$ the (i, j) entry of M_b implies $P_k \cong e_iR$ and $Q_k \cong e_jR$. Then let $S(b)$ be the set of all k 's which correspond to an entry of M_b , that is, $k \leftrightarrow (,)_b$. Since a permutation matrix contains exactly one nonzero entry in each row and one in each column, (3) holds, and the proof is complete.

4. Remark. We have not dealt with uniqueness of the diagonal matrices obtained here because, as observed in [L-R, 4.3], if R is any semiperfect ring, $\text{diag}(c_1, \dots, c_n)$ is equivalent to $\text{diag}(d_1, \dots, d_n) \Leftrightarrow \bigoplus R/c_iR \cong \bigoplus R/d_iR$ (isomorphism of R -modules).

REFERENCES

- [B] H. Bass, *Finitistic dimension and a homological generalization of semi-primary rings*, Trans. Amer. Math. Soc. 95 (1960), 466–488. MR 28 #1212.
- [BO] N. Bourbaki, *Éléments de mathématique*. XXIII. Part 1. *Les structures fondamentales de l'analyse*. Livre II: *Algèbre*. Chap. 8: *Modules et anneaux semi-simples*, Actualités Sci. Indust., no. 1261, Hermann, Paris, 1958. MR 20 #4576.
- [E–G] D. Eisenbud and P. Griffith, *Serial rings*, J. Algebra 17 (1971), 389–400. MR 43 #2021.
- [H] Marshall Hall, Jr., *Combinatorial theory*, Blaisdell, Waltham, Mass., 1967. MR 37 #80.
- [L] J. Lambek, *Lectures on rings and modules*, Blaisdell, Waltham, Mass., 1966. MR 34 #5857.
- [L–R] L. S. Levy and J. C. Robson, *Matrices and pairs of modules*, J. Algebra 29 (1974), 427–454.
- [M] B. J. Muller, *On semi-perfect rings*, Illinois J. Math. 14 (1970), 464–467. MR 41 #6909.
- [W] R. B. Warfield, *Serial rings, semihereditary rings, and finitely presented modules*, J. Algebra (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706