A STURM-LIOUVILLE THEOREM FOR SOME ODD MULTIVALUED MAPS

JOÃO-PAULO DIAS AND JESÚS HERNÁNDEZ

ABSTRACT. Let $T: H \rightarrow 2^H$ be the subdifferential of a real l. s. c. convex function on an infinite dimensional, separable, real Hilbert space H . Assuming that T is odd (i.e. $T(-u) = -Tu, \forall u \in H$), $0 \in T(0)$, $(I + T)^{-1}$ is compact and $T(0)$ satisfies a geometrical condition, we prove that T has an infinite sequence $\{\lambda_n\}$ of eigenvalues such that $0 \leq \lambda_n \rightarrow +\infty$.

1. Introduction. Let H be an infinite-dimensional, separable, real Hilbert space with inner-product $(\cdot, \cdot)$ and norm $\|\cdot\|$, and let $\phi: H \rightarrow]-\infty, +\infty]$ be a lower semicontinuous (l. s. c.) convex function such that

$$(1.1) \quad \phi(u) \geq 0, \quad \forall u \in H, \quad \text{and} \quad \phi(0) = 0.$$

Let us consider the multivalued map $T = \partial\phi: H \rightarrow 2^H$ defined by

$$\partial\phi(u) = \{z \in H \mid \phi(v) - \phi(u) \leq (z, v - u), \forall v \in H\}.$$

We call T the subdifferential of ϕ . This is a maximal monotone operator (cf. [3], for example).

Let us put $\mathcal{C} = \{C \subset H \mid 0 \notin C, C \text{ is closed and symmetric}\}$.

Following [7] we define the genus of $C \in \mathcal{C}$, $\text{gen}(C)$, as the smallest integer N such that there exists an odd continuous map $f: C \rightarrow \mathbb{R}^N \setminus \{0\}$. By definition $\text{gen}(\emptyset) = 0$ and if there is no such N , $\text{gen}(C) = \infty$. We have $\text{gen}(\{x \in \mathbb{R}^N \mid \|x\| = 1\}) = N$ (cf. [7]).

If $A \subset H \setminus \{0\}$ is symmetric, we put (cf. [1]) $\gamma(A) = \sup\{\text{gen}(C) \mid C \subset A, C \text{ is a compact and symmetric}\}$.

We want to prove the following result:

Theorem 1. *Suppose that we have (1.1) and*

$$(1.2) \quad \phi(-u) = \phi(u), \quad \forall u \in H;$$

(1.3) *The operator $P = (I + T)^{-1}: H \rightarrow H$ is compact;*

(1.4) *For a given $R > 0$ we have $\gamma(\{u \in H \mid \|u\| = R, u \notin T(0)\}) = \infty$.*

Then, there exists an infinite sequence $(u_n, \lambda_n) \in H \times \mathbb{R}$, verifying

$$(1.5) \quad 0 \leq \lambda_n \rightarrow +\infty, \quad \|u_n\| = R(1 + \lambda_n)^{-1} \rightarrow 0, \quad \lambda_n u_n \in Tu_n, \\ n = 1, 2, \dots$$

Received by the editors September 23, 1974.

AMS (MOS) subject classifications (1970). Primary 47H05; Secondary 35P05.

Key words and phrases. Subdifferential, maximal monotone operator, genus of a closed symmetric subset, Sobolev spaces.

Remark. Condition (1.3) is easily verified if we have

(1.6) The set $\{u \in H \mid \|u\| \leq c, \phi(u) \leq c\}$ is compact, $\forall c > 0$. Condition (1.4) is trivially verified if $T(0)$ is bounded and so Theorem 1 extends Theorem 1 of [5] (where $T(0) = \{0\}$).

As an application of Theorem 1 let us consider $H = L^2(\Omega)$, where Ω is a bounded open subset of $\mathbb{R}^N$ with smooth boundary Γ , and let $j: \mathbb{R} \rightarrow]-\infty, +\infty]$ be a l. s. c. convex function such that

$$(1.7) \quad j(r) \geq 0, \quad \forall r \in \mathbb{R}, \quad \text{and} \quad j(0) = 0.$$

We define $\phi: H \rightarrow]-\infty, +\infty]$ by

$$(1.8) \quad \phi(u) = \begin{cases} \frac{1}{2} \int_{\Omega} |\text{grad } u|^2 dx + \int_{\Omega} j(u) dx & \text{if } u \in H_0^1(\Omega) \text{ and } j(u) \in L^1(\Omega), \\ +\infty & \text{otherwise,} \end{cases}$$

where $H_0^1(\Omega) = \{u \in H^1(\Omega) \mid u = 0 \text{ on } \Gamma\}$, $H^m(\Omega)$ is the usual Sobolev space and $\text{grad } u = (\partial u / \partial x_1, \dots, \partial u / \partial x_N)$.

It has been proved by H. Brézis (cf. [2]) that ϕ is a l. s. c. convex function which satisfies (1.1) and that

$$\partial\phi(u) = \begin{cases} -\Delta u + \beta(u) & \text{if } u \in H^2(\Omega) \cap H_0^1(\Omega) \text{ and there exists } f \in L^2(\Omega) \\ & \text{such that } f(x) \in \beta(u(x)) \text{ a.e. on } \Omega, \\ \emptyset & \text{otherwise,} \end{cases}$$

where $\beta = \partial j: \mathbb{R} \rightarrow 2^{\mathbb{R}}$.

Since the injection $H^1(\Omega) \hookrightarrow L^2(\Omega)$ is compact, the function ϕ verifies (1.6). Furthermore, if $T = \partial\phi$, we have $T(0) \equiv \beta(0)$, and so we have the following consequence of Theorem 1:

Theorem 2. Suppose that we have (1.7) and

$$(1.9) \quad j(-r) = j(r), \quad \forall r \in \mathbb{R}.$$

Then, given $R > 0$, there exists an infinite sequence $(u_n, \lambda_n) \in H \times \mathbb{R}$, verifying (1.5) for $T = \partial\phi$, ϕ defined by (1.8).

2. Proof of Theorem 1. It is well known that $P = (I + T)^{-1}$ is a single-valued map with domain H and range $D(T) = \{u \in H \mid Tu \neq \emptyset\}$, and that P is monotone and contracting (cf. [3], for example).

Furthermore, (1.1) implies that $0 \in T(0)$ and so $P(0) = 0$.

Let $G: H \rightarrow \mathbb{R}$ be the function defined by

$$G(u) = (u, Pu) - \frac{1}{2} \|Pu\|^2 - \phi(Pu).$$

The function G is Fréchet differentiable and $\nabla G(u) = Pu$, $\forall u \in H$, where $\nabla G(u)$ denotes the Fréchet derivative of G at the point u (cf. [4], [5]).

Hence G is convex, $G(0) = 0$ and $G(u) \geq 0, \forall u \in H$.

If we put $\mu = 1 + \lambda$, $v = \mu u$, we have $\lambda u \in Tu \Leftrightarrow v = \mu Pu$.

Since we have $T(0) = \{u \in H | Pu = 0\} = \{u \in H | G(u) = 0\}$, we can write

$$(2.1) \quad G(u) \neq 0 \Leftrightarrow Pu \neq 0,$$

$\{u \in S_R | G(u) \neq 0\} = \{u \in S_R | u \notin T(0)\}$, where $S_R = \{u \in H | \|u\| = R\}$.

Furthermore, the operator P is odd by (1.2) and maximal monotone.

Hence we have, by (1.3)

$$u_n \rightharpoonup u \Rightarrow Pu_n \rightharpoonup Pu,$$

where $\rightharpoonup$ denotes weak convergence in H .

We can now apply theorem A of [1] to operators $A \equiv I$ and $B \equiv P$. We conclude that, if $R > 0$ verifies (1.4), there exists an infinite sequence $(v_n, \mu_n) \in H \times \mathbb{R}$ such that

$$v_n \in S_R, \quad \beta_n = G(v_n) > 0, \quad v_n = \mu_n P v_n, \quad n = 1, 2, \dots,$$

where $\beta_n = \sup_{C \in \mathcal{C}_n} \min_{u \in C} G(u)$, $\mathcal{C}_n = \{C \subset S_R | C \text{ is compact, symmetric and } \text{gen}(C) \geq n\}$. It is easy to see (by adaptation of the proof of Lemma 4.6 of [6, Chapter VI]) that $\beta_n \xrightarrow[n]{\downarrow} 0$. Furthermore $\mu_n = R^2(Pv_n, v_n)^{-1} \geq 1, n = 1, 2, \dots$.

Let v_k be a subsequence of v_n such that $v_k \rightharpoonup v_0$. We have $G(v_0) = \lim_k G(v_k) = 0$ and so $Pv_0 = 0$ by (2.1).

Hence $(Pv_k, v_k) \xrightarrow[k]{\rightarrow} (Pv_0, v_0) = 0$ and so $\mu_n \xrightarrow[n]{\rightarrow} +\infty$.

This completes the proof of Theorem 1.

REFERENCES

1. H. Amann, *Lusternik-Schnirelman theory and nonlinear eigenvalue problems*, Math. Ann. 199 (1972), 55–72.
2. H. Brézis, *Monotonicity methods in Hilbert spaces and some applications to nonlinear partial differential equations*, Contributions to Nonlinear Functional Analysis (E. H. Zarantonello, ed.), Academic Press, New York, 1971, pp. 101–156.
3. ———, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland, Amsterdam, 1973.
4. J. P. Dias, *Variational inequalities and eigenvalue problems for nonlinear maximal monotone operators in a Hilbert space*, Amer. J. Math. (to appear).
5. ———, *Un théorème de Sturm-Liouville pour une classe d'opérateurs non linéaires maximaux monotones*, J. Math. Anal. Appl. 47 (1974), 400–405.
6. M. A. Krasnosel'skiĭ, *Topological methods in the theory of nonlinear integral equations*, GITTL, Moscow, 1956; English transl., Macmillan, New York, 1964. MR 20 #3464; 28 #2414.
7. P. H. Rabinowitz, *Some aspects of nonlinear eigenvalue problems*, Rocky Mountain J. Math. 3 (1973), 161–202. MR 47 #9383.

INSTITUTO DE FISICA E MATEMÁTICA, LISBOA, 4, PORTUGAL

ANALYSE NUMÉRIQUE, UNIVERSITÉ DE PARIS VI, 75-PARIS, 5, FRANCE