A NOTE ON A DIFFERENTIAL CONCOMITANT

P. R. EISEMAN AND A. P. STONE

ABSTRACT. If h and k are vector 1-forms, the vanishing of the concomitant $[h, k]$ is an integrability condition for certain problems on manifolds. In the case that $h = k$ the vanishing of the Nijenhuis tensor $[h, h]$ implies $d(\text{tr } h)$ is a conservation law for h , provided that $\text{tr } h$ is not constant. When the trace of h is constant, a conservation law for h exists if one can find a vector 1-form k with nonconstant trace such that $[h, k] = 0$.

1. Introduction. For any vector 1-form h there is an associated operator d_h which is an antiderivation of degree one. In the case that h is nonsingular and has vanishing Nijenhuis tensor $[h, h]$, the operator d_h satisfies a Poincaré lemma and thus provides an h -dependent version of de Rham's theorem. Such an operator is an example of a special type of derivation whose basic properties were studied in a definitive paper by A. Frölicher and A. Nijenhuis [3]. When h is the identity on 1-forms, the operator d_h reduces to the usual exterior differentiation operator d . This derivation d_h and other operators related to it may be used to give an alternative description of the differential concomitant $[h, k]$ of vector 1-forms h and k . The vanishing of this concomitant yields the main result of this paper. It is an identity which simply says that the composition $(d \circ \text{trace})$ acts as a first order derivation on the composition hk .

The derivational property of $d \circ \text{trace}$ is applicable in the study of conservation laws. With it one can establish the existence of conservation laws in a constructive manner which has the added advantage of giving a global result. Since conservation laws have usually been mathematically studied in only a local manner, global results are few. For example, in general relativity the effect of a global conservation law (Einstein's equations) on the manifold topology is of current physical interest (see [2]). Implicit in any global conservation law problem is an interaction with topology; the derivational identity yields a possible method which can be applied in such global problems.

2. Preliminaries. Let A denote the algebra of C^∞ functions on a compact, orientable, n -dimensional Riemannian manifold M without boundary.

Presented to the Society, January 15, 1974; received by the editors December 9, 1974.

AMS (MOS) subject classifications (1970). Primary 53C15; Secondary 53A55.

Key words and phrases. Vector 1-form, differential concomitant, differential form, conservation law.

Let E denote the A -module of differential 1-forms on M . A vector 1-form $h \in \text{End}_A E$ induces endomorphisms $h^{(q)} \in \text{End}_A (\wedge^q E)$ for any nonnegative integer q , and the $h^{(q)}$ are defined by setting $h^{(q)} = 0$ if $q > p$ and

$$h^{(q)}(\phi^1 \wedge \cdots \wedge \phi^p) = \frac{1}{(p-q)!q!} \sum_{\pi} |\pi| \{h\phi^{\pi(1)} \wedge \cdots \wedge h\phi^{\pi(q)}\} \\ \wedge \phi^{\pi(q+1)} \wedge \cdots \wedge \phi^{\pi(p)}$$

if $0 \leq q \leq p$, where $\phi^i \in E$, π runs through all permutations of $(1, \dots, p)$ and $|\pi|$ denotes the sign of the permutation. The transformation $h^{(0)}$ is taken to be the identity mapping on $\wedge^p E$.

In the case where $q = p \leq n$, the operator $h^{(p)}$ is locally represented by an $\binom{n}{p} \times \binom{n}{p}$ matrix $[h^{(p)}]$ relative to some local basis of p -forms. If $[h]$ denotes an $n \times n$ matrix which locally represents h , then it can be shown that

$$\det[h^{(p)}] = (\det[h])^{\binom{n-1}{p-1}}$$

and hence h is invertible on 1-forms if and only if $h^{(p)}$ is invertible on p -forms.

An alternating derivation $d_h: \wedge E \rightarrow \wedge E$ is obtained from h and an exterior derivation d by setting $d_h = h^{(1)}d - dh^{(1)}$. Thus when h is the identity, d_h reduces to d . The Nijenhuis tensor can be extended as a derivation on $\wedge E$. On p -forms one thus obtains the formula

$$[h, h] = -h^{(2)}d + d_h h^{(1)} + dh^{(2)},$$

which can be rewritten in the form

$$[h, h] = \frac{1}{2}\{d_{h^2} - (d_h)_h\}.$$

In a similar way the concomitant $[h, k]$ of vector 1-forms defined by the equation

$$[h, k] = \frac{1}{2}\{-[h^{(1)}k^{(1)} - (hk)^{(1)}]d \\ + [h^{(1)}dk^{(1)} + k^{(1)}dh^{(1)}] - [d[(hk)^{(1)} + k^{(1)}h^{(1)}]]\}$$

can be rewritten in the form

$$[h, k] = \frac{1}{2}\{d_{hk} - (d_h)_k\}.$$

3. The main identity. Let $\omega^1, \dots, \omega^n$ be a local orthonormal basis of differential 1-forms and suppose that $\omega^1 \wedge \cdots \wedge \omega^n$ agrees with an orientation of M . Suppose also that the vector 1-form h is locally specified by setting $h\omega^j = h_i^j \omega^i$, where $h_i^j \in A$ and $1 \leq i, j \leq n$, and the Einstein summation convention has been invoked. Since the orientation is specified, the Hodge star operator $*$: $\wedge^p E \rightarrow \wedge^{n-p} E$ is determined by setting

$$*(\omega^{i_1} \wedge \cdots \wedge \omega^{i_p}) = \epsilon_{i_1 \dots i_p} \omega^{j_1} \wedge \cdots \wedge \omega^{j_{n-p}}$$

where $i_1 < \dots < i_p$ and $j_1 < \dots < j_{n-p}$ are complementary sets of positive integers and $\epsilon_{i_1 \dots i_p}$ is +1 or -1 if the permutation $(i_1, \dots, i_p, j_1, \dots, j_{n-p})$ of the integers $(1, 2, \dots, n)$ is even or odd respectively. In the sequel the trace of $\mathbf{h}$ and the transpose of $\mathbf{h}$ will be denoted by $\text{tr } \mathbf{h}$ and $\mathbf{h}_t$ respectively. The following lemma is a key to establishing our identity. The proof of the lemma appears in [1].

Lemma 3.1. *For any vector 1-form $\mathbf{h}$,*

$$h^{(1)*} + *h_t^{(1)} = (\text{tr } \mathbf{h}) *.$$

Since the manifold M is assumed to be compact, an inner product on $\wedge^p E$ is defined by setting

$$(\alpha, \beta) = \int_M \alpha \wedge * \beta$$

for any p -forms α and β . With respect to this inner product the adjoint of $h^{(p)}$ is $h_t^{(p)}$ for $p = 0, 1, \dots, n$.

The codifferential δ is defined to be the adjoint of the exterior derivative d . The adjoint of $d_h = h^{(1)}d - dh^{(1)}$ is then easily seen to be $\delta_h = \delta h_t^{(1)} - h_t^{(1)}\delta$. The operator δ_h is called the $\mathbf{h}$ -codifferential. As a consequence of Stokes' theorem one obtains the relation $\delta = (-1)^{np+n+1} * d *$ on p -forms. The corresponding expression for δ_h is obtained in the following lemma.

Lemma 3.2. *If $\mathbf{h}$ is a vector 1-form, then on p -forms*

$$\delta_h = (-1)^{np+n+1} * \{d_h + d(\text{tr } \mathbf{h}) \wedge\} *.$$

Proof. On p -forms one obtains

$$\delta_h = \delta h_t^{(1)} - h_t^{(1)}\delta = (-1)^{np+n+1} \{ * d * h_t^{(1)} - h_t^{(1)} * d * \},$$

and thus Lemma 3.1 may then be applied to yield

$$\begin{aligned} \delta_h &= (-1)^{np+n+1} \{ * d [\text{tr } \mathbf{h} * - h^{(1)} *] - [* \text{tr } \mathbf{h} - * h^{(1)}] d * \} \\ &= (-1)^{np+n+1} \{ * d (\text{tr } \mathbf{h}) * - * d h^{(1)} * + * h^{(1)} d * \} \\ &= (-1)^{np+n+1} * \{ d_h + d(\text{tr } \mathbf{h}) \wedge \} *. \end{aligned}$$

The next lemma establishes a formula for $\text{adj}[\mathbf{h}, \mathbf{k}]$ for any $\mathbf{h}, \mathbf{k} \in \text{End}_A E$; this formula is analogous to the formula for δ_h in the preceding lemma.

Lemma 3.3. *For any $\mathbf{h}, \mathbf{k} \in \text{End}_A E$,*

$$\text{adj}[\mathbf{h}, \mathbf{k}] = (-1)^{np+1} * \{ [\mathbf{h}, \mathbf{k}] + \frac{1}{2} \{ d(\text{tr } \mathbf{h} \mathbf{k}) - \mathbf{h} d(\text{tr } \mathbf{k}) - \mathbf{k} d(\text{tr } \mathbf{h}) \} \wedge \} *$$

on p -forms.

Proof. Since

$$*[h, k] = \frac{1}{2} * \{d_{hk} - (d_h)_k\} = \frac{(-1)^{p(n-p)}}{2} \{ *d_{hk} * - *(d_h)_k * \} *$$

on p -forms, it suffices to compute $*d_{hk}*$ and $*(d_h)_k*$ on $(n-p)$ 1-forms. Thus, as a consequence of Lemma 3.2, one has

$$*d_{hk} * = (-1)^{np+1} \delta_{hk} - *d(\text{tr } hk) *$$

and

$$\begin{aligned} *(d_h)_k * &= * \{ k^{(1)} d_h - d_h k^{(1)} \} * \\ &= \{ \text{tr } k - k_t^{(1)} \} * d_h * - * d_h * \{ \text{tr } k - k_t^{(1)} \} \end{aligned}$$

by Lemma 3.1. The expression for $*(d_h)_k*$ then reduces to

$$*(d_h)_k * = (-1)^{np+1} (\delta_h)_k - * \{ kd(\text{tr } h) * + hd(\text{tr } k) \} *$$

and, hence,

$$\begin{aligned} *[h, k] &= (-1)^{p+1} \frac{1}{2} \{ \delta_{hk} - (\delta_h)_k \} * \\ &\quad + * \frac{1}{2} \{ kd(\text{tr } h) + hd(\text{tr } k) - d(\text{tr } hk) \} \wedge *. \end{aligned}$$

Thus

$$*[h, k] = (-1)^{p+1} \text{adj}[h, k] * + \frac{1}{2} * \{ kd(\text{tr } h) + hd(\text{tr } k) - d(\text{tr } hk) \} \wedge *$$

and the desired result is then obtained by an application of the Hodge star operator $*$, which is an isomorphism, on the right side of the above expression.

If $[h, k]$ vanishes, then Lemma 3.3 yields an identity which is the main result of this paper. The identity is given in the following proposition.

Proposition 3.4. *If $[h, k] = 0$, then $d(\text{tr } hk) = hd(\text{tr } k) + kd(\text{tr } h)$. Thus $D = d \circ \text{trace}: \text{End}_A E \rightarrow E$ is a first order derivation on compositions of vector 1-forms h and k whose differential concomitant $[h, k]$ vanishes.*

One can then study equations of the form $Dh = \alpha$ for some $\alpha \in E$ where a solution $h \in \text{End}_A E$ is desired. This equation is analogous to first order differential equations on the real line. The study of such equations will not be pursued here. However, the fact that D acts analogously to first derivatives on polynomial functions of one real variable will be shown in the following corollaries to Proposition 3.4.

Corollary 3.5. *If $[h, h] = 0$, then for any nonnegative integers i and j ,*

$$h^i d(\text{tr } h^j) + h^j d(\text{tr } h^i) = d(\text{tr } h^{i+j}).$$

Proof. The vanishing of $[h, h]$ implies the vanishing of the concomitant $[h^i, h^j]$ for any nonnegative integers i and j . Thus, as a consequence of Proposition 3.4 with h^i and h^j in place of h and k , the corollary is established.

Corollary 3.6. *If $[\mathbf{h}, \mathbf{h}] = 0$, then for any nonnegative integer i , $\mathbf{h}^i d(\text{tr } \mathbf{h}) = (i+1)^{-1} d(\text{tr } \mathbf{h}^{i+1})$.*

Proof. If $i = 1$, then $\mathbf{h}d(\text{tr } \mathbf{h}) = \frac{1}{2}d(\text{tr } \mathbf{h}^2)$ as a consequence of Corollary 3.5. If $\mathbf{h}^j d(\text{tr } \mathbf{h}) = (j+1)^{-1} d(\text{tr } \mathbf{h}^{j+1})$ for some positive integer $j > 1$, then

$$\mathbf{h}^{j+1} d(\text{tr } \mathbf{h}) = \frac{1}{j+1} \mathbf{h} d(\text{tr } \mathbf{h}^{j+1}) = \frac{1}{j+1} \{-\mathbf{h}^{j+1} d(\text{tr } \mathbf{h}) + d(\text{tr } \mathbf{h}^{j+2})\}$$

as another consequence of Corollary 3.5. Hence

$$\mathbf{h}^{j+1} d(\text{tr } \mathbf{h}) = (j+2)^{-1} d(\text{tr } \mathbf{h}^{j+2}).$$

This induction argument establishes the corollary.

Corollary 3.7. *If $[\mathbf{h}, \mathbf{h}] = 0$, and if $\mathbf{h}$ has constant trace, then the coefficients in the characteristic polynomial of $\mathbf{h}$ are constant.*

Corollary 3.7 is an immediate consequence of the fact that if $\mathbf{h}$ has constant trace, then all the positive powers of $\mathbf{h}$ also have constant trace.

In the case that the coefficients of the characteristic polynomial of a vector 1-form $\mathbf{h}$ are all constant, a result analogous to the formula $\delta^* = (-1)^{p+1} d$ on p -forms is obtained. Thus we have

Corollary 3.8. *If the coefficients of the characteristic polynomial of $\mathbf{h}$ are constant, then*

$$*[\mathbf{h}, \mathbf{h}] = (-1)^{p+1} (\text{adj}[\mathbf{h}, \mathbf{h}])^*.$$

Proof. Let $\mathbf{h}^n = a_0 I + a_1 \mathbf{h} + \dots + a_{n-1} \mathbf{h}^{n-1}$ be the characteristic equation of $\mathbf{h}$. Since $a_{n-2} = \frac{1}{2} \{\text{tr}(\mathbf{h}^2) - a_{n-1}^2\}$, and $\text{tr } \mathbf{h} = a_{n-1}$, it follows that $\text{tr}(\mathbf{h}^2)$ is constant and, hence, $\mathbf{h}d(\text{tr } \mathbf{h}) = \frac{1}{2}d(\text{tr } \mathbf{h}^2) = 0$. Thus $*[\mathbf{h}, \mathbf{h}] = (-1)^{p+1} \text{adj}[\mathbf{h}, \mathbf{h}]^*$ on p -forms as a consequence of Lemma 3.3.

This section is concluded with an alternate version of the identity in Corollary 3.6. Since the traces of the various powers of $\mathbf{h}$ are expressible in terms of the coefficients a_i , the following proposition may be obtained.

Proposition 3.9. *If $[\mathbf{h}, \mathbf{h}] = 0$ and $\mathbf{h}^n = a_0 I + a_1 \mathbf{h} + \dots + a_{n-1} \mathbf{h}^{n-1}$ is the characteristic equation of $\mathbf{h}$, then*

$$\mathbf{h} da_i = da_{i-1} + a_i da_n, \quad i = 0, 1, \dots, n-1 \quad (da_{-1} \equiv 0).$$

Proof. For the case that $\mathbf{h}$ is cyclic, a proof appears in [6]. Otherwise the identity in Corollary 3.6 may be used. Thus $\mathbf{h}d(\text{tr } \mathbf{h}) = \frac{1}{2}d(\text{tr } \mathbf{h}^2)$ and, since $\text{tr } \mathbf{h} = a_{n-1}$ and $\text{tr } \mathbf{h}^2 = 2a_{n-2} + a_{n-1}^2$, one obtains $\mathbf{h}da_{n-1} = da_{n-2} + a_{n-1} da_{n-1}$. Hence, the case $i = n-1$ in the proposition is established. If $\mathbf{h}$ is applied to the above result, the relation obtained is

$$\begin{aligned} \mathbf{h} da_{n-2} &= \mathbf{h}^2 da_{n-1} - a_{n-1} \mathbf{h} da_{n-1} \\ &= \mathbf{h}^2 d(\text{tr } \mathbf{h}) - a_{n-1} \mathbf{h} d(\text{tr } \mathbf{h}) = (1/3) d(\text{tr } \mathbf{h}^3) - (a_{n-1}/2) d(\text{tr } \mathbf{h}^2); \end{aligned}$$

since $\text{tr } h^3 = 3a_{n-3} + 3a_{n-1}a_{n-2} + a_{n-1}^3$, the case $i = n - 2$ is established. Thus

$$hda_{n-2} = da_{n-3} + a_{n-2}da_{n-1}.$$

The remaining identities can be established by repeated use of Corollary 3.6, though the calculation is tedious because of the form of the expressions for $\text{tr } h^i$.

Corollary 3.10. *If the Nijenhuis tensor $[h, h]$ vanishes identically, then the coefficients a_i in the characteristic equation for h are functionally independent if and only h is cyclic with generator $d(\text{tr } h)$.*

Proof. It is sufficient to establish the relation

$$d(\text{tr } h) \wedge h d(\text{tr } h) \wedge \cdots \wedge h^{n-1} d(\text{tr } h) = da_{n-1} \wedge da_{n-2} \wedge \cdots \wedge da_0,$$

which is a consequence of the proposition.

4. Conservation laws. The conservation laws of physics can be expressed in conservation law form as a differential equation [4], and also as a relation between vector 1-forms and differential forms [5]. The use of vector 1-forms and differential forms trivially extends the problem to a global one on manifolds. Global conservation laws are of interest in physics as is evident from [2], but to date do not appear to have been studied in a general mathematical context. With the identity stated in Corollary 3.4 the global conservation law problem can be studied.

On a differentiable manifold a conservation law for a vector 1-form h is defined as any exact differential form θ such that $h\theta$ is also exact. Thus if $\theta = df$ for some differentiable function f , then θ is a conservation law for h if there is a differentiable function g such that $hdf = dg$. The analysis may be divided into two cases insofar as the existence of conservation laws is concerned. The cases correspond to constant and nonconstant trace for h .

If h has trace which is not constant, then Corollary 3.6 says that $d(\text{tr } h)$ is a conservation law for any positive power of h . Moreover if h is cyclic with $d(\text{tr } h)$ as a generator, then Corollary 3.1 implies that a basis of conservation laws can be obtained. This discussion is summarized in the following statement.

Proposition 4.1. *Let M be a compact orientable n -dimensional Riemannian manifold without boundary and let h be a vector 1-form whose trace is not constant on M . If the concomitant $[h, h]$ vanishes on M , then $d(\text{tr } h^i)$ with $i = 1, 2, \dots$ is a conservation law for h , and for fixed i , $d(\text{tr } h^i)$ is a conservation law for h^j , with $j = 1, 2, \dots$. Finally if h is cyclic with generator $d(\text{tr } h)$, then the collection $d(\text{tr } h^i)$, $i = 1, 2, \dots, n$, is a basis of conservation laws for h .*

The analysis for the case in which $\text{tr } h$ is constant is not as simple.

The above reasoning can be extended provided that a nontrivial solution can be obtained for a certain system of differential equations. The solution of global differential equations is, of course, limited by topological considerations. In particular, if it is possible to find another vector 1-form $\mathbf{k}$ with non-constant trace such that the concomitant $[\mathbf{h}, \mathbf{k}]$ vanishes, then the identity of Proposition 3.4 yields the statement that $d(\text{tr } \mathbf{h})$ is a conservation law for $\mathbf{h}$. In local coordinates the vanishing of $[\mathbf{h}, \mathbf{k}]$ leads to a system of $n\binom{n}{2}$ equations (corresponding to n basis elements and $\binom{n}{2}$ elements in $E \wedge E$) in n^2 unknowns. This system may be written out explicitly if we assume that $(dx^1, \dots, dx^n)$ is a local basis of 1-forms, that $\mathbf{h}dx^i = h_\alpha^i dx^\alpha$, $\mathbf{k}dx^i = k_\alpha^i dx^\alpha$, and that $(\ , \)_{,i} = \partial(\)/\partial x^i$. Thus the condition $[\mathbf{h}, \mathbf{k}] = 0$ leads to a local system of the form

$$\begin{aligned} 2(h_j^\alpha \delta_l^\beta - h_l^\beta \delta_j^\alpha) k_{\beta, \alpha}^i \\ = (2h_{l,j}^\alpha \delta_\beta^i + h_{j,\beta}^i \delta_l^\alpha + h_{\beta,l}^i \delta_j^\alpha - 2h_{j,l}^\alpha \delta_\beta^i - h_{l,\beta}^i \delta_j^\alpha - h_{s,j}^i \delta_l^\alpha) k_\alpha^\beta \end{aligned}$$

for $1 \leq j < l \leq n$ and $1 \leq i \leq n$. For $n \geq 3$ the system $[\mathbf{h}, \mathbf{k}] = 0$ is generally overdetermined but may still have nontrivial solutions. To summarize this section one has

Proposition 4.2. *Let M be a compact orientable n -dimensional Riemannian manifold without boundary and let $\mathbf{h}$ be a vector 1-form on M . If there is a vector 1-form $\mathbf{k}$ with nonconstant trace and vanishing concomitant $[\mathbf{h}, \mathbf{k}]$, then $d(\text{tr } \mathbf{k})$ is a conservation law for $\mathbf{h}$, provided $\text{tr } \mathbf{h}$ is constant.*

REFERENCES

1. P. R. Eiseman and A. P. Stone, *A generalized Hodge theory*, J. Differential Geometry 9 (1974), 169–176. MR 49 #1534.
2. R. H. Gowdy, *Vacuum spacetimes with two-parameter spacelike isometry groups and compact invariant hypersurfaces: topologies and boundary conditions*, Ann. Phys. 83 (1974), 208–241.
3. A. Frölicher and A. Nijenhuis, *Theory of vector-valued differential forms. I. Derivations in the graded ring of differential forms*, Nederl. Akad. Wetensch. Proc. Ser. A 59 = Indag. Math. 18 (1956), 338–359. MR 18, 569.
4. P. D. Lax, *Hyperbolic systems of conservation laws. II*, Comm. Pure Appl. Math. 10 (1957), 537–566. MR 20 #176.
5. A. P. Stone, *Analytic conservation laws*, Ann. Inst. Fourier (Grenoble) 16 (1966), fasc. 2, 319–327. MR 35 #6160.
6. ———, *On the integrability of a vector 1-form*, J. London Math. Soc. (2) 6 (1973), 403–411. MR 47 #7650.

UNITED TECHNOLOGIES RESEARCH CENTER, EAST HARTFORD, CONNECTICUT 06108

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NEW MEXICO 87131