ON THE LEFSCHETZ NUMBER FOR ITERATES OF CONTINUOUS MAPPINGS¹

H. O. PEITGEN

ABSTRACT. We give an elementary proof of the (mod p)-congruence between the Lefschetz numbers of a continuous mapping and its p th iterate, p prime, and deduce some results about periodic mappings.

The following proposition can be obtained in a less general situation from theorems of Zabreiko and Krasnosel'skiĭ [10] and Steinlein [9] by using the normalization property of the fixed point index. However, the proofs in [9] and [10] are analytical and quite difficult. Moreover, the normalization property requires some deeper considerations (cf. Dold [1]). In this paper we present a short and appropriate proof using only basic algebraic and topological arguments.

PROPOSITION. *Let X be a compact Hausdorff space such that $\check{H}^*(X; \mathbb{Z})$ is finitely generated. Then $\Lambda(f) \equiv \Lambda(f^p) \pmod{p}$, whenever f is a continuous map on X and p is prime.*

For the proof of the proposition we need the following basic

LEMMA. *Let A be a matrix of integers and let p be prime. Then $\text{trace } A \equiv \text{trace } A^p \pmod{p}$.*

PROOF. If A is in $\mathbb{Z}^{n^2}$, we denote by $\Pi : \mathbb{Z}^{n^2} \rightarrow \mathbb{Z}_p^{n^2}$ and by $\pi : \mathbb{Z} \rightarrow \mathbb{Z}_p$ the corresponding projections. We have $\Pi(A^p) = (\Pi(A))^p$ and $\pi(\text{trace } A) = \text{trace } \Pi(A)$. Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of $\Pi(A)$. Since the λ_i are roots of a polynomial with coefficients in $\mathbb{Z}_p$, then $\sum_{i=1}^n \lambda_i$ is defined in the extension field $\mathbb{Z}_p(\lambda_1, \dots, \lambda_n)$ of $\mathbb{Z}_p$. Notice that $\mathbb{Z}_p(\lambda_1, \dots, \lambda_n)$ is a field of characteristic p , hence $(\sum_{i=1}^n \lambda_i)^p = \sum_{i=1}^n \lambda_i^p$. Since $\text{trace } \Pi(A) = \sum \lambda_i$ and $\text{trace } (\Pi(A))^p = \sum \lambda_i^p$ are in $\mathbb{Z}_p$, we have $\sum \lambda_i = (\sum \lambda_i)^p = \sum \lambda_i^p$ in $\mathbb{Z}_p$. Thus we get

$$\pi(\text{trace } A) = \text{trace } \Pi(A) = \text{trace } (\Pi(A))^p = \pi(\text{trace } A^p). \quad \square$$

PROOF OF PROPOSITION. From Spanier [8, p. 335], we have that the following diagram with exact rows commutes:

$$\begin{array}{ccccccc} 0 & \rightarrow & \check{H}^n(X; \mathbb{Z}) \otimes \mathbb{Q} & \xrightarrow{j} & \check{H}^n(X; \mathbb{Q}) & \rightarrow & \text{Tor}(\check{H}^{n+1}(X; \mathbb{Z}), \mathbb{Q}) \\ & & \downarrow \mathbb{Z} f_n^* \otimes \text{id}_{\mathbb{Q}} & & \downarrow \mathbb{Q} f_n^* & & \downarrow \\ 0 & \rightarrow & \check{H}^n(X; \mathbb{Z}) \otimes \mathbb{Q} & \xrightarrow{j} & \check{H}^n(X; \mathbb{Q}) & \rightarrow & \text{Tor}(\check{H}^{n+1}(X; \mathbb{Z}), \mathbb{Q}) \end{array}$$

Received by the editors August 13, 1974.

AMS (MOS) subject classifications (1970). Primary 55C20; Secondary 55C25, 55C35.

Key words and phrases. Lefschetz number, iterates, fixed point index, periodic mappings.

¹ The contents of this paper were presented at the meeting on Problems in Nonlinear Functional Analysis held in Bonn, July 22-26, 1974.

Since $\text{Tor}(A, \mathbf{Q}) = 0$ for any abelian group A , we have that J is an isomorphism and, therefore,

$$\text{trace}(\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}}) = \text{trace}(J^{-1} \circ \mathbf{Q}f_n^* \circ J) = \text{trace}(\mathbf{Q}f_n^*).$$

Let F be a free subgroup and T the torsion subgroup of $\check{H}^n(X; \mathbf{Z})$ such that $\check{H}^n(X; \mathbf{Z}) = F \oplus T$, and let $e_1, \dots, e_r$ be a basis for F . Then a basis for $\check{H}^n(X; \mathbf{Z}) \otimes \mathbf{Q}$ is given by $e_1 \otimes 1, \dots, e_r \otimes 1$. If we choose a_{ij} in $\mathbf{Z}$ and t in T such that $\mathbf{Z}f_n^*(e_i) = \sum_{j=1}^r a_{ij}e_j + t$, then we have

$$\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}}(e_i \otimes 1) = \sum_{j=1}^r a_{ij}e_j \otimes 1.$$

Hence $\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}}$ is given by a matrix of integers. Now

$$\Lambda(f) = \sum_n (-1)^n \text{trace } \mathbf{Q}f_n^* = \sum_n (-1)^n \text{trace } \mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}}.$$

From the Lemma we have

$$\text{trace}(\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}}) \equiv \text{trace}(\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}})^p \pmod{p}.$$

Finally,

$$\begin{aligned} \text{trace}(\mathbf{Z}f_n^* \otimes \text{id}_{\mathbf{Q}})^p &= \text{trace}(J^{-1} \circ \mathbf{Q}f_n^* \circ J)^p \\ &= \text{trace}(J^{-1} \circ (\mathbf{Q}f_n^*)^p \circ J) = \text{trace}(\mathbf{Q}f_n^*)^p, \end{aligned}$$

which proves the proposition. $\square$

It can be easily seen that the same proof works in the case of multivalued mappings in the sense of Powers [7].

Our interest in the proposition arose from the attempt of proving the (mod p)-theorem of Zabreiko and Krasnosel'skiĭ [10] and Steinlein [9] by purely topological means and from its usefulness in asymptotic fixed point theory (cf. Fenske and Peitgen [2] and Peitgen [6]).

We close with a few observations and remarks.

(1) The following is a generalization of results due to Floyd [3], [4] and Halpern [5]. The proofs in [3] and [4] use the Smith special homology groups, and the proof in [5] is based on the Kelley-Spanier characteristic.

THEOREM. *Let X be a compact Lefschetz space such that $\check{H}^*(X; \mathbf{Z})$ is finitely generated and let T be a fixed point free periodic transformation of period $s = p^r$, p prime, operating on X . Then p divides the Euler characteristic $\chi(X)$.*

PROOF. From the proposition we have that

$$\chi(X) = \Lambda(T^{p^r}) \equiv \Lambda(T^{p^{r-1}}) \equiv \dots \equiv \Lambda(T^p) \equiv \Lambda(T) = 0,$$

where all congruences are mod p . $\square$

We add in passing that the same argument yields the divisibility result in the case when T is a fixed point free transformation on X and $\mathbf{Q}(T^s)^* = \text{id}$.

COROLLARY . *Let X be a compact Lefschetz space such that $\check{H}^*(X; \mathbf{Z})$ is finitely generated, and let T be a periodic transformation of period $s = p^r$, p prime, operating on X . Assume that $\chi(X) \not\equiv 0 \pmod{p}$; then T has a fixed point.*

(2) Set $\lambda_p = p^{-1}(\Lambda(g^p) - \Lambda(g))$. The proposition tells us that λ_p is an integer invariant for g . It seems to be of some interest to determine λ_p especially in view of the following observations. Set

$$\chi_p(X) = \sum_n (-1)^n \dim_{\mathbf{Z}_p} \check{H}^n(X; \mathbf{Z}_p),$$

whenever this number is defined. Assume that X is a finite-dimensional compact Hausdorff space such that $\check{H}^*(X; \mathbf{Z})$ is finitely generated. Notice that a standard argument using Theorem 10 in Spanier [8, p. 335], yields $\chi(X) = \chi_p(X)$. Consider the very special case when g is a fixed point free transformation of prime period p operating on X . Then we have that $\lambda_p = \chi_p(X/G)$. Here G is the group generated by g and X/G is the orbit space. This follows from Floyd [3], [4]. Furthermore, combining another result of Floyd [3] and the proposition, we have that

$$\Lambda(g) \equiv \Lambda(g^p) = \chi(X) \equiv \chi_p(X^G) \pmod{p}$$

when g is not necessarily fixed point free.

We would like to remark that a purely topological approach for a proof of the (mod p)-theorem for the fixed point index is the following.

Let $f: U \rightarrow \mathbf{R}^n$ be continuous, where U is open in $\mathbf{R}^n$. Let p be prime and denote by F (resp. F_p) the fixed point set of f (resp. f^p) in U . Assume that F and F_p are compact and that f^p has no fixed points on the boundary of U and $f(F_p) \subset U$. Then the Krasnosel'skiĭ-Zabreiko-Steinlein theorem is equivalent to the commutativity of the following diagram:

$$\begin{array}{ccccc} H_n(S^n; \mathbf{Z}_p) & \longrightarrow & H_n(S^n, S^n - F; \mathbf{Z}_p) & \longrightarrow & H_n(U, U - F; \mathbf{Z}_p) \\ \downarrow & & & & \xrightarrow{(i-f)^*} H_n(\mathbf{R}^n, \mathbf{R}^n - \{0\}; \mathbf{Z}_p) \\ H_n(S^n; \mathbf{Z}_p) & \longrightarrow & H_n(S^n, S^n - F_p; \mathbf{Z}_p) & \longrightarrow & H_n(U, U - F_p; \mathbf{Z}_p) \\ & & & & \downarrow \\ & & & & \xrightarrow{(i-f^p)^*} H_n(\mathbf{R}^n, \mathbf{R}^n - \{0\}; \mathbf{Z}_p) \end{array}$$

Using coefficients in $\mathbf{Z}$ instead of $\mathbf{Z}_p$, the rows define a fixed point index for f (resp. f^p) (cf. Dold [1]). It is immediate from Dold's paper that it suffices to prove commutativity of the right rectangle where F is replaced by F_p .

REFERENCES

1. A. Dold, *Fixed point index and fixed point theorems for Euclidean neighborhood retracts*, Topology **4** (1965), 1–8. MR 33 #1850.
2. C. Fenske and H. O. Peitgen, *Repulsive fixed points of multivalued transformations and the fixed point index*, Math. Ann. (to appear).

3. E. E. Floyd, *On periodic maps and the Euler characteristics of associated spaces*, Trans. Amer. Math. Soc. **72** (1952), 138—147. MR **13**, 673.
4. ———, *Periodic maps via Smith theory*, Seminar on Transformation Groups, Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N.J., 1960, Chap. III.
5. B. R. Halpern, *Fixed points for iterates*, Pacific J. Math. **25** (1968), 255—275. MR **41** #1039.
6. H. O. Peitgen, *Asymptotic fixed point theorems and stability*, J. Math. Anal. Appl. **47** (1974), 32—42.
7. M. J. Powers, *Lefschetz fixed point theorems for a new class of multivalued maps*, Pacific J. Math. **42** (1972), 211—220.
8. E. H. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966. MR **35** #1007.
9. H. Steinlein, *Ein Satz über den Leray-Schauderschen Abbildungsgrad*, Math. Z. **126** (1972), 176—208. MR **47** #5667.
10. P. P. Zabreiko and M. A. Krasnosel'skiĭ, *Iterations of operators and fixed points*, Dokl. Akad. Nauk SSSR **196** (1971), 1006—1009 = Soviet Math. Dokl. **12** (1971), 294—298. MR **47** #4082.

INSTITUT FÜR ANGEWANDTE MATHEMATIK, UNIVERSITÄT BONN, BONN, FEDERAL REPUBLIC OF GERMANY