RATIONAL CHEBYSHEV APPROXIMATION TO CERTAIN ENTIRE FUNCTIONS OF ZERO ORDER ON THE POSITIVE REAL AXIS. II

A. R. REDDY

ABSTRACT. A sufficient condition is given on the growth of certain entire functions which guarantee a certain rate of convergence of the error in approximating reciprocals of a class of entire functions by reciprocals of polynomials under the uniform norm on the positive real axis.

Recently we proved

THEOREM 1 [1, THEOREM 7]. *Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$, $a_0 > 0$ and $a_k \geq 0$ ($k \geq 1$) be an entire function of zero order satisfying the assumptions that*

$$(1) \quad 1 < \limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \Lambda + 1 = \rho_l < \infty$$

and

$$(2) \quad 0 < \liminf_{r \rightarrow \infty} \frac{\log M(r)}{(\log r)^{\Lambda+1}} = b_l, \quad \limsup_{r \rightarrow \infty} \frac{\log M(r)}{(\log r)^{\Lambda+1}} = B_l < \infty,$$

where $M(r) = \max_{|z|=r} |f(z)|$. Then there exists a sequence of polynomials $\{P_n(x)\}_{n=0}^{\infty}$ for which

$$(3) \quad \lim_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{1}{P_n(x)} \right\|_{L_{\infty}[0, \infty)} \right\}^{1/n} = 0.$$

One sees that (2) implies the limit in (1), for suppose, to the contrary,

$$\begin{aligned} 0 &< \liminf_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \sigma < \Lambda + 1 \\ &= \limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} < \infty. \end{aligned}$$

Then for each $\epsilon > 0$, there is a sequence of values of r for which

$$(4) \quad \log M(r) \leq (\log r)^{\sigma+\epsilon}.$$

From (4) we obtain

Received by the editors August 27, 1974.

AMS (MOS) subject classifications (1970). Primary 41A20.

Key words and phrases. Rational approximation, zero order, positive real axis.

$$(\log M(r))/(\log r)^{\Lambda+1} \leq (\log r)^{\sigma+\epsilon}/(\log r)^{\Lambda+1},$$

and we may choose ϵ so that $\sigma + \epsilon < \Lambda + 1$. Then clearly we have

$$\liminf_{r \rightarrow \infty} \frac{\log M(r)}{(\log r)^{\Lambda+1}} = b_l = 0,$$

which contradicts (2).

On the other hand the limit in (1) may not imply (2). For example let

$$f_1(z) = 1 + \sum_{n=2}^{\infty} \frac{z^n}{\exp[(n \log n)^2]},$$

$$f_2(z) = 1 + \sum_{n=2}^{\infty} \frac{z^n}{\exp[(n/\log n)^2]}.$$

It is easy to calculate for

$$f_1(z), \quad \Lambda = 1 = \sigma - 1, \quad B_l = 0, \quad f_2(z), \quad \Lambda = 1 = \sigma - 1, \quad B_l = \infty.$$

Hence it is natural to ask whether one can get (3) under the only assumption

$$(5) \quad 0 < \lim_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \Lambda + 1 < \infty.$$

In fact we prove the more general

THEOREM 2. *Let $f(z) = \sum_{k=0}^{\infty} a_k z^k$, $a_0 > 0$, $a_k \geq 0$ ($k \geq 1$) be an entire function of zero order satisfying the assumption that*

$$0 < \Lambda < \liminf_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \sigma < \infty.$$

Then there exists a sequence of polynomials $\{P_n(x)\}_{n=0}^{\infty}$ for which

$$\lim_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{1}{P_n(x)} \right\|_{L_{\infty}[0, \infty)} \right\}^{1/n} = 0.$$

REMARK. The method used here is much simpler than the one used in [1].

PROOF. By assumption

$$\limsup_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \Lambda + 1.$$

Then it is known [2, Theorems 1 and 3] that

$$(6) \quad \limsup_{n \rightarrow \infty} \frac{\log n}{\log \{n^{-1} \log |a_n|^{-1}\}} = \Lambda.$$

From (6) it is easy to get for each $\epsilon > 0$ there is an n_0 depending on ϵ such that for all $n \geq n_0(\epsilon)$,

$$(7) \quad |a_n| \leq \exp(-n^{1+1/(\Lambda+\epsilon)}).$$

Let $S_n(z) = \sum_{k=0}^n a_k z^k$; then clearly for $0 \leq x \leq r$,

$$(8) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq \frac{f(x) - S_n(x)}{f(x)S_n(x)} \leq a_0^{-2} \sum_{k=n+1}^{\infty} a_k r^k.$$

From (7) and (8) we get, for $0 \leq x \leq r$,

$$(9) \quad \begin{aligned} 0 &\leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq a_0^{-2} \sum_{k=n+1}^{\infty} \exp(-k^{1/(1+(\Lambda+\epsilon))} + k \log r) \\ &= a_0^{-2} \sum_{k=n+1}^{\infty} \exp[(-n^{1/(\Lambda+\epsilon)}) + \log r)k]. \end{aligned}$$

Choose

$$(10) \quad 2 \log r = n^{1/(\Lambda+\epsilon)}.$$

Then we have from (9) that

$$(11) \quad \begin{aligned} 0 &\leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq a_0^{-2} \sum_{k=\Lambda+1}^{\infty} \exp\left[\left(\frac{-n^{1/(\Lambda+\epsilon)}}{2}\right)k\right] \\ &\leq C_1 \exp\left(-n^{1+1/(\Lambda+\epsilon)} C_2\right), \end{aligned}$$

where C_1, C_2 are some constants. On the other hand for all $x \geq r$, we get

$$(12) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq \frac{1}{S_n(x)} \leq \frac{1}{S_n(r)} = \frac{1}{f(r) - \sum_{k=n+1}^{\infty} a_k r^k}.$$

By assumption

$$0 < \liminf_{r \rightarrow \infty} \frac{\log \log M(r)}{\log \log r} = \sigma < \infty.$$

From this we get for all $r \geq r_0(\epsilon)$,

$$(13) \quad f(r) = M(r) \geq \exp((\log r)^{\sigma-\epsilon}).$$

On the other hand we know from (8) and (11) that

$$(14) \quad \sum_{k=n+1}^{\infty} a_k r^k \leq \exp(-C_2 n^{1/(\Lambda+\epsilon)}).$$

Hence from (12), (13) and (14) we get, with a little manipulation, for all $x \geq r$,

$$(15) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq \frac{1}{S_n(r)} \leq \exp\left(-\frac{(\log r)^{\sigma-\epsilon}}{2}\right).$$

From (15) and (10) we obtain for all $x \geq r$,

$$(16) \quad 0 \leq \frac{1}{S_n(x)} - \frac{1}{f(x)} \leq \frac{1}{S_n(r)} \leq \exp(-C_3 (n)^{(\sigma-\epsilon)/(\Lambda+\epsilon)}),$$

where C_3 is a suitable constant. By choosing ϵ such that $\sigma - \epsilon > \Lambda + \epsilon$, we

get from (11) and (16) the required result, i.e.

$$\lim_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{1}{P_n(x)} \right\|_{L_\infty[0, \infty)} \right\}^{1/n} = 0.$$

I would like to thank the referee for his suggestions.

REFERENCES

1. G. Meinardus, A. R. Reddy, G. D. Taylor and R. S. Varga, *Converse theorems and extensions in Chebyshev rational approximation to certain entire functions in $[0, +\infty)$* , Trans. Amer. Math. Soc. **170** (1972), 171—185. MR **46** #9603.
2. A. R. Reddy, *On entire Dirichlet series of zero order*, Tôhoku Math. J. (2) **18** (1966), 144—155. MR **35** #5614.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48823

Current address: Department of Mathematics, Yeshiva University, New York, New York 10033