SUBMANIFOLDS OF A RIEMANNIAN MANIFOLD WITH SEMISYMMETRIC METRIC CONNECTIONS

ZENSHO NAKAO

ABSTRACT. We derive the Gauss curvature equation and the Codazzi-Mainardi equation with respect to a semisymmetric metric connection on a Riemannian manifold and the induced one on a submanifold. We then generalize the theorema egregium of Gauss.

1. Introduction. K. Yano [6] proved that a Riemannian manifold admits a semisymmetric metric connection with vanishing curvature tensor if and only if the manifold is conformally flat. Later, T. Imai [3], [4] studied some properties of hypersurfaces of a Riemannian manifold with a semisymmetric metric connection, and also obtained the Gauss curvature equation and the Codazzi-Mainardi equation with respect to a semisymmetric metric connection on a Riemannian manifold and the induced one on a hypersurface. The object of this paper is to derive the above two equations with respect to a semisymmetric metric connection on an $(n + p)$ -dimensional Riemannian manifold and the induced one on an n -dimensional submanifold, and also to generalize the theorema egregium of Gauss. The notation of [5] will be used for the most part.

2. Gauss equation and Weingarten equation. Let M be an n -dimensional Riemannian manifold isometrically imbedded in an $(n + p)$ -dimensional Riemannian manifold M' . We denote by g the Riemannian metric tensor on M' as well as the induced one on M . Since M has codimension p we can locally choose p cross sections ξ_i , $i = 1, 2, \dots, p$, of the normal bundle $T(M)^\perp$ of M in M' which are orthonormal at each point of M .

A linear connection $\hat{\nabla}'$ on M' is called a semisymmetric metric connection if $\hat{\nabla}'g = 0$ (metric) and the torsion tensor $\hat{T}'$ of $\hat{\nabla}'$ satisfies $\hat{T}'(X', Y') = \pi(Y')X' - \pi(X')Y'$ (semisymmetric) for $X', Y' \in \mathfrak{X}(M')$, where π is a 1-form on M' [6].

We now assume that a semisymmetric metric connection ∇' is given on M' by

$$(1) \quad \hat{\nabla}'_{X'} Y' = \nabla'_{X'} Y' + \pi(Y')X' - g(X', Y')P'$$

for $X', Y' \in \mathfrak{X}(M')$, where ∇' denotes the Riemannian connection with respect to g and P' a vector field on M' defined by $g(P', X') = \pi(X')$ for X'

Received by the editors August 27, 1973 and, in revised form, December 30, 1974.

AMS (MOS) subject classifications (1970). Primary 53B15, 53B25; Secondary 53C05.

Key words and phrases. Codazzi-Mainardi equation, conformally flat space, Gauss curvature equation, mean curvature normal, sectional curvature, semisymmetric metric connection, totally umbilical submanifold.

$\in \mathfrak{X}(M')$ [6]. On M we define a vector field P and real-valued functions λ_i , $i = 1, 2, \dots, p$, by decomposing P' into its unique tangential and normal components, thus

$$(2) \quad P' = P + \sum_{i=1}^p \lambda_i \xi_i.$$

If we denote by ∇ the induced Riemannian connection on M from ∇' on M' , then we have the Gauss equation (with respect to ∇'),

$$(3) \quad \nabla'_X Y = \nabla_X Y + \sum_{i=1}^p h_i(X, Y) \xi_i$$

for $X, Y \in \mathfrak{X}(M)$, where h_i are the second fundamental forms on M [5, pp. 10–21]. Let a connection $\hat{\nabla}$ on M be induced from the semisymmetric metric connection $\hat{\nabla}'$ on M' by the equation which may be called the Gauss equation with respect to $\hat{\nabla}'$,

$$(4) \quad \hat{\nabla}'_X Y = \hat{\nabla}_X Y + \sum_{i=1}^p \hat{h}_i(X, Y) \xi_i$$

for $X, Y \in \mathfrak{X}(M)$, where $\hat{h}_i$ are tensors of type (0,2) on M .

From (1), using (2), (3) and (4), we obtain

$$(5) \quad \begin{aligned} \hat{\nabla}_X Y + \sum_{i=1}^p \hat{h}_i(X, Y) \xi_i &= \nabla_X Y + \sum_{i=1}^p h_i(X, Y) \xi_i \\ &\quad + \pi(Y)X - g(X, Y) \left(P + \sum_{i=1}^p \lambda_i \xi_i \right), \end{aligned}$$

from which we get

$$(6) \quad \hat{\nabla}_X Y = \nabla_X Y + \pi(Y)X - g(X, Y)P$$

for $X, Y \in \mathfrak{X}(M)$, and we also have

$$(7) \quad \hat{h}_i = h_i - \lambda_i g.$$

Using (6), we get $\hat{\nabla}_X \{g(Y, Z)\} = (\hat{\nabla}_X g)(Y, Z) + \nabla_X \{g(Y, Z)\}$, from which follows $(\hat{\nabla}_X g)(Y, Z) = 0$ for $X, Y, Z \in \mathfrak{X}(M)$, i.e.,

$$(8) \quad \hat{\nabla} g = 0,$$

and

$$(9) \quad \hat{T}(X, Y) = T(X, Y) + \pi(Y)X - \pi(X)Y = \pi(Y)X - \pi(X)Y$$

for $X, Y \in \mathfrak{X}(M)$, where $\hat{T}$ and T denote the torsion tensors of connections $\hat{\nabla}$ and ∇ , respectively. Then from (8) and (9) we have [4]

THEOREM 1. *The induced connection on a submanifold of a Riemannian manifold with a semisymmetric metric connection is also a semisymmetric metric connection.*

The Weingarten equation (with respect to ∇') is given by

$$(10) \quad \nabla'_X \xi_i = -A_i(X) + D_X \xi_i$$

for $X \in \mathfrak{X}(M)$, where A_i are tensors of type (1,1) on M and D is a (metric) connection in the normal bundle $T(M)^\perp$ with respect to the fibre metric induced from g [5, pp. 10–21].

Note that from (3) and (10) we have $h_i(X, Y) = g(Y, A_i(X))$, and thus we get

$$(11) \quad h_i(X, Y) = g(X, A_i(Y)) = g(A_i(X), Y)$$

for $X, Y \in \mathfrak{X}(M)$ since h_i are symmetric. While from (1) and (2) we have $\hat{\nabla}'_X \xi_i = \nabla'_X \xi_i + \lambda_i X$, which together with (10) implies $\hat{\nabla}'_X \xi_i = -(A_i - \lambda_i I) \cdot (X) + D_X \xi_i$, where I is the identity tensor. Defining tensors $\hat{A}_i$ of type (1,1) on M by $\hat{A}_i = A_i - \lambda_i I$, we get a more concise expression,

$$(12) \quad \hat{\nabla}'_X \xi_i = -\hat{A}_i(X) + D_X \xi_i$$

for $X \in \mathfrak{X}(M)$, which may be called the Weingarten equation with respect to $\hat{\nabla}'$.

We obtain, using (7) and (11), the following result which will be used later [1], [5, pp. 10–21]:

LEMMA. *The induced linear transformations, also denoted by A_i and $\hat{A}_i$, of the tangent space $T_m(M)$ at $m \in M$ defined by (10) and (12) satisfy, respectively, $h_i(X, Y) = g(A_i(X), Y)$ and $\hat{h}_i(X, Y) = g(\hat{A}_i(X), Y)$ for $X, Y \in T_m(M)$, and thus are symmetric with respect to g , i.e., $g(A_i(X), Y) = g(X, A_i(Y))$ and $g(\hat{A}_i(X), Y) = g(X, \hat{A}_i(Y))$ for $X, Y \in T_m(M)$.*

The mean curvature normal H of M (with respect to ∇) is given by $H = (1/n) \sum_{i=1}^p (\text{trace } A_i) \xi_i$ [5, pp. 29–42]. We define similarly the mean curvature normal $\hat{H}$ of M with respect to $\hat{\nabla}$ by $\hat{H} = (1/n) \sum_{i=1}^p (\text{trace } \hat{A}_i) \xi_i$. Let $X_j, j = 1, 2, \dots, n$, be n orthonormal local vector fields on M . Then H and $\hat{H}$ can be expressed as

$$(13) \quad H = \frac{1}{n} \sum_{i=1}^p \left\{ \sum_{j=1}^n h_i(X_j, X_j) \right\} \xi_i, \quad \hat{H} = \frac{1}{n} \sum_{i=1}^p \left\{ \sum_{j=1}^n \hat{h}_i(X_j, X_j) \right\} \xi_i.$$

If $h_i = k_i g$, where k_i are real-valued functions on M , then M is said to be totally umbilical (with respect to ∇). Similarly, if $\hat{h}_i = k_i g$, then M is said to be totally umbilical with respect to $\hat{\nabla}$ [4], [7, pp. 91–93].

We get from (2), (7) and (13) the following results [4]:

THEOREM 2. *The mean curvature normal of M and that of M with respect to the semisymmetric metric connection $\hat{\nabla}$ coincide if and only if the vector field P' is tangent to M .*

THEOREM 3. *A submanifold M of a Riemannian manifold M' is totally umbilical if and only if it is totally umbilical with respect to the semisymmetric metric connection $\hat{\nabla}$.*

3. Gauss curvature equation and Codazzi-Mainardi equation. We denote by $R'(X', Y')Z' = \nabla'_{X'} \nabla'_{Y'} Z' - \nabla'_{Y'} \nabla'_{X'} Z' - \nabla'_{[X', Y']} Z'$ for $X', Y', Z' \in \mathfrak{X}(M')$ and $R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z$ for $X, Y, Z \in \mathfrak{X}(M)$ the curvature tensors of ∇' and ∇ , respectively. Then the Gauss curvature equation (with respect to ∇' and ∇) is given by

$$(14) \quad R'(W, Z, X, Y) = R(W, Z, X, Y) + \sum_{i=1}^p \{h_i(X, Z)h_i(Y, W) - h_i(Y, Z)h_i(X, W)\}$$

for $X, Y, Z, W \in \mathfrak{X}(M)$, where

$$R'(W', Z', X', Y') = g(R'(X', Y')Z', W').$$

for $X', Y', Z', W' \in \mathfrak{X}(M')$ and

$$R(W, Z, X, Y) = g(R(X, Y)Z, W)$$

for $X, Y, Z, W \in \mathfrak{X}(M)$ are, respectively, the Riemann-Christoffel curvature tensors of M' and M (with respect to ∇' and ∇), and the Codazzi-Mainardi equation (with respect to ∇' and ∇) is given by

$$(15) \quad R'(\xi_i, Z, X, Y) = (\nabla_X h_i)(Y, Z) - (\nabla_Y h_i)(X, Z) + \sum_{j=1}^p g(\{h_j(Y, Z)D_X \xi_j - h_j(X, Z)D_Y \xi_j\}, \xi_i)$$

for $X, Y, Z \in \mathfrak{X}(M)$ [1], [5, pp. 22–29].

Next we shall find the Gauss curvature equation and the Codazzi-Mainardi equation with respect to the semisymmetric metric connections $\tilde{\nabla}'$ and $\tilde{\nabla}$. The curvature tensors of $\tilde{\nabla}'$ and $\tilde{\nabla}$ are defined, respectively, by

$$\tilde{R}'(X', Y')Z' = \tilde{\nabla}'_{X'} \tilde{\nabla}'_{Y'} Z' - \tilde{\nabla}'_{Y'} \tilde{\nabla}'_{X'} Z' - \tilde{\nabla}'_{[X', Y']} Z'$$

for $X', Y', Z' \in \mathfrak{X}(M')$ and $\tilde{R}(X, Y)Z = \tilde{\nabla}_X \tilde{\nabla}_Y Z - \tilde{\nabla}_Y \tilde{\nabla}_X Z - \tilde{\nabla}_{[X, Y]} Z$ for $X, Y, Z \in \mathfrak{X}(M)$. Then from (4), (9) and (12) we get

$$(16) \quad \begin{aligned} \tilde{R}'(X, Y)Z &= \tilde{R}(X, Y)Z + \sum_{i=1}^p \{\tilde{h}_i(X, Z)\tilde{A}_i(Y) - \tilde{h}_i(Y, Z)\tilde{A}_i(X)\} \\ &+ \sum_{i=1}^p \{(\tilde{\nabla}_X \tilde{h}_i)(Y, Z) - (\tilde{\nabla}_Y \tilde{h}_i)(X, Z) \\ &\quad + \tilde{h}_i(\pi(Y)X - \pi(X)Y, Z)\}\xi_i \\ &+ \sum_{i=1}^p \{\tilde{h}_i(Y, Z)D_X \xi_i - \tilde{h}_i(X, Z)D_Y \xi_i\} \end{aligned}$$

for $X, Y, Z \in \mathfrak{X}(M)$.

We define the Riemann-Christoffel curvature tensors of M' and M with respect to $\tilde{\nabla}'$ and $\tilde{\nabla}$, respectively, by $\tilde{R}'(W', Z', X', Y') = g(\tilde{R}'(X', Y')Z', W')$ and $\tilde{R}(W, Z, X, Y) = g(\tilde{R}(X, Y)Z, W)$ for $X', Y', Z', W' \in \mathfrak{X}(M')$ and $X, Y, Z, W \in \mathfrak{X}(M)$. Then from (16) and the Lemma, we obtain the Gauss curvature equation with respect to $\tilde{\nabla}'$ and $\tilde{\nabla}$ [4]:

$$(17) \quad \begin{aligned} \mathring{R}'(W, Z, X, Y) &= \mathring{R}(W, Z, X, Y) \\ &+ \sum_{i=1}^p \{ \mathring{h}_i(X, Z) \mathring{h}_i(Y, W) - \mathring{h}_i(Y, Z) \mathring{h}_i(X, W) \} \end{aligned}$$

for $X, Y, Z, W \in \mathfrak{X}(M)$. We also have from (16) the Codazzi-Mainardi equation with respect to $\mathring{\nabla}'$ and $\mathring{\nabla}$ [4]:

$$(18) \quad \begin{aligned} \mathring{R}'(\xi_i, Z, X, Y) &= (\mathring{\nabla}_X \mathring{h}_i)(Y, Z) - (\mathring{\nabla}_Y \mathring{h}_i)(X, Z) + \mathring{h}_i(\pi(Y)X - \pi(X)Y, Z) \\ &+ \sum_{j=1}^p g(\{ \mathring{h}_j(Y, Z) D_X \xi_j - \mathring{h}_j(X, Z) D_Y \xi_j \}, \xi_i) \end{aligned}$$

for $X, Y, Z \in \mathfrak{X}(M)$.

Now we suppose the Riemannian manifold M' is conformally flat and that the submanifold M is totally umbilical, then we can assume $\mathring{R}' = 0$ [6], and we also have $\mathring{h}_i = k_i g$, since M is totally umbilical with respect to $\mathring{\nabla}$ by Theorem 3. Then from (17) we get

$$(19) \quad \mathring{R}(W, Z, X, Y) = \{ g(Y, Z)g(X, W) - g(X, Z)g(Y, W) \} \sum_{i=1}^p (k_i)^2$$

for $X, Y, Z, W \in \mathfrak{X}(M)$, which implies that M is also conformally flat ($n > 3$) [3]. Thus we have [4]

THEOREM 4. *A totally umbilical submanifold in a conformally flat Riemannian manifold is conformally flat.*

4. Theorema egregium. We obtain a generalization of the theorema egregium of Gauss with respect to semisymmetric metric connection by the method of N. Hicks [1].

From the Gauss curvature equation (17) and the Lemma, we get

$$\begin{aligned} \mathring{R}'(X, Y, X, Y) &= \mathring{R}(X, Y, X, Y) \\ &+ \sum_{i=1}^p \{ g(\mathring{A}_i(X), Y)^2 - g(\mathring{A}_i(X), X)g(\mathring{A}_i(Y), Y) \} \end{aligned}$$

for $X, Y \in \mathfrak{X}(M)$. Therefore we have

THEOREM 5. *Let $\mathfrak{P}$ be a 2-dimensional subspace of $T_m(M)$, and let $\mathring{R}'(\mathfrak{P})$ and $\mathring{R}(\mathfrak{P})$ be, respectively, the sectional curvatures of $\mathfrak{P}$ in M' and M with respect to the semisymmetric connections $\mathring{\nabla}'$ and $\mathring{\nabla}$. If X and Y form an orthonormal base of $\mathfrak{P}$, then*

$$(20) \quad \mathring{R}'(\mathfrak{P}) = \mathring{R}(\mathfrak{P}) + \sum_{i=1}^p \{ g(\mathring{A}_i(X), Y)^2 - g(\mathring{A}_i(X), X)g(\mathring{A}_i(Y), Y) \}.$$

As immediate consequences of Theorem 5 we get [6]

COROLLARY 1. *If $\dim M' = 3$ and M is a surface in M' , then the determinant of $\mathring{A}_i$ (where there is now only one such map) is independent of $\mathring{A}_i$ but depends only on the Riemannian metric tensor g and the semisymmetric metric connections $\mathring{\nabla}'$ and $\mathring{\nabla}$.*

COROLLARY 2. If M' is a conformally flat Riemannian manifold of dimension 3 and M is a surface in M' , then there exists a semisymmetric metric connection $\tilde{\nabla}$ on M for which $\det \tilde{A}_i$ is an intrinsic invariant of M , and, when P' is tangent to M , $\det \tilde{A}_i (= \tilde{R}(\mathfrak{P}))$ is equal to $\det A_i$ which is the Gaussian curvature of M .

REFERENCES

1. N. J. Hicks, *Submanifolds of semi-Riemannian manifolds*, Rend. Circ. Mat. Palermo (2) **12** (1963), 137–149. MR **29** #1601.
2. —, *Notes on differential geometry*, Van Nostrand Math. Studies, no. 3, Van Nostrand, Princeton, N. J., 1965. MR **31** #3936.
3. T. Imai, *Notes on semi-symmetric metric connections*, Tensor **24** (1972), 293–296.
4. —, *Hypersurfaces of a Riemannian manifold with semi-symmetric metric connection*, Tensor **23** (1972), 300–306.
5. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Vol. II, Interscience Tracts in Pure and Appl. Math., no. 15, Vol. II, Interscience, New York, 1969. MR **38** #6501.
6. K. Yano, *On semi-symmetric metric connection*, Rev. Roumaine Math. Pures Appl. **15** (1970), 1579–1586. MR **43** #1078.
7. —, *Integral formulas in Riemannian geometry*, Pure and Appl. Math., no. 1, Dekker, New York, 1970. MR **44** #2174.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OKLAHOMA 73069