ARTIN ROOT NUMBERS FOR REAL CHARACTERS

JERRY GECHTER

ABSTRACT. For K and L number fields, χ a real-valued character on $\text{Gal}(K/L)$, the Artin root number $W(\chi)$ is ± 1 . We analyze the question of sign for χ a degree 2 character over $\mathbf{Q}$ induced from an abelian character on a quadratic extension.

Let $K/\mathbf{Q}$ be a finite Galois extension, and χ some character on $\text{Gal}(K/\mathbf{Q})$. The Artin L -function [2] $L(s, \chi)$ satisfies a functional equation of form $\xi(s, \chi) = W(\chi)\xi(1-s, \bar{\chi})$ where ξ is L multiplied by Γ factors and (constant)^s, and $W(\chi)$ is the Artin root number associated to χ . Iterating this functional equation gives $W(\chi)W(\bar{\chi}) = 1$. Further, if χ is real, then $W(\chi)^2 = 1$ or $W(\chi) = \pm 1$.

Until recently $W(\chi)$ was assumed always $+1$. The first examples to the contrary were given by Serre (unpublished) and Armitage [1] using a construction of Serre. The question of sign is of considerable interest: first because an L -function with $W(\chi) = -1$ gives a number field whose zeta function has a zero at $\frac{1}{2}$ (see [1]), and second because the question of sign is tied up with the existence of a normal basis for the ring of integers O_K over $\mathbf{Z}$ (see [4] and Fröhlich's Vancouver talk).

Here we consider the following set up: We start with k a quadratic extension of $\mathbf{Q}$ and ψ a primitive abelian character on k . Assume that K , the field defined by ψ via class field theory, is normal $/\mathbf{Q}$ and that the induced character $\chi = \psi^*$ on $\text{Gal}(K/\mathbf{Q})$ is real valued. Then we prove:

- (1) For k imaginary, $W(\chi) = +1$ always.
- (2) For k real, tamely ramified $/\mathbf{Q}$, $W(\chi) = \pm 1$, and both occur infinitely often with given k .

This result is subsumed in a recent paper of Fröhlich [5] (where $\mathbf{Q}$ is replaced by an arbitrary number field). The proof here is less elegant but more elementary.

The main idea of the proof is to look at ψ restricted to $\mathbf{Z} \subset O_k$. This is a Dirichlet character on $\mathbf{Z}$ and in fact we have

PROPOSITION 1. $\psi|_{\mathbf{Z}}$ is a not necessarily primitive character cotrained either with the identity or with the Kronecker symbol (d/n) , for $d = \text{discriminant of } k$.

PROOF. Since $\chi = \psi^*$ is real, $L(s, \psi)$ has real coefficients as a Dirichlet series. That in turn says the product of the Euler factors above a single rational prime has real coefficients. Immediately we get

Received by the editors February 4, 1975.

AMS (MOS) subject classifications (1970). Primary 10H10, 12A70; Secondary 12C25.

Key words and phrases. Artin root number, Gauss sum, Artin L -series, normal basis.

© American Mathematical Society 1976

$$\psi((p)) = \pm 1, 0 \quad \text{for } (d/p) = -1,$$

$$\psi(\underline{P}) = \pm 1, 0 \quad \text{for } (d/p) = 0, \underline{P}^2 = (p).$$

For $(d/p) = +1$,

$$(p) = \underline{P} \underline{P}^\sigma \quad \text{for } \sigma \in \text{Gal}(k/Q).$$

$$\begin{aligned} (1 - (\underline{P})p^{-s})^{-1} (1 - (\underline{P}^\sigma)p^{-s})^{-1} \\ = (1 - [(\underline{P}) + (\underline{P}^\sigma)]p^{-s} + ((p))p^{-2s})^{-1}, \end{aligned}$$

so $\psi(\underline{P}) + \psi(\underline{P}^\sigma) \in R$. Furthermore since K is normal $/Q$, $\psi(\underline{P})$ and $\psi(\underline{P}^\sigma)$ must be roots of unity of the same order. This implies

$$\psi(\underline{P}^\sigma) = \overline{\psi(\underline{P})}, \quad \text{and} \quad \psi((p)) = \psi(\underline{P})\psi(\underline{P}^\sigma) = 1.$$

We thus see that $\psi((p)) = +1$ or 0 unless $(d/p) = -1$. This means $\psi|Z$ factors through the Artin symbol

$$\left[\frac{k/Q}{p} \right],$$

which is the statement of the proposition.

Characters ψ such that $\psi|Z$ is cotrained with the identity are precisely the classical ring class characters.

PROPOSITION 2. *Let ψ be a primitive ring class character on a quadratic field. Then $W(\psi) = 1$.*

PROOF. In [3] we explicitly evaluate the local factors in Tate's formula for $W(\psi)$. This is somewhat tedious but straightforward. Fröhlich's approach is similar. Stark (unpublished) has suggested a purely analytic proof based on the form of the functional equation for ring class characters.

THEOREM 1. *If k is imaginary quadratic $W(\chi) = +1$.*

PROOF. Since the Artin L -function is invariant under inducing characters, $W(\chi) = W(\psi)$. Next, since k is totally imaginary, ψ is unramified at ∞ , so $\psi(-1) = +1$. By Proposition 1, $\psi|Z$ is cotrained with the identity or with (d/n) . For an imaginary quadratic field $(d/-1) = -1$, so ψ must be cotrained with the identity. Then by Proposition 2, $W(\psi) = +1$.

We now move on to the real quadratic case. Since Proposition 2 still applies, we need first to find characters ψ so that $\psi|Z$ is cotrained with (d/n) .

PROPOSITION 3. *If k is real, tamely ramified $/Q$, there exist infinitely many characters ψ on k satisfying*

- (a) ψ is primitive,
- (b) $\text{Ker } \psi$ is invariant under conjugation,
- (c) $\psi|Z$ is cotrained with (d/n) .

PROOF. We explicitly construct ψ as a product $\psi_p \psi_d$, where ψ_p and ψ_d are numerical characters on O_k (not necessarily trivial on the ray class of the fundamental unit) with the following properties: ψ_p is defined modulo some

rational prime p and is trivial on integers; $\psi_d|_{\mathbf{Z}}$ is (d/n) . In fact for k tamely ramified we can take ψ_d to be (d/n) as defined on $O_k \bmod(\sqrt{d})$.

Let u be the fundamental unit. $\psi_d(u) = \pm 1$. If it is $+1$, then ψ_d itself extends to a ray class character. If $\psi_d(u) = -1$, then we look for a rational prime p and a character ψ_p such that $\psi_p(u) = -1$. $\psi_p\psi_d$ then will extend to a ray class character ψ .

The existence of the character ψ_p is an exercise in class field theory (see [3]). That there are infinitely many ψ follows since the existence proof finds infinitely many ψ_p . If ψ is any character with the desired properties, $\psi\psi_p\psi_{p'}$ is another.

As yet we do not have $W(\psi) = -1$. For that we use a very special case of a theorem of Langlands (see [6], or [4] for this form). Let θ, ν be ray class characters with conductors (finite parts) f_θ, f_ν . Let $M(\theta)$ be the number of real primes at which θ is ramified. If $(f_\theta, f_\nu) = 1$ and if $M(\theta) + M(\nu) \equiv M(\theta\nu) \pmod{4}$, then $W(\theta)W(\nu) = W(\theta\nu)\nu(f_\theta)\theta(f_\nu)$.

We now use class field theory to construct a ring class character ψ' such that

$$(f_\psi, f_{\psi'}) = 1, \quad \psi(f_\psi) = -1, \quad \psi'(f_\psi) = 1.$$

Since ψ' is a ring class character it is either unramified or totally ramified at ∞ . Further, the same is true for ψ by (b) of Proposition 3. Hence the condition at the infinite primes in Langlands' theorem is satisfied. $(f_\psi, f_{\psi'}) = 1$ takes care of the other condition.

Collecting information we have $W(\psi)W(\psi') = W(\psi\psi')\psi(f_\psi)\psi'(f_\psi)$ by Langlands; $W(\psi') = 1$ by Proposition 2; $\psi(f_\psi) = -1$ and $\psi'(f_\psi) = 1$ by construction. Therefore

$$W(\psi) = -W(\psi\psi').$$

$\psi\psi'$ also satisfies (a), (b), and (c) of Proposition 3, so we have

THEOREM 2. *Let k be a real quadratic field, tamely ramified / $\mathbf{Q}$. Then there exist infinitely many ray class characters ψ on k such that*

- (a) ψ is primitive,
- (b) $\text{Ker } \psi$ is invariant under conjugation,
- (c) $\psi|_{\mathbf{Z}}$ is cotrained with (d/n) ,
- (d) $W(\psi) = -1$.

The same is true with (d) replaced by $W(\psi) = +1$.

Only condition (c) needs comment, for it shows (given (b)) that ψ^* is a real character. To see this we reverse the reasoning in Proposition 1.

$$\begin{aligned} \psi^* \text{ is real} &\Leftrightarrow \psi^*([(K/\mathbf{Q})/P]) \in R \text{ for all primes } P \text{ of } K, \\ &\Leftrightarrow L(s, \psi^*) \text{ has real coefficients as a Dirichlet series,} \\ &\Leftrightarrow L(s, \psi) \text{ has real coefficients as a Dirichlet series,} \\ &\Leftrightarrow \psi((p)) = \pm 1, 0 \quad \text{for } (d/p) = -1, \\ &\quad \psi((p)) = +1, 0 \quad \text{for } (d/p) = 1, 0, \end{aligned}$$

and this last condition follows from (c). Consequently Theorem 2 is precisely the result asserted in the case of real k .

REFERENCES

1. J. V. Armitage, *Zeta functions with a zero at $s = \frac{1}{2}$* , Invent. Math. **20**(1973), 125–138.
2. E. Artin, *Zur Theorie der L-Reihen mit allgemeinen Gruppencharakteren*, Hamb. Abh. **8**(1930), 292–306.
3. J. Gechter, *Gaussian sums for Artin L-series*, Thesis, M.I.T., Cambridge, Mass., 1974.
4. A. Fröhlich, *Artin root numbers and normal integral bases for quaternion fields*, Invent. Math. **17**(1972), 143–166. MR48 #2115.
5. ———, *Artin root numbers, conductors, and representations for generalized quaternion groups*, Proc. London Math. Soc. **28**(1974), 402–438.
6. A. Weil, *Dirichlet series and automorphic forms*, Lecture Notes in Math., vol. 189, Springer-Verlag, New York, 1971.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON 98195