JOIN-PRINCIPAL ELEMENT LATTICES

E. W. JOHNSON

ABSTRACT. Let $(\mathcal{L}, M)$ be a local Noether lattice. If the maximal element M is meet principal, it is well known and easily seen that every element of $\mathcal{L}$ is meet principal. In this note, we obtain the corresponding result for M join-principal. We also consider join-principal elements generally under the assumption of the weak union condition and show, for example, that the square of a join-principal element is principal.

Throughout we assume that $\mathcal{L}$ is a local Noether lattice with maximal element M .

THEOREM 1. *If $A \in \mathcal{L}$ is join-principal, then, for each $n \geq 1$, A^n has a unique minimal basis in $\mathcal{L}/((0:A) \wedge A)$.*

PROOF. Use Theorem 1 of [2] to choose a minimal base $E_1, \dots, E_k$ for A such that $E_i E_j = 0$ whenever $i \neq j$.

Let F be any principal element $\leq A^n \vee ((0:A) \wedge A)$. Then by Lemma 1 of [3], there exist principal elements $F_1, \dots, F_{k+1}$ such that

$$F \vee \left(\bigvee_{i \neq j} F_i E_i^n \right) = \bigvee_i F_i E_i^n$$

for $j = 1, \dots, k+1$, where $E_{k+1} = I$ and $F_{k+1} \leq (0:A) \wedge A$.

Then $FE_j = F_j E_j^{n+1}$ ($1 \leq j \leq k$), so

$$FA = \bigvee_{1 \leq j \leq k} F_j E_j^{n+1} = \bigvee_{1 \leq j \leq k} F_j E_j^n A.$$

Hence

$$F \vee (0:A) = \left(\bigvee_{1 \leq j \leq k} F_j E_j^n \right) \vee (0:A).$$

So, since principal elements are join-irreducible,

$$F \vee ((0:A) \wedge A) = F_j E_j^n \vee ((0:A) \wedge A),$$

for some j . Hence either $F \vee ((0:A) \wedge A) = E_j^n \vee ((0:A) \wedge A)$, or $F \leq MA^n \vee ((0:A) \wedge A)$ and F cannot be used in a minimal base for A^n in

Received by the editors February 25, 1974.

AMS (MOS) subject classifications (1970). Primary 06A20.

© American Mathematical Society 1976

$\mathcal{L}/(0:A) \wedge A$. It follows that the only minimal bases for A^n in $\mathcal{L}/((0:A) \wedge A)$ are subsets of $E_1^n \vee ((0:A) \wedge A), \dots, E_k^n \vee ((0:A) \wedge A)$. On the other hand, if $E_j^n \leq E_1^n \vee \dots \vee E_j^n \vee \dots \vee E_k^n \vee ((0:A) \wedge A)$, then $E_j^n A = E_j^{n+1} \leq (E_1^n \vee \dots \vee E_j^n \vee \dots \vee E_k^n \vee ((0:A) \wedge A))E_j = 0$. Hence A^n has the unique minimal base in $\mathcal{L}/A \wedge (0:A)$ consisting of those $E_j^n \vee (A \wedge (0:A))$ with $E_j^n A \neq 0$. Q.E.D.

COROLLARY 2. *If $\mathcal{L}$ satisfies the weak union condition and if $A \in \mathcal{L}$ is join-principal, then $A = E \vee ((0:A) \wedge A)$ for some principal element E . In particular, $A^2 = E^2$ is principal.*

PROOF. If the weak union condition is satisfied, only principal elements have unique minimal bases. Since $\mathcal{L}/A \wedge (0:A)$ inherits the hypothesis from $\mathcal{L}$, the result follows.

COROLLARY 3. *Let (R, M) be a local ring in which M is join-principal. Then, as an R -module, M is the direct sum of a cyclic R -module and an R/M -vector space.*

In the general setting, we get the following

THEOREM 4. *If the maximal element M of $\mathcal{L}$ is join-principal, then $\mathcal{L}/(0:M)$ is distributive.*

PROOF. Let F be any principal element of $\mathcal{L}$, $F \not\leq 0:M$. By Theorem 1 and Corollary 1 of [2], there exist principal elements $E_1, \dots, E_k$ in $\mathcal{L}$ such that $E_1 \vee (0:M), \dots, E_k \vee (0:M)$ are an independent minimal base of M in $\mathcal{L}/(0:M)$. Choose n so that $F \leq M^n \vee (0:M)$ and $F \not\leq M^{n+1} \vee (0:M)$. Since M is join-principal in $\mathcal{L}$, it follows that $F \vee (0:M) = E_i^n \vee (0:M)$ for some i . Since the $E_j \vee (0:M)$ are independent, it follows that $\mathcal{L}/(0:M)$ is distributive. Q.E.D.

COROLLARY 5. *If the maximal element M of $\mathcal{L}$ is join-principal, then every element of $\mathcal{L}$ is join-principal.*

PROOF. Let B and A be any elements of $\mathcal{L}$. Since M is join-principal in $\mathcal{L}/C$ for every C , it suffices to show that $BA:A = B \vee (0:A)$.

Let $E_1, \dots, E_k$ be independent principal elements such that $E_1 \vee (0:M), \dots, E_k \vee (0:M)$ form the minimal base for M in $\mathcal{L}/0:M$. Then from the proof of Theorem 4, $AB:A$ and $B \vee (0:A)$ are both joins of powers of the $E_i \vee (0:M)$ in $\mathcal{L}/0:M$.

Set $B \vee (0:A) = E_1^{s_1} \vee \dots \vee E_k^{s_k} \vee (0:M)$, and assume $E_i^n \leq BA:A$. Then $E_i^n A \leq A(B \vee (0:A)) = AE_1^{s_1} \vee \dots \vee AE_k^{s_k}$. Since $E_j X$ is a power of E_j for every $X \neq 1$ and since the E_j are independent, it follows that $E_i^n A \leq E_i^{s_i} A$. If $n < s_i$, then $A \leq E_i^{s_i-n} A \vee (0:E_i^n)$. However, in this case, $A \leq 0:E_i^n$ and $E_i^n \leq 0:A$. If $n \geq s_i$, then $E_i^n \leq E_1^{s_1} \vee \dots \vee E_k^{s_k}$. In either case, $E_i^n \leq B \vee (0:A)$. Hence $BA:A \leq B \vee (0:A)$. Q.E.D.

REFERENCES

1. E. W. Johnson and J. P. Lediaev, *Structure of Noether lattices into join-principal maximal*

elements, Pacific J. Math. **37** (1971), 101–108. MR **46** #7108.

2. ———, *Join-principal elements in Noether lattices*, Proc. Amer. Math. Soc. **36** (1972), 73–78. MR **46** #5301.

3. E. W. Johnson and Michael Detlefsen, *Prime sequences and distributivity in local Noether lattices*, Fund. Math. **86** (1974), 149–156.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA 52242