SOME l -SIMPLE PATHOLOGICAL LATTICE-ORDERED GROUPS

A. M. W. GLASS¹ AND STEPHEN H. MCCLEARY

ABSTRACT. The purpose of this note is to explore two recently discovered lattice-ordered groups, and to establish that they are l -simple, i.e., have no proper l -ideals. These groups are "pathologically o -2-transitive" ordered permutation groups acting on the real line. The two groups exhibit even nastier properties than previous pathological groups, and serve as counterexamples to several conjectures about l -groups.

1. Introduction. $\mathbf{R}$ will denote the set of real numbers ordered in the usual way, and $A(\mathbf{R})$ the lattice-ordered group of all order-preserving permutations of $\mathbf{R}$, with $f \leq g$ if and only if $xf \leq xg$ for all $x \in \mathbf{R}$. Let F be an l -subgroup of $A(\mathbf{R})$, i.e., a subgroup which is simultaneously a sublattice. F is said to be o -2-transitive if for all $u < v$ and $x < y$ in $\mathbf{R}$, there exists $f \in F$ such that $uf = x$ and $vf = y$. For $f \in F$, let $\text{support}(f) = \{x \in \mathbf{R}: xf \neq x\}$. An o -2-transitive l -subgroup F of $A(\mathbf{R})$ having no element of bounded support other than the identity element 1 is called *pathologically o -2-transitive*.

The grandfather of all pathologically o -2-transitive groups, which was exhibited by Charles Holland in [4], was $\{g \in A(\mathbf{R}): \exists n = n(g) \in \mathbf{Z}^+ \text{ such that } \forall x \in \mathbf{R}, \forall k \in \mathbf{Z}, (x + kn)g = xg + kn\}$. Here $\mathbf{Z}$ denotes the integers and $\mathbf{Z}^+$ the positive integers. This group will be denoted throughout by G . G was shown in [5] to be l -simple, i.e., to have no proper l -ideals (normal convex l -subgroups). In [7], another pathological group was introduced, namely $\{g \in A(\mathbf{R}): \forall x \in \mathbf{R}, \exists n = n(g, x) \in \mathbf{Z}^+ \text{ such that } \forall k \in \mathbf{Z}, (x + kn)g = xg + kn\}$, here denoted by H , and H was shown to be l -simple. In [7] and [6] there were established certain properties of this class of groups which led to the name "pathological". It was found [2] that these groups are not always l -simple. Pathological groups played a crucial role in [3] in the theory of α^* -extensions and $\dagger$ -extensions, where the two groups $\tilde{G}$ and $\tilde{H}$ (defined below) arose quite naturally. $(\tilde{H}, \mathbf{R})$ turned out to be the unique $\dagger$ -closure of the groups $(G, \mathbf{R})$, $(H, \mathbf{R})$, and $(\tilde{G}, \mathbf{R})$.

2. The l -groups $\tilde{G}$ and $\tilde{H}$. For $f \in A(\mathbf{R})$ and $x \in \mathbf{R}$, let $\Delta(f, x) = xf - x$. Observe that

$$|\Delta(f, x + kn) - \Delta(f, x)| = |(x + kn)f - (xf + kn)|;$$

this fact will be used heavily.

Let

Received by the editors August 21, 1975.

AMS (MOS) subject classifications (1970). Primary 06A55.

Key words and phrases. Lattice-ordered group, l -simple, ordered permutation group.

¹Research supported in part by a grant from the Bowling Green State University Faculty Research Committee.

© American Mathematical Society 1976

$$\tilde{G} = \{g \in A(\mathbf{R}): \forall \varepsilon > 0, \exists n = n(g, \varepsilon) \in \mathbf{Z}^+ \text{ s.t. } \forall x \in \mathbf{R}, \forall k \in \mathbf{Z}, \\ |\Delta(g, x + kn) - \Delta(g, x)| < \varepsilon \text{ and } |\Delta(g^{-1}, x + kn) - \Delta(g^{-1}, x)| < \varepsilon\}.$$

Let

$$\tilde{H} = \{g \in A(\mathbf{R}): \forall \varepsilon > 0, \forall x \in \mathbf{R}, \exists n = n(g, \varepsilon, x) \in \mathbf{Z}^+ \text{ s.t. } \forall k \in \mathbf{Z}, \\ |\Delta(g, x + kn) - \Delta(g, x)| < \varepsilon \text{ and } |\Delta(g^{-1}, x + kn) - \Delta(g^{-1}, x)| < \varepsilon\}.$$

These groups were mentioned briefly in [3] in connection with $\dagger$ -extensions, but without proof even of the fact they are groups. Here we fill in some details, and establish their l -simplicity. Clearly $G \subseteq \tilde{G} \subseteq \tilde{H}$ and $G \subseteq H \subseteq \tilde{H}$.

THEOREM. $\tilde{G}$ and $\tilde{H}$ are l -simple lattice-ordered groups.

PROOF. We begin by showing that $\tilde{H}$ is closed under product. Let $g, h \in \tilde{H}$, and let $x \in \mathbf{R}$ and $\varepsilon > 0$. Pick $t_1 < xg < t_2$ such that $t_2h - t_1h < \varepsilon/2$. Let $\varepsilon_1 = t_2h - t_1h < \varepsilon/2$ and $\varepsilon_2 = \min\{t_2 - xg, xg - t_1, \varepsilon/2\}$. Let

$$n = n(h, \varepsilon_1, t_1)n(h, \varepsilon_1, t_2)n(g, \varepsilon_2, x),$$

and let $k \in \mathbf{Z}$. Now

$$|(x + kn)g - (xg + kn)| = |\Delta(g, x + kn) - \Delta(g, x)| < \varepsilon_2,$$

so that $t_1 + kn < (x + kn)g < t_2 + kn$, and thus

$$(i) \quad (t_1 + kn)h < (x + kn)gh < (t_2 + kn)h.$$

Also,

$$(ii) \quad t_1h + kn < xgh + kn < t_2h + kn,$$

and by the definition of ε_1 ,

$$(iii) \quad (t_1 + kn)h < t_2h + kn,$$

and

$$(iv) \quad t_1h + kn < (t_2 + kn)h.$$

Since $(t_2h + kn) - (t_1h + kn) = t_2h - t_1h < \varepsilon/2$, and since $|(t_i + kn)h - (t_ih + kn)| < \varepsilon/2$ ($i = 1, 2$), we have

$$|(t_2 + kn)h - (t_1h + kn)| < \varepsilon \quad \text{and} \quad |(t_2h + kn) - (t_1 + kn)h| < \varepsilon.$$

In view of the numbered facts, this yields

$$|\Delta(gh, x + kn) - \Delta(gh, x)| = |(x + kn)gh - (xgh + kn)| < \varepsilon.$$

Since $(gh)^{-1} = h^{-1}g^{-1}$, there exists $m \in \mathbf{Z}^+$ such that for all $k \in \mathbf{Z}$, $|\Delta((gh)^{-1}, x + km) - \Delta((gh)^{-1}, x)| < \varepsilon$, and then nm works for both gh and $(gh)^{-1}$. Therefore $gh \in \tilde{H}$.

In the above argument, if $g, h \in \tilde{G}$, then n and m can be chosen independently of x , so that $gh \in \tilde{G}$, provided h is uniformly continuous, which we now show. Let $\varepsilon > 0$ and $n = n(h, \varepsilon/3)$. There exists $\delta > 0$ such that if $x, y \in [0, 2n]$ and $|x - y| < \delta$, then $|xh - yh| < \varepsilon/3$. But $|\Delta(h, x + kn) - \Delta(h, x)| < \varepsilon/3$ for all $k \in \mathbf{Z}$. A simple computation shows that if $u, v \in \mathbf{R}$ and $|u - v| < \delta$, then $|uh - vh| < \varepsilon$. Hence h is uniformly continuous.

We have shown that $\tilde{H}$ and $\tilde{G}$ are subgroups of $A(\mathbf{R})$, and clearly they are *l*-subgroups. Since $G \subseteq \tilde{G} \subseteq \tilde{H}$ and G is *o*-2-transitive, so are $\tilde{H}$ and $\tilde{G}$. Moreover, $\tilde{H}$ (and thus also $\tilde{G}$) are pathological. For suppose $1 \neq h \in \tilde{H}$, so that there exists $x \in \mathbf{R}$ such that $xh \neq x$. Let $\epsilon = |\Delta(h, x)|/2 > 0$. Then there exists $n = n(h, \epsilon, x) \in \mathbf{Z}^+$ such that $|\Delta(h, x + kn) - \Delta(h, x)| < \epsilon$ for all $k \in \mathbf{Z}$. Thus $(x + kn)h \neq x + kn$ for all $k \in \mathbf{Z}$.

Recall that G is *l*-simple [4]. Thus to prove that $\tilde{H}$ is *l*-simple, it is enough to show that if $M \neq \{1\}$ is an *l*-ideal of $\tilde{H}$, then some element of M exceeds a (strictly) positive element of G , since then M will contain all translations. (Let $1 < f \in \tilde{H}$ and $n = n(f, 1, 0)$. Then $xf - x < 0f + n + 1$ for all $x \in \mathbf{R}$, so f is bounded by a translation.) Let $1 < h \in M$. There exists $x \in \mathbf{R}$ such that $x < xh$. Let $\epsilon = (xh - x)/2 > 0$ and $n = n(h, \epsilon, x)$. Then $(x + kn)h > xh + kn - \epsilon$ for all $k \in \mathbf{Z}$. Hence $(x + kn)h > x + kn + \epsilon$. Thus for all $s \in [x + kn, x + kn + \epsilon/2]$, we have

$$sh - s > (x + kn)h - (x + kn + \epsilon/2) > \epsilon/2.$$

Then h exceeds a positive $g \in G$ which moves points near each $x + kn$ and agrees with the identity elsewhere. As indicated above, it follows that $\tilde{H}$ is *l*-simple.

This argument also shows that $\tilde{G}$ is *l*-simple. Alternatively, the *l*-simplicity of $\tilde{G}$ can be deduced from a routine modification of the argument in [5] for that of G .

G is actually simple as a group [8], and so is H . It is not known whether $\tilde{G}$ and $\tilde{H}$ are simple as groups.

PROPOSITION. *($\tilde{H}, \mathbf{R}$) and ($\tilde{G}, \mathbf{R}$) are not isomorphic as ordered permutation groups.*

PROOF. It suffices to show that $\tilde{H} \neq \tilde{G}$. For it was shown in [3] (without determining whether $\tilde{H} = \tilde{G}$) that $(\tilde{H}, \mathbf{R})$ is the unique $+$ -closure of $(\tilde{G}, \mathbf{R})$, so that $(\tilde{H}, \mathbf{R})$ is $+$ -closed, while if $\tilde{H} \neq \tilde{G}$, $(\tilde{G}, \mathbf{R})$ is not $+$ -closed. (See [3] for definitions.)

To construct $h \in \tilde{H} \setminus \tilde{G}$, we proceed as follows: Let $S_1 = \{k \in \mathbf{Z} : k \equiv 0 \pmod{2}\}$. For $i > 1$, let s_i be that integer not in $S_1 \cup \dots \cup S_{i-1}$ having smallest absolute value (taking the positive choice in case of ties), and let S_i be $\{k \in \mathbf{Z} : k \equiv s_i \pmod{2^i}\}$. The S_i 's partition $\mathbf{Z}$. Let g_i be the order-preserving permutation of $[0, 1]$ defined by $xg_i = x$ for $x \in [0, 1/(i+2)]$ and $xg_i = (x+1)/2$ for $x \in [1/(i+1), 1]$, with the remainder filled in linearly. For $y_i = ((1/i+1) + (1/i+2))/2$, $|y_i g_i - y_j g_j| > 1/8$ for all $i \neq j \in \mathbf{Z}^+$. Now define h on $[k, k+1]$, $k \in \mathbf{Z}$, by the rule $xh = (x-k)g_i + k$, where $k \in S_i$.

Then $h \in H \subseteq \tilde{H}$ since when $x \in [k, k+1]$ and $k \in S_i$, $n(h, x) = 2^i$. Moreover, $h \notin \tilde{G}$, for when $k \in S_i$, 2^i is the smallest value for $n(h, 1/8, x = k + y_i)$. Hence $\tilde{H} \neq \tilde{G}$, proving the proposition.

Presumably $\tilde{H}$ and $\tilde{G}$ are not isomorphic as *l*-groups, but we have not been able to prove this. G is not isomorphic even as a group to $\tilde{G}$, $\tilde{H}$ or H since there exists $1 \neq z \in G$ such that each $g \in G$ commutes with some positive power of z , whereas it can be shown that this is false for the other three groups. Also, H is not isomorphic as an *l*-group to either $\tilde{G}$ or $\tilde{H}$. This can be

seen in terms of order units. If F is an l -group, $u \in F^+$ is called a *strong order unit* if every $f \in F$ is exceeded by some power of u , and we shall call u a *mild order unit* if every $f \in F$ fails to exceed some power of u . In all four of our pathological groups, positive translations are strong order units. In H , every mild order unit is a strong order unit, the two conditions being satisfied by those u 's in H^+ which have no fixed points. On the other hand, in the next section we shall construct $g \in \tilde{G}^+ \subseteq \tilde{H}^+$ fixing precisely one point. Then g is a mild order unit, but not a strong order unit.

3. Pathological groups as counterexamples. In [4, Theorem 7 and p. 432], Charles Holland proved a theorem and stated a conjecture about transitive ordered permutation groups acting on arbitrary totally ordered sets S . Here we consider the special case in which the group acts on $\mathbf{R}$.

A *prime* subgroup of an l -group F is a convex l -subgroup P such that $f \wedge g = 1$ implies $f \in P$ or $g \in P$. A prime subgroup containing no l -ideal of F other than $\{1\}$ is called a *representing* subgroup. For any transitive lattice-ordered permutation group (F, S) , the point stabilizers $F_s = \{f \in F: sf = s\}$ are representing subgroups of F . The condition that these stabilizers be *maximal* representing subgroups of F is of interest because it is equivalent to a certain kind of "minimality" of S .

THEOREM (HOLLAND). *Let $(F, \mathbf{R})$ be a transitive lattice-ordered permutation group having a nonidentity element of bounded support, and let θ be an l -isomorphism of F onto some transitive l -subgroup of some $A(S)$ such that the point stabilizers of $(F\theta, S)$ are maximal representing subgroups of $F\theta$. Then there exists an o -isomorphism ψ from $\mathbf{R}$ onto S which induces θ in the sense that for all $f \in F$, $f\theta = \psi^{-1}f\psi$.*

Holland says that "it seems likely that the theorem remains true" without the hypothesis that F contains a nonidentity element of bounded support. However, such is not the case. The conclusion of the theorem requires that every maximal representing subgroup of F be a point stabilizer F_r for some $r \in \mathbf{R}$ (see [4]). We now construct a maximal representing subgroup K of the original pathological group G which is not a stabilizer of $(G, \mathbf{R})$. (The construction works equally well for the groups H , $\tilde{G}$, and $\tilde{H}$.)

Let $T_i = b_i + 4^i\mathbf{Z}$, $i = 1, 2, \dots$, where $b_1 = 1$ and for $i > 1$, b_i is the second smallest positive integer in T_{i-1} . Then $T_1 \supset T_2 \supset \dots$ forms a descending tower having empty intersection. Enlarge this tower to an ultrafilter $\mathfrak{F}$ on $\mathbf{R}$. Let $P = \{g \in G: \exists F_g \in \mathfrak{F} \text{ such that } rg = r\forall r \in F_g\}$, a prime subgroup of G . Given any $r \in \mathbf{R}$, we can pick a T_i that does not contain r and construct an order-preserving permutation g which moves r , has period 4^i (so that $g \in G$), and fixes each $t \in T_i$ (so that $g \in P$). Hence P is not contained in any stabilizer G_r . $P \neq G$ since each $g \in P$ fixes some $r \in \mathbf{R}$. Since G has a strong order unit, Zorn's lemma guarantees that P is contained in a maximal proper prime subgroup K of G . Since K is proper and G is l -simple, K is a (maximal) representing subgroup of G . K is not equal to any G_r , for then $P \subseteq G_r$. Thus the key hypothesis of Holland's theorem is in fact needed. We remark without proof that the right regular representation of G on the totally ordered set G/K of right cosets of K is also pathologically o -2-transitive.

Now we show that $\tilde{G}$ and $\tilde{H}$ enjoy a very unpleasant property making them even more perverse than the earlier pathological groups: They contain an element g which fixes precisely one point. This answers a question of existence posed by McCleary and resolves in the negative a conjecture of Richard Ball (see [1]).

First we construct $\bar{g} \in A(\mathbf{R})$ by setting

$$x\bar{g} = \begin{cases} x & \text{if } x \text{ is an integer divisible by 4,} \\ x + \frac{1}{2} & \text{if } x \in [2p - \frac{1}{4}, 2p + \frac{1}{4}], \quad p \text{ an odd integer,} \end{cases}$$

and then filling in the remainder linearly. Now, for $n \geq 2$ and q odd, we set $xg = x + 1/2^n$ throughout the interval containing 2^nq and extending in each direction until the graph of $x + 1/2^n$ first crosses the graph of $\bar{g}$, and we set $xg = x\bar{g}$ for all other x . It is easily seen that for $2^m < \varepsilon$, we may take $n(g, \varepsilon) = 2^m$, so that $g \in \tilde{G} \subseteq \tilde{H}$. By construction, g fixes only 0.

Let K be an l -group. A convex l -subgroup C is said to be *closed* if whenever a subset of C has a supremum in K , that supremum lies in C . If $k \in K$, $K(k)^*$ denotes the closed convex l -subgroup generated by k . Now let (K, S) be a lattice-ordered permutation group, and let $\bar{S}$ be the completion of S by Dedekind cuts (without endpoints). If $T \subseteq \bar{S}$, let $K_T = \{k \in K: tk = t \forall t \in T\}$. If $D \subseteq G$, let $\bar{F}x D = \{s \in \bar{S}: sd = s \forall d \in D\}$. For any $k \in K$, we clearly have

$$\bar{F}x(k) \cap \{s \in \bar{S}: G_s \text{ is closed}\} \subseteq \bar{F}x(K(k)^*) \subseteq \bar{F}x(k).$$

In [1], Ball introduced the concept of a *full* convex l -subgroup of an l -group K , meaning one which is a union of closed convex l -subgroups. When (K, S) is a permutation group, F denotes $\{s \in \bar{S}: G_s \text{ is full}\}$. Clearly we can improve the above result to say $\bar{F}x(k) \cap F \subseteq \bar{F}x(K(k)^*) \subseteq \bar{F}x(k)$. Ball conjectured [1, 5.1.10] that for transitive groups, $\bar{F}x(k) \cap F = \bar{F}x(K(k)^*)$ for all $k \in K^+$. In view of the fact that $K_{\bar{F}x C} = C$ when C is closed in K , this would imply that $K(k)^* = K_{\bar{F}x(k) \cap F}$ for all $k \in K^+$ [1, 5.1.9].

However, it has been observed in a discussion between Ball and the present authors that $\tilde{G}$ and $\tilde{H}$ provide counterexamples to these conjectures. As Ball observed in [1], $(G, \mathbf{R})$ has full stabilizers, for $G_0 = \bigcup \{G_{nz}: n \in \mathbf{Z}^+\}$, a union of closed subgroups. The same is true of $(H, \mathbf{R})$. But for $\tilde{G}$ and $\tilde{H}$, the stabilizers are not full. Let $g \in \tilde{G}$ fix only 0, as above. Let z denote the translation $x \mapsto x + 1$, and let $r \in \mathbf{R} \setminus [-1, 0]$. Since g fixes only 0, there exists a power g^n of g such that $rg^n > r + 1$. Then $z = \sup\{g^n \wedge z\}$ since if $g^n \wedge z \leq f < z$ for all r , zf^{-1} would have bounded support. Thus, letting K be either $\tilde{G}$ or $\tilde{H}$, we have $z \in K(g)^*$, so that $K(g)^* = K$. Since $g \in K_0$, K_0 cannot be full. Hence F is empty, so that Ball's weaker conjecture would here say that for any $k \in K^+$, $K(k)^* = K$. This fails for any k lying in a proper closed convex l -subgroup of K , e.g., in K_z .

4. Other pathological groups. Our four examples of pathological groups all have strong order units. The examples below will lack strong order units, and thus will not be isomorphic as l -groups to any of the above examples. First, the restriction $(G_0|\mathbf{R}^+, \mathbf{R}^+)$ of the stabilizer G_0 is pathological. Those of its elements which are bounded below form a proper l -ideal, so this example is

not l -simple. A smaller proper l -ideal is given by $M = \{g \in G_0 | \mathbf{R}^+ : \exists n = n(g) \in \mathbf{Z}^+ \text{ such that } \forall x \in \mathbf{R}^+, (x + kn)g = xg + kn \text{ for all } k \in \mathbf{Z}^+ \text{ and there exists a neighborhood of } 0 \text{ throughout which } xg = x\}$. M is l -simple. For let $1 < g$, $h \in M^+$, and let $n = n(g)n(h)$. A standard compactness argument shows that g is exceeded by a product of conjugates (by elements of period n) of h .

5. Epilogue. For convenience, we tabulate the properties of the six pathological groups mentioned above. No two of these are isomorphic as l -groups except, conceivably, $\tilde{G}$ and $\tilde{H}$, and even these are not isomorphic as ordered permutation groups.

	l -simple	strong order unit	element fixing precisely one point	full stabilizers
G	Yes	Yes	No	Yes
H	Yes	Yes	No	Yes
$\tilde{G}$	Yes	Yes	Yes	No
$\tilde{H}$	Yes	Yes	Yes	No
$G_0 \mathbf{R}^+$	No	No	No	Yes
M	Yes	No	No	Yes

REFERENCES

1. Richard Ball, *Full convex l -subgroups of a lattice ordered group*, Ph. D. Dissertation, University of Wisconsin, Madison, Wis., 1974.
2. A. M. W. Glass, *l -simple lattice-ordered groups*, Proc. Edinburgh Math. Soc. **19** (1974), 133–138.
3. A. M. W. Glass, W. Charles Holland and Stephen H. McCleary, *a^* -closures of completely distributive lattice-ordered groups*, Pacific J. Math. **59** (1975), 43–67.
4. W. Charles Holland, *Transitive lattice-ordered permutation groups*, Math. Z. **87** (1965), 420–433. MR **31** #2310.
5. Justin T. Lloyd, *Complete distributivity in certain infinite permutation groups*, Michigan Math. J. **14** (1967), 393–400. MR **36** #2544.
6. Stephen H. McCleary, *Closed subgroups of lattice-ordered permutation groups*, Trans. Amer. Math. Soc. **173** (1972), 303–314. MR **47** #97.
7. ———, *o -2-transitive ordered permutation groups*, Pacific J. Math. **49** (1973), 425–429. MR **50** #2018.
8. ———, *Some simple homeomorphism groups having nonsolvable outer automorphism groups* (to appear).

DEPARTMENT OF MATHEMATICS, BOWLING GREEN STATE UNIVERSITY BOWLING GREEN, OHIO 43403

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GEORGIA 30602