SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

A NEW PROOF OF GIESY'S THEOREM ON NON- B -CONVEXITY

K. P. R. SASTRY AND S. VENKATA RATNAM NAIDU

ABSTRACT. A new proof of Giesy's theorem on non- B -convexity is given.

The following theorem was proved by D. P. Giesy in [2, Lemma 6, §1] and has been often used since (see, for example [1]):

A necessary and sufficient condition for non- D -convexity of a normed linear space X is that given $k \geq 2$, $\varepsilon > 0$, there exist $x_1, x_2, \dots, x_k \in S$ such that for each choice of scalars $\alpha_1, \alpha_2, \dots, \alpha_k$,

$$(1 - \varepsilon) \sum_{i=1}^k |\alpha_i| \leq \left\| \sum_{i=1}^k \alpha_i x_i \right\|.$$

It is known that D -convexity is equivalent to B -convexity [2]. In this note a simple proof for the necessity part is given. The sufficiency is trivial, of course. The notation is the same as that of Giesy [2].

PROOF OF THE NECESSITY. Let $k \geq 2$ and $0 < \varepsilon < 1$. Let $\delta = k^{-1}\varepsilon$.

X is not D -convex $\Rightarrow X$ is not D, x, δ -convex

$\Rightarrow \exists x_1, x_2, \dots, x_k \in S \ni$ for any $\lambda_1, \lambda_2, \dots, \lambda_k \in D$,

$$k^{-1} \left\| \sum_{i=1}^k \lambda_i x_i \right\| > 1 - \delta.$$

Let $\alpha_1, \alpha_2, \dots, \alpha_k$ be scalars such that $\sum_{i=1}^k |\alpha_i| = 1$. Consider

$$\begin{aligned} \left\| \sum_{i=1}^k \alpha_i x_i \right\| &\geq \left\| \sum_{i=1}^k sg(\alpha_i) x_i \right\| - \left\| \sum_{i=1}^k (sg(\alpha_i) - \alpha_i) x_i \right\| \\ &> k(1 - \delta) - \sum_{i=1}^k |sg(\alpha_i) - \alpha_i| = k - \varepsilon - \sum_{i=1}^k (1 - |\alpha_i|) \\ &= k - \varepsilon - (k - 1) = 1 - \varepsilon. \end{aligned}$$

Received by the editors November 24, 1975.

AMS (MOS) subject classifications (1970). Primary 46B10.

Key words and phrases. B -convexity, D -convexity.

© American Mathematical Society 1976

Hence $(1 - \epsilon) \sum_{i=1}^k |\alpha_i| \leq \|\sum_{i=1}^k \alpha_i x_i\|$ for any choice of scalars $\alpha_1, \alpha_2, \dots, \alpha_k$.

REFERENCES

1. D. R. Brown, *B-convexity and reflexivity in Banach spaces*, Trans. Amer. Math. Soc. **187** (1974), 69–76.
2. D. P. Giesy, *On a convexity condition in normed linear spaces*, Trans. Amer. Math. Soc. **125** (1966), 114–146; corrections, *ibid.* **140** (1969), 511–512. MR **34** #4866; **39** #1945.

DEPARTMENT OF MATHEMATICS, ANDHRA UNIVERSITY, WALT AIR 530003, INDIA