SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

ON A LIPSCHITZ ESTIMATE FOR CONFORMAL MAPS IN THE PLANE

R. MICHAEL RANGE

By combining a weak form of Koebe's $\frac{1}{4}$ -Theorem with elementary facts about subharmonic functions, one obtains a brief proof of the following statement.

THEOREM. *Suppose D and D^* are bounded domains in $\mathbb{C}$ with C^2 boundary. Then any conformal map $f: D \rightarrow D^*$ is Lipschitz, i.e. there is $K < \infty$, such that $|f(z) - f(\zeta)| \leq K|z - \zeta|$ for $z, \zeta \in D$.*

By a C^2 boundary I mean that ∂D is a not necessarily connected, compact, 1-dimensional C^2 submanifold of $\mathbb{C}$, or, equivalently, that ∂D consists of finitely many disjoint closed Jordan curves whose parametrizations in terms of arclength are twice continuously differentiable.

Stronger results, with more complicated proofs, have been known for a long time; for a more recent proof, one may consult Warschawski [4], where one can also find references to the older literature. I thank Professor H. Grunsky for pointing out this reference to me.

PROOF. Fix $z \in D$ and restrict f to the disc with center z and radius $d(z, \partial D)$. By Koebe's Theorem [1, Chapter IV, Satz 59], the image of this disc contains a disc with center $f(z)$ and radius $\rho \cdot |f'(z)| \cdot d(z, \partial D)$, where $\rho > 0$ is independent of z . This implies

$$(*) \quad |f'(z)| \leq 1/\rho \, d(f(z), \partial D^*) \cdot d(z, \partial D)^{-1} \quad \text{for } z \in D.$$

Choose a C^2 function ϕ in a neighborhood of $\bar{D}$, such that $\phi < 0$ on D , $\phi = 0$ on ∂D and $\text{grad } \phi \neq 0$ on ∂D . Locally, the existence of ϕ is an immediate consequence of the definition of a C^2 submanifold, and the global existence follows by a partition of unity argument. Replacing ϕ by $\exp(A\phi) - 1$ with sufficiently large $A > 0$, one can also assume that $\partial^2 \phi / \partial z \partial \bar{z} > 0$ near ∂D . Moreover, there is $k < \infty$ such that

$$(**) \quad |\phi(z)| \leq k \cdot d(z, \partial D) \quad \text{for } z \in D.$$

As suggested by Henkin [2], one now uses the following lemma.

Received by the editors March 18, 1975.

AMS (MOS) subject classifications (1970). Primary 30A30; Secondary 30A36, 31A05.

Copyright © 1976, American Mathematical Society

LEMMA (CF. [3]). Suppose $D \subset \mathbb{C}$ has C^2 boundary. Let u be continuous on $\overline{D}$, $u < 0$ on D , $u = 0$ on ∂D , and u is subharmonic on $U \cap D$ for some neighborhood U of ∂D . Then there is $c > 0$, such that

$$|u(w)| \geq c \cdot d(w, \partial D) \quad \text{for } w \in D.$$

Applying this Lemma to $\phi \circ f^{-1}$ and D^* , together with (**), one obtains $d(f(z), \partial D^*) \leq k/c \cdot d(z, \partial D)$. This estimate and (*) show that $|f'|$ is bounded, and hence f is Lipschitz. Q.E.D.

PROOF OF THE LEMMA. Choose $r > 0$, such that for each $\zeta \in \partial D$ the interior $\Delta_\zeta(r)$ of the circle of radius r through ζ with center on the inner normal to ∂D at ζ lies in $U \cap D$. Let $M = \sup\{u(z) : z \in D, d(z, \partial D) \geq r\}$; M is < 0 . For $z \in D$ with $d(z, \partial D) < r$ choose $\zeta \in \partial D$ with $|z - \zeta| = d(z, \partial D)$. Let P_r be the Poisson kernel for the disc $\Delta_\zeta(r)$, and let $d\sigma$ be arclength on $\partial\Delta_\zeta(r)$. Then

$$u(z) \leq \int_{\partial\Delta_\zeta} u(\lambda) P_r(z, \lambda) d\sigma(\lambda) \leq M \int_{\partial\Delta_\zeta \cap \{z: d(z, \partial D) \geq r\}} P_r(z, \lambda) d\sigma(\lambda),$$

and the Lemma now follows directly from properties of P_r .

REFERENCES

1. H. Behnke and F. Sommer, *Theorie der analytischen Funktionen einer komplexen Veränderlichen*, 2nd rev. ed., Die Grundlehren der math. Wissenschaften, Band 77, Springer-Verlag, Berlin, 1962. MR 26 #5137.
2. G. M. Henkin, *An analytic polyhedron is not holomorphically equivalent to a strictly pseudoconvex domain*, Dokl. Akad. Nauk SSSR 210 (1973), 1026–1029 = Soviet Math. Dokl. 14 (1973), 858–862. MR 48 #6467.
3. M. Keldyš and M. Lavrent'ev, *Sur l'unicité de la solution du problème de Neumann*, C. R. (Dokl.) Acad. Sci. URSS 26 (1937), 141–142.
4. S. E. Warschawski, *On differentiability at the boundary in conformal mapping*, Proc. Amer. Math. Soc. 12 (1961), 614–620. MR 24 #A1374.

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, ALBANY, NEW YORK 12222