ON 4-MANIFOLDS CROSS I

DAVID G. WRIGHT

ABSTRACT. In this note we set forth conditions under which for a given 4-manifold M there exist a countably infinite number of 4-manifolds M_i such that $\pi_1(\text{Bd } M_i)$ are distinct indecomposable groups and each $M_i \times I$ is homeomorphic with $M \times I$.

1. Introduction. Poenaru [6] and Mazur [4] have each shown that there exists a compact 4-manifold M such that $M \times I$ is homeomorphic with $B^4 \times I$, but M is not homeomorphic with B^4 . In this note we set forth conditions under which for a given 4-manifold M there exist a countably infinite number of 4-manifolds M_i such that $\pi_1(\text{Bd } M_i)$ are distinct indecomposable groups and each $M_i \times I$ is homeomorphic with $M \times I$. An interesting aspect of this note is that we never have to compute a fundamental group.

We express our indebtedness to D. R. McMillan, Jr. for discussions.

2. Definitions and notations. We let I , B^n , E^n and S^n denote the interval $[-1, 1]$, the n -ball $[-1, 1]^n$, Euclidean n -space, and the n -sphere respectively. If M is an n -manifold, then $\text{Int } M$ and $\text{Bd } M$ will denote the interior and boundary of M . The closure of a subset A of a topological space will be represented by $\text{Cl } A$. We recall that a 3-manifold is *irreducible* if every embedded S^2 bounds a B^3 . Let F be a surface in a 3-manifold M . If F is not a 2-sphere, then it is called *incompressible in M* if every simple closed curve on F which bounds an (open) disk in $M - F$ also bounds a disk in F .

3. Statement of the theorem.

THEOREM. Suppose M is a compact 4-manifold which is obtained from the 4-manifold N by adding a 2-handle H . If $\text{Cl}(\text{Bd } M - H)$ is an orientable, irreducible 3-manifold with incompressible boundary, then there exists a countably infinite collection of compact 4-manifolds M_i such that:

- (1) $\text{Bd } M_i$ is not homeomorphic with $\text{Bd } M_j$ for $i \neq j$,
- (2) $\pi_1(\text{Bd } M_i)$ is an indecomposable group and not infinite cyclic,
- (3) $\pi_1(\text{Bd } M_i) \neq \pi_1(\text{Bd } M_j)$ for $i \neq j$ and, hence, $\text{Int } M_i$ is not homeomorphic with $\text{Int } M_j$,
- (4) $M_i \times I$ is homeomorphic with $M \times I$.

Received by the editors January 2, 1975.

AMS (MOS) subject classifications (1970). Primary 57A15, 57C35; Secondary 57C40, 57A10.

Key words and phrases. Dunce hat, contractible 4-manifolds, homeomorphic under products.

© American Mathematical Society 1976

Examples of such manifolds are $(B^3 \times S^1) \cup H$ where H is a 2-handle attached to a solid torus T in $\text{Bd}(B^3 \times S^1) = S^2 \times S^1$ such that T has geometric index at least 2 in $S^2 \times S^1$.

To insure condition (4) we will need the following lemmas. We define $\text{id}: I \rightarrow I$ to be the identity map.

LEMMA 1. *Let $T_1, T_2 \subset \text{Int } B^3$ be solid tori. Then there exists a homeomorphism $h: T_1 \rightarrow T_2$ such that $h \times \text{id}: T_1 \times I \rightarrow T_2 \times I$ extends to a homeomorphism $H: B^3 \times [-2, 2] \rightarrow B^3 \times [-2, 2]$ which leaves the boundary of $B^3 \times [-2, 2]$ point-wise fixed.*

LEMMA 2. *Let $T_1, T_2 \subset \text{Bd } B^4$ be solid tori. Then there exists a homeomorphism $h: T_1 \rightarrow T_2$ such that $h \times \text{id}: T_1 \times I \rightarrow T_2 \times I$ extends to a homeomorphism $H: B^4 \times I \rightarrow B^4 \times I$.*

PROOF. Lemma 1 follows from well-known unknotting techniques. From Lemma 1 we obtain a homeomorphism $h: T_1 \rightarrow T_2$ such that $h \times \text{id}: T_1 \times I \rightarrow T_2 \times I$ extends to a homeomorphism $f: \text{Bd}(B^4 \times I) \rightarrow \text{Bd}(B^4 \times I)$ which in turn can be extended to a homeomorphism $H: B^4 \times I \rightarrow B^4 \times I$.

4. Construction of the M_i . We form 4-manifolds M_i by adding a pseudo 2-handle to N . Let H be the 2-handle and T_i be the regular neighborhood of the composite of i trefoil knots in $\text{Bd } H$. We set T to be $N \cap H$ which is an unknotted solid torus in $\text{Bd } H$. Let $h_i: T_i \rightarrow T$ be the homeomorphism promised by Lemma 2. Let $f: T \rightarrow N$ be the inclusion map. Then $M_i = N \cup_{f_i} H$ where $f_i = f \circ h_i$.

The boundary of M_i is the union of two irreducible 3-manifolds attached along a torus which is incompressible in each. Hence $\text{Bd } M_i$ is irreducible.

5. Proof of the theorem.

(1) The 3-manifold $\text{Bd } M_i$ has a Haken number [3] (maximal number of disjoint, nonparallel, incompressible surfaces) of at least $i + 1$ since there exist $i + 1$ disjoint, nonparallel, incompressible tori in $\text{Bd } H - T_i$. Hence we may pick out a subsequence of the M_i (which we still denote by M_i) such that for $i \neq j$, $\text{Bd } M_i$ and $\text{Bd } M_j$ have distinct Haken numbers. Hence $\text{Bd } M_i$ is not homeomorphic with $\text{Bd } M_j$.

(2) Suppose $\pi_1(\text{Bd } M_i)$ is decomposable or infinite cyclic. Then by [10] there exists an essential 2-sphere in $\text{Bd } M_i$. This contradicts the irreducibility of $\text{Bd } M_i$ and (2) is proved.

(3) Suppose $\pi_1(\text{Bd } M_i) = \pi_1(\text{Bd } M_j)$ for $i \neq j$. By the sphere theorem [5] $\pi_2(\text{Bd } M_i) = 0$. Hence by well-known techniques [8] there exists a map $f: \text{Bd } M_i \rightarrow \text{Bd } M_j$ which induces an isomorphism on fundamental group. Since $\text{Bd } M_i$ and $\text{Bd } M_j$ are irreducible and contain incompressible surfaces, we can apply Waldhausen's theorem [9, Theorem 6.1, p. 77] to conclude that $\text{Bd } M_i$ is homeomorphic with $\text{Bd } M_j$ which is a contradiction.

Since $\pi_1(\text{Bd } M_i) \neq \pi_1(\text{Bd } M_j)$, we conclude [1] that $\text{Int } M_i$ is not homeomorphic with $\text{Int } M_j$.

(4) We define a homeomorphism $h: M_i \times I \rightarrow M \times I$ by first noting that

$$M_i \times I = N \times I \cup_{f_i \times \text{id}} H \times I$$

and

$$M \times I = N \times I \cup_{f \times \text{id}} H \times I.$$

We define $h|N \times I$ to be the identity. By the construction of M_i the desired homeomorphism is guaranteed.

PROPOSITION. *There exists a 4-manifold, namely, $\text{Bd}(M \times I)$, in which M and each M_i can be embedded.*

This follows since $M_i \times \{-1\} \subset \text{Bd}(M_i \times I)$ which is homeomorphic to $\text{Bd}(M \times I)$. In fact, the double of each M_i is homeomorphic with $\text{Bd}(M \times I)$.

6. Contractible 4-manifolds.

DEFINITION. For each ordered n -tuple of integers $\beta = (m_1, \dots, m_n)$ such that $\sum m_i = 1$, let D_β be the contractible 2-complex formed by attaching a disk D to a circle α by the formula $\alpha^{m_1} \alpha^{m_2} \dots \alpha^{m_n}$. We call D_β a *generalized dunce hat*.

LEMMA 3. *If M is an n -dimensional manifold ($n \geq 5$) with a generalized dunce hat as spine, then M is an n -ball.*

We first note that a generalized dunce hat can be embedded in B^3 as a spine. To see this let A_1 and A_2 be transverse annuli on a torus T such that $A_1 \cap A_2$ is a disk. Let J_1 be an arc in A_1 with endpoints in $A_1 \cap A_2$ which goes around A_1 as described by β ; i.e., J_1 first winds around A_1 m_1 times, then m_2 times, etc. This is done in the most obvious manner so that J_1 can be completed to a simple closed curve J by connecting the endpoints of J_1 by an arc that runs around A_2 . Now let $f: T \rightarrow S^1$ be a map which wraps T around S^1 such that the inverse of a point is either A_2 or a simple closed curve parallel to A_2 . The mapping cylinder M_f of f is a solid torus with J a longitudinal curve of M_f . If care has been taken, $M_{f|J}$ is D_β minus a disk. However attaching a 2-handle to M_f along J yields B^3 .

Since D_β can be embedded as a spine of B^3 , by taking products, D_β can be embedded in B^n ($n \geq 3$) as a spine. However by Price [7], [2, Lemma 2] there is only one way to embed D_β in E^n ($n \geq 5$). Hence our lemma follows.

We are now in a position to prove Glaser's theorem [2] which states that there are uncountably many distinct contractible open 4-manifolds.

Let M be the 4-manifold M which is $B^3 \times S^1$ plus a 2-handle attached to a solid torus in $\text{Bd}(B^3 \times S^1)$ which has algebraic index 1 and geometric index not equal to 1. The manifold M has a generalized dunce hat as a spine. The proof of this fact is similar to the proof of [11, Theorem 5]. Hence $M \times I$ is homeomorphic with B^5 .

Now the theorem applied to the manifold M yields a countably infinite

collection of manifolds M_i with $\pi_1(\text{Bd } M_i)$ distinct indecomposable groups and not infinite cyclic. The existence of such a collection is the key step in Glaser's proof.

To finish the proof we form infinite sums M_α of the M_i 's in uncountably many different ways such that in two different ones M_i occurs more in one than in the other for some i . Since $\pi_1(\text{Bd } M_\alpha) \neq \pi_1(\text{Bd } M_\beta)$ for $\alpha \neq \beta$, $\text{Int } M_\alpha$ is not homeomorphic with $\text{Int } M_\beta$ [1].

REFERENCES

1. M. L. Curtis and K. W. Kwun, *Infinite sums of manifolds*, *Topology* **3** (1965), 31–42. MR **31** #729.
2. L. C. Glaser, *Uncountably many contractible open 4-manifolds*, *Topology* **6** (1966), 37–42. MR **34** #5072.
3. W. Haken, *Some results on surfaces in 3-manifolds*, *Studies in Modern Topology*, Math. Assoc. Amer.; distributed by Prentice-Hall, Englewood Cliffs, N.J., 1968, pp. 39–98. MR **36** #7118.
4. B. Mazur, *A note on some contractible 4-manifolds*, *Ann. of Math. (2)* **73** (1961), 221–228. MR **23** #A2873.
5. C. D. Papakyriakopoulos, *On Dehn's lemma and the asphericity of knots*, *Ann. of Math. (2)* **66** (1957), 1–26. MR **19**, 761.
6. V. Poenaru, *Les decompositions de l'hypercube en produit topologique*, *Bull. Soc. Math. France* **88** (1960), 113–129. MR **23** #A2872a.
7. T. M. Price, *Equivalence of embeddings of k -complexes in E^n for $n \leq 2k + 1$* , *Michigan Math. J.* **13** (1966), 65–69. MR **34** #3558.
8. J. R. Stallings, *On fibering certain 3-manifolds*, *Topology of 3-Manifolds and Related Topics* (Proc. Univ. of Georgia Inst., 1961), Prentice-Hall, Englewood Cliffs, N.J., 1962, pp. 95–100. MR **28** #1600.
9. F. Waldhausen, *On irreducible 3-manifolds which are sufficiently large*, *Ann. of Math. (2)* **87** (1968), 56–88. MR **36** #7146.
10. J. H. C. Whitehead, *On finite cocycles and the sphere theorem*, *Colloq. Math.* **6** (1958), 271–281. MR **20** #7271.
11. E. C. Zeeman, *On the dunce hat*, *Topology* **2** (1963), 341–358. MR **27** #6275.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824

Current address: Department of Mathematics, Utah State University, Logan, Utah 84322