COMPRESSIBLE MAPS

JAY E. GOLDFEATHER

ABSTRACT. Weingram has shown that if G is a finitely generated abelian group, then every nontrivial map $f: \Omega S^{2n+1} \rightarrow K(G, 2n)$ is incompressible; that is, f is not homotopic to a map whose image is contained in some finite-dimensional skeleton.

It is shown that a nontrivial map $\Omega S^{2n+1} \rightarrow K(G, 2n)$ may be compressible if G is not finitely generated. This result leads to some understanding of the obstructions to compressibility in Weingram's Theorem.

A map $f: X \rightarrow Y$ is said to compress into $A \subseteq Y$ if there is a map $f': X \rightarrow A$ such that $f \simeq i \cdot f'$, where i is inclusion. Weingram [4] has shown that if G is a finitely generated abelian group, then every nontrivial map $f: \Omega S^{2n+1} \rightarrow K(G, 2n)$ is incompressible, that is, f is not homotopic to a map whose image is contained in some finite-dimensional skeleton.

It will be shown that a nontrivial map $\Omega S^{2n+1} \rightarrow K(G, 2n)$ may be compressible if G is not finitely generated. Specifically, if P is a set of primes and $Z_P \subseteq Q$ is the subgroup of P -local integers, then any map $\Omega S^{2n+1} \rightarrow K(Q/Z_P, 2n)$ compresses into the $(2n+1)$ -skeleton, provided $2 \notin P$ or $n = 1, 3$.

Let $M(G, k)$ denote a Moore space of type (G, k) . For the remainder of this paper, it will be assumed that $2 \notin P$ or $2 \in P$ and $n = 1, 3$.

The following two well-known theorems will be stated without proof:

THEOREM (ADAMS [1]). $M(Z_P, 2n+1)$ is an H -space.

THEOREM (STASHEFF [3]). Let X and W be H -spaces and let $F: X \rightarrow W$ be an H -map. Let $\Omega W \rightarrow Y \rightarrow X$ be the fibration induced by f . Then Y is an H -space.

LEMMA 1. Any map $M(Z_P, 2n+1) \rightarrow K(Q, 2n+1)$ is homotopic to an H -map.

PROOF. The obstructions to a map $f: X \rightarrow Y$ being homotopic to an H -map lie in $H^i(X \wedge X; \pi_i(Y))$. Since

$$\pi_i(K(Q, 2n+1)) = 0 \quad \text{for } i \neq 2n+1$$

and

$$H^{2n+1}(M(Z_P, 2n+1) \wedge M(Z_P, 2n+1); G) = 0 \quad \text{for any } G,$$

all obstructions lie in zero groups.

Received by the editors January 29, 1976 and, in revised form, March 22, 1976.

AMS (MOS) subject classifications (1970). Primary 55G36; Secondary 55F05, 55D45.

Key words and phrases. H -space mod p .

Copyright © 1977, American Mathematical Society

In view of the preceding lemma, it will be assumed that any given map $f: M(Z_p, 2n+1) \rightarrow K(Q, 2n+1)$ is an H -map.

Let f be induced by the natural embedding $Z_p \subseteq Q$.

PROPOSITION 2. *Let $K(Q, 2n) \rightarrow E \rightarrow M(Z_p, 2n+1)$ be the fibration induced by f . Then E is an H -space.*

PROOF. This is an immediate consequence of the above two theorems and Lemma 1.

THEOREM 3.

$$\tilde{H}_i(E; Z) = \begin{cases} Q/Z_p, & i = 2n, \\ 0, & \text{otherwise.} \end{cases}$$

PROOF. (Porter [2] has proved a similar theorem when the base space is S^3 .) $H_*(K(Q, 2n); Z)$ is a polynomial algebra over Z on one generator in dimension $2n$ tensored with Q . Since f is induced by the natural embedding of $Z_p \subseteq Q$, the Serre exact sequence

$$H_{2n+1}(E) \rightarrow H_{2n+1}(M(Z_p, 2n+1)) \rightarrow H_{2n}(K(Q, 2n)) \rightarrow H_{2n}(E) \rightarrow 0$$

reduces to

$$0 \rightarrow Z_p \rightarrow Q \rightarrow H_{2n}(E) \rightarrow 0$$

and, hence, $H_{2n}(E) = Q/Z_p$.

In the homology spectral sequence, the only nonzero differential is d^{2n+1} which is an isomorphism for total degrees greater than $2n+1$ since $Z_p \otimes Q \cong Q$. Hence, $H_i(E) = 0$ for $i > 2n+1$.

REMARK. E is homotopic to a $(2n+1)$ -dimensional Moore space

$$M(Q/Z_p, 2n).$$

PROPOSITION 4. $K(Q/Z_p, 2n)$ is not homotopic to a finite-dimensional complex.

PROOF. The exact sequence $0 \rightarrow Z_p \rightarrow Q \rightarrow Q/Z_p \rightarrow 0$ induces a fibration

$$K(Q/Z_p, 2n) \rightarrow K(Z_p, 2n+1) \rightarrow K(Q, 2n+1).$$

It is well known that $K(Q, 2n+1)$ is homotopic to a finite-dimensional complex. Hence if $K(Q/Z_p, 2n)$ were also homotopic to a finite-dimensional complex, it would imply that $H^k(K(Z_p, 2n+1); Z_p) = 0$ for all k greater than some integer. It suffices to show, then, that $H^k(K(Z_p, 2n+1); Z_p)$ is nonzero for infinitely many k , where $p \in P$.

Let ι be the fundamental class in $H^{2n+1}(K(Z, 2n+1); Z_p) = \text{Hom}(Z, Z_p)$. It is well known that there are an infinite number of Steenrod operations $\mathcal{P}^I$ such that $\mathcal{P}^I \iota \neq 0$. (I denotes an admissible sequence.) Hence it suffices to show that if $f: K(Z, 2n+1) \rightarrow K(Z_p, 2n+1)$ is the natural inclusion, then

$f^*(i) = i$ for some $i \in H^{2n+1}(K(Z_p, 2n+1); Z_p) = \text{Hom}(Z_p, Z_p)$. But $f^*: \text{Hom}(Z_p, Z_p) \rightarrow \text{Hom}(Z, Z_p)$ is onto and the proposition is proved.

THEOREM 5. *Every nontrivial map $\Omega S^{2n+1} \rightarrow K(Q/Z_p, 2n)$ compresses into the $(2n+1)$ -skeleton.*

PROOF. Any map $f: \Omega S^{2n+1} \rightarrow K(Q/Z_p, 2n)$ is homotopic to a map $r \cdot \Omega Sg$, where $g: S^{2n} \rightarrow K(Q/Z_p, 2n)$ and r is the retraction $\Omega SK(Q/Z_p, 2n) \rightarrow K(Q/Z_p, 2n)$.

But every such map g factors through $M(Q/Z_p, 2n)$ so that

$$S^{2n} \xrightarrow{g_1} M(Q/Z_p, 2n) \xrightarrow{g_2} K(Q/Z_p, 2n)$$

is nontrivial where

$$g \simeq g_2 \cdot g_1.$$

By Theorem 3 and Proposition 2, $M(Q/Z_p, 2n)$ is an H -space so that g_1 extends to $\bar{g}_1: \Omega S^{2n+1} \rightarrow M(Q/Z_p, 2n)$. Then $f \simeq g_2 \cdot \bar{g}_1$, so f compresses into the $(2n+1)$ -skeleton.

Observe that if $P = \{p\}$, then $Q/Z_p = Z_{p^\infty} = \varinjlim_r Z_{p^r}$ induced by the inclusion $Z_{p^r} \rightarrow Z_{p^{r+1}}$. Let G be a finitely-generated odd torsion group so that $G = \bigoplus_{i=1}^m Z_{p_i^{a_i}}$. Let $P = \{p_1, \dots, p_m\}$ and $G_{k+1} = \bigoplus_{i=1}^m Z_{p_i^{a_i+k}}$. Then $Q/Z_p \cong \varinjlim_k G_k$.

Since $\pi_{2n+1}(M(G_r, 2n)) = G_r \otimes Z_2 = 0$, and $M(G_r, 2n)$ can be thought of as the $(2n+1)$ -skeleton of $K(G_r, 2n)$, the obstructions to extending

$$S^{2n} \rightarrow K(G_r, 2n)^{(2n+1)}$$

to $\Omega S^{2n+1} \rightarrow K(G_r, 2n)^{(2n+1)}$ are the obstructions to extending

$$S^{2n} \rightarrow M(G_r, 2n)$$

to $\Omega S^{2n+1} \rightarrow M(G_r, 2n)$.

Let $j_{r,m}: M(G_r, 2n) \rightarrow M(G_{r+m}, 2n)$ be inclusion and let $j: S^{2n} \rightarrow M(G_r, 2n)$.

THEOREM 6. *For every k and r , there is an m such that $j_{r,m} \cdot j$ extends to $(S^{2n})_k$, the k th reduced product of S^{2n} .*

PROOF. Let $j_{r,\infty}: M(G_r, 2n) \rightarrow M(Q/Z_p, 2n)$ be the inclusion. Since $M(Q/Z_p, 2n)$ is an H -space, it follows that $j_{r,\infty} \cdot j$ extends to $(S^{2n})_k$ for all k . Let $\bar{j}$ be such an extension. Since $(S^{2n})_k$ is compact, $\text{image } \bar{j} \subseteq M(G_{r+m}, 2n)$ for some m and, hence, $\bar{j}$ is an extension of $j_{r,m} \cdot j$.

BIBLIOGRAPHY

1. J. F. Adams, *The sphere, considered as an H -space mod p* , Quart J. Math. Oxford Ser. (2) **12** (1961), 52–60. MR **23** #A651.
2. R. D. Porter, *An H -space with finite dimensional homology whose loop space has torsion*, Proc. Amer. Math. Soc. **37** (1973), 291–292. MR **46** #9970.
3. J. D. Stasheff, *On extensions of H -spaces*, Trans. Amer. Math. Soc. **105** (1962), 126–135. MR **31** #2726.
4. S. Weingram, *On the incompressibility of certain maps*, Ann. of Math. (2) **93** (1971), 476–485. MR **46** #890.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MILWAUKEE, WISCONSIN 53201