COHOMOLOGY OF ASSOCIATIVE TRIPLE SYSTEMS

RENATE CARLSSON

ABSTRACT. In the following a cohomology for associative triple systems is defined. The classical corollaries can be extended to the associative triples.

1. Introduction. One distinguishes two associative triple systems. The associative triples of the first kind or ternary rings were introduced by Lister [6]. Those of the second kind or Hestenes ternary rings were first investigated by Hestenes and as special alternative triples by Loos [7]. Typical examples for the latter are the $p \times q$ matrices in an associative algebra with an involution, the triple product given by $(x, y, z) \mapsto x' \bar{y} z$, $\bar{y}$ the transposed-conjugate matrix.

In this note we define a canonic cohomology for associative triples. Cohomologies are well known for certain algebras and Lie triples [2]. The corollaries for associative triples are analogous to the classical results.

In the following K is a commutative and associative ring with unit 1, the modules taken unitary over K . If K is explicitly assumed as a field then the dimensions over K are finite.

2. Definitions. The universal (j -) imbedding. A K -module A with a trilinear inner composition $A \times A \times A \rightarrow A$, with $(x, y, z) \mapsto \langle xyz \rangle$, is an *associative triple system of the 1st kind* if

$$\langle \langle xyz \rangle uv \rangle = \langle x \langle yzu \rangle v \rangle = \langle xy \langle zuv \rangle \rangle$$

for $x, y, z, u, v \in A$. A is an *associative triple of the 2nd kind*, when

$$\langle \langle xyz \rangle uv \rangle = \langle x \langle uzy \rangle v \rangle = \langle xy \langle zuv \rangle \rangle.$$

A *trimodule* M over a triple system A is a K -module M together with 3 trilinear compositions $A \times A \times M \rightarrow M$, the products denoted $\langle xym \rangle$, $\langle xmy \rangle$, $\langle mxy \rangle$, where $m \in M$. The *split null extension* or *semidirect sum* $A + M$ of M over A is the K -module direct sum with

$$\langle (x + m_1)(y + m_2)(z + m_3) \rangle := \langle xyz \rangle + \langle m_1 yz \rangle + \langle x m_2 z \rangle + \langle x y m_3 \rangle$$

for $x, y, z \in A$, $m_i \in M$, $i = 1, 2, 3$. A trimodule M is *associative of the 1st kind* over A if the semidirect sum is an associative triple of the 1st kind, equally for the 2nd kind. In the following A always denotes an associative triple, and M an associative trimodule over A of the same kind if nothing

Received by the editors September 16, 1975.

AMS (MOS) subject classifications (1970). Primary 16A78, 18H15, 18H40; Secondary 08A25, 16A21, 16A28.

Key words and phrases. Cohomology groups of associative triple systems, universal (j -) imbedding, Whitehead lemmata, Wedderburn principal theorem, invariant submodules, trimodules.

Copyright © 1977, American Mathematical Society

other is stated, where associative means associative of the 1st or 2nd kind. For K -modules N, P let $N \oplus P$ denote the K -module direct sum.

Let B be an associative algebra, the multiplication designed by " $\circ$ ", and $\iota: A \rightarrow B$ a K -module monomorphism. (B, ι) is an *imbedding* of an associative triple A of the 1st kind if $B = \iota A + \iota A \circ \iota A$ with $\iota x \circ \iota y \circ \iota z = \iota \langle xyz \rangle$, $x, y, z \in A$, and the imbedding is *direct* if the sum is direct [6]. (B, ι) is *universal* if for each homomorphism $f: A \rightarrow A'$ onto a triple A' , and any imbedding (B', κ) of A' with monomorphism κ there exists a unique algebra homomorphism $\tilde{f}: B \rightarrow B'$ onto B' so that the diagram

$$\begin{array}{ccc} B & \xrightarrow{\tilde{f}} & B' \\ \iota \uparrow & & \uparrow \kappa \\ A & \xrightarrow{f} & A' \end{array}$$

is commutative, $\hookrightarrow$ denoting a K -module monomorphism.

Let A be associative of the 2nd kind, j an involution of B with $\overline{\iota A} := j(\iota A)$, $\bar{b} := j(b)$, $b \in B$. (B, ι, j) is a *j -imbedding* of A if $B = \iota A + \overline{\iota A} + \iota A \circ \iota A + \iota A \circ \iota A$ with $\iota A \circ \iota A = \{0\}$ and $\iota x \circ \bar{\iota y} \circ \iota z = \iota \langle xyz \rangle$ for $x, y, z \in A$. The *j -imbedding* is *direct* if the above sum is direct. (B, ι, j) is *universal* if for any homomorphism f onto A' , (B', κ, j') a *j' -imbedding* of A' , there exists a unique homomorphism $\tilde{f}: B \rightarrow B'$ so that the above diagram is commutative and $j'\tilde{f} = \tilde{f}j$. The standard imbedding of A is a direct imbedding if A is associative of the 1st kind [6], and it contains a direct *j -imbedding* otherwise, as is easily noticed (cf. [7]). One immediately has

THEOREM 1. *Let A be an associative triple over K . Then there exists a universal imbedding $(U(A), \iota)$, respectively a universal j -imbedding $(U(A), \iota, j)$ of A . If $(V(A), \kappa)$, respectively $(V(A), \kappa, j')$, is a universal (j' -) imbedding too, then there is a unique isomorphism g of $U(A)$ onto $V(A)$ with $\kappa = g\iota$.*

PROOF. Let $T(A) := K \cdot 1 \oplus T^*(A)$ be the tensor algebra of the K -module A with $T^*(A) := \bigoplus (\otimes^n A)$, $n \in \mathbb{N}$. Then $T(A)$ is a universal enveloping associative algebra for A with unit 1. We construct $U(A)$ as a factor algebra: In case of the 2nd kind let $\bar{A}$ be a K -module isomorphic to A , the isomorphism given by $x \mapsto \bar{x}$. Let Q denote the ideal of $T^*(A \oplus \bar{A})$ generated by $x \otimes y, \bar{x} \otimes \bar{y}, x \otimes \bar{y} \otimes z - \langle xyz \rangle, \bar{x} \otimes y \otimes \bar{z} - \langle \bar{z}yx \rangle$ for $x, y, z \in A$. Set $U(A) := T^*(A \oplus \bar{A})/Q$ and $\iota: x \mapsto x + Q$. Observing the elementary properties of the tensor product one notes the existence of an involution j of $U(A)$ with $j(x + Q) = \bar{x} + Q$. Thus $(U(A), \iota, j)$ is a universal *j -imbedding* of A . For the 1st kind take $T^*(A)$ modulo the ideal Q generated by the elements $x \otimes y \otimes z - \langle xyz \rangle$.

The proof of the second part of the statement is standard. $\square$

In the subsequent we may suppose that ι is the inclusion map. For an algebra C let $C^{[n]}$ denote the n -fold cartesian product. Let $U(A)$ design a universal (*j -) imbedding* of A . Obviously $U(A)$ is direct.

For a direct imbedding $B = A \oplus A \circ A$ let $\epsilon_1: B \rightarrow A, \epsilon_2: B \rightarrow A \circ A$ be

the canonic projections. If B is a direct j -imbedding let $\varepsilon_1, \varepsilon_2, \varepsilon_{-1}, \varepsilon_{-2}$ be its canonic projections onto $A, A \circ \bar{A}, \bar{A}, \bar{A} \circ A$. Set $B_k := \varepsilon_k(B)$, $k \in J := \{1, 2\}$, respectively $J := \{-2, -1, 1, 2\}$. The *formal multiplication* of the B -components B_k is obviously declared, e.g. for the 2nd kind $A \cdot (\bar{A} \circ A) := A$, $(A \circ \bar{A}) \cdot (A \circ \bar{A}) := A \circ \bar{A}$, $A \cdot \bar{A} := \{0\}, \dots$. Similar to the case of Lie triples [2] we will consider some homogeneous multilinear mappings of universal (j -) imbeddings. For $x := (x_1, \dots, x_n) \in B^{[n]}$ with $x \in B_k$, $k = k(i)$, the B -component is $\Phi(x) := \prod B_{k(i)}$, $i = 1, \dots, n$, the product formal. Let C be a subalgebra of B with $\varepsilon_k(C) \subset C$, and $B' a$ (j' -) imbedding of A' with A' of the same kind as A . Then $f: C^{[n]} \rightarrow B'$ is *isovarying* if

$$\Phi(x) \subset B_k \Rightarrow f(x) \subset B'_k,$$

and *antivarying* if for $\Phi(x) \neq \{0\}$

$$\Phi(x) \subset B_k \Rightarrow f(x) \subset \bigoplus B'_l, \quad l \neq k, l \in J.$$

Any f has a unique decomposition $f = f_1 + f_2$, f_1 isovarying and f_2 antivarying. Denote $\varepsilon: f \mapsto f_1$. Then $\varepsilon^2 := \varepsilon \circ \varepsilon = \text{id}$.

Let $B := A + M$, and let J be the ideal in $U(B)$ equal to

$$\{a \in B \circ M + M \circ B \mid a \circ A = A \circ a = \{0\}\},$$

respectively in the case of the 2nd kind

$$\begin{aligned} \{a \in B \circ \bar{M} + M \circ \bar{B} + \bar{B} \circ M + \bar{M} \circ B \mid a \circ (A + \bar{A}) \\ = (A + \bar{A}) \circ a = \{0\}\}. \end{aligned}$$

Then the *universal module imbedding* M_U is the ideal of $U(A + M)$ generated by M , respectively $M \oplus \bar{M}$, in the case of 2nd kind associativity, modulo J . The maps corresponding to ε and ε, j for the 2nd kind after the passages from $U(A + M)$ to $U(A)$ and M_U are denoted equally. M_U is an associative bimodule over $U(A)$ in the obvious way.

3. The cohomology. Let C^n be the linear space of the n -cochains of $U(A)$ for M_U [3], $n \in \mathbb{N}_0$. These are the n -linear mappings of $U(A)^{[n]}$ into M_U for $n \in \mathbb{N}$, while $C^0 = M_U$. The coboundary operator is defined by

$$\delta f(a_1) = a_1 \cdot f - f \cdot a_1, \quad \text{if } f \in C^0,$$

$$\begin{aligned} \delta f(a_1, \dots, a_{n+1}) &= a_1 \cdot f(a_2, \dots, a_{n+1}) \\ &+ \sum_{i=1}^n (-1)^i f(a_1, \dots, a_i \circ a_{i+1}, \dots, a_{n+1}) \\ &+ (-1)^{n+1} f(a_1, \dots, a_n) \cdot a_{n+1} \end{aligned}$$

$$\text{for } f \in C^n, n \neq 0, a_i \in U(A).$$

Z^n, B^n denote the K -modules of the n -cocycles and the n -coboundaries where $B^0 := \{0\}$. $H^n(U(A), M_U) := Z^n / B^n$ is the n -dimensional cohomology group. If $x = (x_1, \dots, x_n) \in U(A)^{[n]}$, $n \neq 0$, $U(A)$ a j -imbedding, set $j(x) := (j(x_n), \dots, j(x_1))$. For $f \in C^0$ extend ε by $\varepsilon(f) := \varepsilon_2 f$ if A is of the 1st kind, and if A is of the 2nd kind let $\varepsilon(f) := (\varepsilon_2 + \varepsilon_{-2})f$, $\bar{j}f := f$. Then the K -module

of the n -cochains of the associative triple system A for M is

$$C^n(A, M) := \{f \in C^n \mid \varepsilon(f) = f\}$$

for $n \in \mathbb{N}_0$ if A and M are associative of the 1st kind. In case of the 2nd kind define

$$C^n(A, M) := \{f \in C^n \mid \varepsilon(f) = f, jff = (-1)^{n(n+1)/2+1}f\}$$

for $n \in \mathbb{N}_0$ if $\frac{1}{2} \in K$. If $2 \cdot K = \{0\}$ modify this definition by $C^0(A, M) := \{f \in C^0 \mid \varepsilon(f) = f, jf = f + z \text{ with } z \in M_U, a \cdot z = z \cdot a \text{ for all } a \in U(A)\}$.

The coboundary operator ∂ is the restriction of δ from C^n onto $C^n(A, M)$. The modules $C^n(A, M)$ together with ∂ define a cochain complex. $Z^n(A, M)$ is the submodule of n -cocycles of $C^n(A, M)$, $B^n(A, M)$ that of n -coboundaries. Let $H^n(A, M)$ be the n -dimensional cohomology group with $H^n(A, M) := Z^n(A, M)/B^n(A, M)$. Set $Z_+^n := Z^n \cap C^n(A, M)$, and let $B_+^n := B^n \cap C^n(A, M)$. We have

LEMMA 1. Let A be an associative triple system, M an associative trimodule over A , and $\frac{1}{2} \in K$ in case of the 2nd kind. Then

$$H^n(A, M) = Z_+^n / B_+^n, \quad \text{and} \quad H^n(U(A), M_U) \cong H^n(A, M) \oplus H(n),$$

$H(n)$ a submodule of $H^n(U(A), M_U)$, $n \in \mathbb{N}_0$.

PROOF. Let A and M be associative of the 2nd kind, and $\frac{1}{2} \in K$. Then for $f \in C^n$ we have $f = \varepsilon(f) + (f - \varepsilon(f))$ and $f = \frac{1}{2} \cdot ((f + jff) + (f - jff))$. These are the decompositions of f into eigenfunctions for the endomorphisms ε and $\tilde{j}$: $f \mapsto jff$. An easy verification shows that ε and $\tilde{j}$ commute. Thus there is a weight space decomposition $C^n = \bigoplus C_{\alpha, \beta}^n$, α the root of ε , β the root of $\tilde{j}$ for $C_{\alpha, \beta}^n$. Now $C^n(A, M) = C_{1, (-1)^{n(n+1)/2+1}}^n$. Since $\delta: C_{\alpha, \beta}^{n-1} \rightarrow C_{\alpha, \beta}^n$, $\beta' = \beta \cdot (-1)^n$, we get $B^n(A, M) = B_+^n$. The rest is obvious. For the case of the 1st kind one proceeds similarly and even more immediately. $\square$

The associative triple A is separable over the field K , if for every base field extension $\tilde{K}$ of K the base field extension $\tilde{A}$ of A is semisimple. Note that as in the binary case the radical of a ternary algebra A coincides with the radical of the ternary ring A when $\frac{1}{2} \in K$ [8]. We receive

COROLLARY 1. Let M be an associative trimodule of the 1st or 2nd kind over A , and A separable over a field K , $\text{char}(K) \neq 2$. Then for all $n \in \mathbb{N}$, $H^n(A, M) = \{0\}$.

PROOF. Because A is semisimple the standard imbedding of A is semisimple too by [6], respectively [7]. Any semisimple imbedding, respectively j -imbedding, is isomorphic to $U(A)$. The isomorphism follows as the corresponding one in case of Lie triples [4]. Thus $U(A)$ is separable and

$$H^n(U(A), M_U) = \{0\} \quad \text{for all } n \in \mathbb{N}. \quad \square$$

Corollary 1 includes the statements corresponding to the two Whitehead lemmas.

A linear mapping $D: A \rightarrow M$ is a derivation of A in M if

$$D(\langle xyz \rangle) = \langle D(x)yz \rangle + \langle xD(y)z \rangle + \langle xyD(z) \rangle, \quad x, y, z \in A.$$

Inner derivations are those generated by the triple multiplications in the semidirect sum $A + M$.

If $\text{Der}(A, M)$ is the K -module of derivations of A in M , then we have $Z^1(A, M) \cong \text{Der}(A, M)$, the elements of $B^1(A, M)$ related to inner derivations of A in M . As isomorphism of $Z^1(A, M)$ onto $\text{Der}(A, M)$ we may choose the restriction of the elements $f \in Z^1(A, M)$ from $U(A)$ onto A . It follows immediately that in case of the 2nd kind, $\text{char}(K) = 2$ and $n = 1$, Corollary 1 is valid too. Hence

COROLLARY 2. *Let A and M be associative of the same kind, A separable over the field K , and $\text{char}(K) \neq 2$ in case of the 1st kind. Then every derivation of A in M is inner.* $\square$

We may remark that in case of $\text{char}(K) = 2$ the trivial 1st kind triple $A := \{0, 1\}$ with $\langle 111 \rangle = 1$ is separable, while the radical of $U(A)$ is $\{0, 1 + 1 \circ 1\} \neq \{0\}$.

Now let C denote a binary algebra, N a C -bimodule. A *semiautomorphism* f of N is an algebra automorphism of the semidirect sum $C + N$ with $f(A) \subset A$, $f(N) \subset N$. It holds:

LEMMA 2. *Let G be a finite group of semiautomorphisms of a bimodule N over an algebra C , n the order of G , and $1/n \in K$. If $N = Q \oplus P$, Q and P are C -submodules with $G(Q) = Q$, then there exists a C -submodule R of N with $G(R) = R$ and $N = Q \oplus R$.* $\square$

The proof follows considering the semidirect sum $C + N$ and going on along the lines of [10].

Irreducible trimodules, complete reducibility, triple (trimodule) homomorphisms are defined as obvious. If A is semisimple, Z the center of $U(A)$, and $Z \otimes Z$ semisimple, A is *c-semisimple*. Denote $W := U(A) \otimes U(A)^{\text{op}}$, $U(A)^{\text{op}}$ the opposite algebra for $U(A)$. For c -semisimple A , W is semisimple. For a direct imbedding B let σ be the natural algebra automorphism with $\sigma(x) = -x$ and fixing $x \circ y$ when $x, y \in A$.

PROPOSITION 1. *Let M be an associative trimodule over A , K a field with $\text{char}(K) \neq 2$, and A c -semisimple. Then M is completely reducible over K .*

PROOF. M_U is an associative left W -module by $x \otimes y \cdot m = x \circ m \circ y$ with $x, y \in U(A)$, $m \in M_U$. Then $M_U = e \circ M_U \circ e \oplus M_0$, e the unit of $U(A)$, M_0 the maximal zero W -submodule, is a decomposition into W -submodules. Set $M := e \circ M_U \circ e$. It may be noted that for the 2nd kind $e = e_1 + e_2$ with $e_1 \in A \circ \bar{A}$, $e_2 \in \bar{A} \circ A$, so that $\varepsilon_1(M) = e_1 \circ M_U \circ e_2$. Thus $M = \varepsilon_1(M) \oplus \varepsilon_1(M_0)$ is a decomposition of M into subtrimodules. Consider now $\varepsilon_1(M_0)$ as a left- and a right-module over the semisimple algebra $A \circ \bar{A}$ over which it decomposes into irreducible submodules [6], and equally for $A \circ \bar{A} \oplus \bar{A} \circ A$ in case of the 2nd kind. Hence $\varepsilon_1(M_0)$ is completely reducible over A .

Set $G := \{\text{Id}, \sigma\}$ for the 1st kind. Then $G(M) \subset M$. If $M \neq \{0\}$, M decomposes by Lemma 2 or [5, p. 117] into minimal G -invariant submodules $P_k \neq \{0\}$ over W , $k = 1, \dots, t$. Therefore $\varepsilon_1(P_k) \subset P_k$. Hence $N_k := \varepsilon_1(P_k)$ is an irreducible subtrimodule. Thus M is a completely reducible A -trimodule.

$\square$

The proposition says equivalently that under its conditions on A any short exact sequence of associative A -trimodules of the same kind, the connecting mappings being trimodule homomorphisms, splits by a trimodule homomorphism.

An *extension of the associative triple A by M* is a short exact sequence of associative triples M, E, A of the same kind $\{0\} \rightarrow M \xrightarrow{e} E \xrightarrow{p} A \rightarrow \{0\}$ with e, p triple homomorphisms. M is in a canonic way a trimodule over A . Let ${}^{(1)}M := \langle MMA \rangle + \langle MAM \rangle + \langle AMM \rangle$. The extension is *singular* if ${}^{(1)}M = \langle MMM \rangle = \{0\}$. The extension is *split* or *trivial* if there exists a triple homomorphism $q: A \rightarrow E$, with $pq = \text{Id}_A$. The extension $\{0\} \rightarrow M \xrightarrow{f} E' \xrightarrow{q} A \rightarrow \{0\}$ is *equivalent* to the first if there is a triple homomorphism $k: E \rightarrow E'$ so that the diagram

$$\begin{array}{ccccccc} & & & E & & & \\ & & e \nearrow & \downarrow k & \searrow p & & \\ \{0\} \rightarrow & M & & & & A & \rightarrow \{0\} \\ & & f \searrow & \downarrow q & \nearrow & & \\ & & & E' & & & \end{array}$$

is commutative. Then k is an isomorphism. Let J' be the ideal of $U(E)$ defined as J , substituting $e(M)$, E for M and $A + M$. Set $U(E)' := U(E)/J'$. Design the maps induced by ε and j on $U(E)'$ equally. For a singular extension there is a commutative diagram:

$$\begin{array}{ccccccccc} \{0\} & \longrightarrow & M & \xrightarrow{e} & E & \xrightarrow{p} & A & \longrightarrow & \{0\} \\ & & \downarrow & & \downarrow & & \downarrow & & \\ \{0\} & \longrightarrow & M_U & \xrightarrow{e'} & U(E)' & \xrightarrow{p'} & U(A) & \longrightarrow & \{0\} \end{array}$$

e', p' algebra homomorphisms invariant relative to ε and respectively for j . Note that the second row is also exact (cf. [2, pp. 158–159]).

A *factor set f of A in M* is, as usual, a trilinear mapping $f: A \times A \times A \rightarrow M$ so that the K -module direct sum $A + M$ with $[xyz] := \langle xyz \rangle + f(x, y, z)$, ${}^{(1)}M := \{0\}$, $\langle MMM \rangle := \{0\}$, and the trimodule structure of M , is an associative triple of the same kind. f is *trivial* if there is $s: A \rightarrow M$, s K -linear with

$$f(x, y, z) = s(\langle xyz \rangle) - \langle s(x)yz \rangle - \langle xs(y)z \rangle - \langle xys(z) \rangle.$$

Let A be K -projective in the usual sense. Then any extension of the triple A is split for the K -module A . Similarly as for algebras, the classes of equivalent singular extensions of A by M constitute a K -module isomorphic to the K -module of factor sets, the trivial factor sets corresponding to the classes of the split extensions. One then obtains, arguing as [2, pp. 159–160] for Lie triples,

THEOREM 2. *If A and the trimodule M over A are associative of the same kind, A K -projective, and $\frac{1}{2} \in K$ in case of the 2nd kind, then there is a module isomorphism $H^2(A, M) \cong E(A, M)/E_t(A, M)$, $E(A, M)$ the K -module of the classes of equivalent singular extensions of A with kernel M , $E_t(A, M)$ the submodule of the classes of split extensions. $\square$*

If $A^{(1)} := \langle AAA \rangle$, $A^{(n+1)} = (A^{(n)})^{(1)}$, $n \in \mathbb{N}$, then A is solvable when $A^{(m)} = \{0\}$ for some m . Set for an ideal I in A , $^{(n+1)}I := ^{(1)}(^{(n)}I)$. I is S -solvable if for one m , $^{(m)}I = \{0\}$. Because $R := \text{Rad}(A) \subset \text{Rad}(U(A))$, R is solvable if K is a field. We have for arbitrary K

LEMMA 3. Let I be an ideal of A . Then:

(a) I is solvable exactly if it is S -solvable.

(b) When I is solvable, $I \neq \{0\}$, then $^{(1)}I \subsetneq I$.

PROOF. (a) From $B^{(1)} \subset ^{(1)}B$ we get $B^{(n)} \subset ^{(n)}B$ for any $n \in \mathbb{N}$. Further one verifies $^{(3)}B \subset B^{(1)}$. Thus $^{(3n)}B \subset B^{(n)}$. Hence we get (a). (a) $\Rightarrow$ (b). $\square$

The proof of the Wedderburn principal theorem reduces by Lemma 3 to the singular case. For associative triples of the 2nd kind the statement is implied by [1]. On behalf of Corollary 1 it follows:

COROLLARY 3. Let A be a finite dimensional associative triple system over a field K , R the radical, A/R separable, and $\text{char}(K) \neq 2$. Then A decomposes $A = B \oplus R$, B a subtriple of A , $B \cong A/R$. $\square$

REFERENCES

1. R. Carlsson, *Der Wedderburnsche Hauptsatz für alternative Paare* (to appear).
2. B. Harris, *Cohomology of Lie triple systems and Lie algebras with involution*, Trans. Amer. Math. Soc. **98** (1961), 148–162. MR **22** #11068.
3. G. Hochschild, *On the cohomology groups of an associative algebra*, Ann. of Math. (2) **46** (1945), 58–67. MR **6**, 114.
4. N. Jacobson, *General representation theory of Jordan algebras*, Trans. Amer. Math. Soc. **70** (1951), 509–530. MR **12**, 797.
5. ———, *Structure of rings*, rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 37, Amer. Math. Soc., Providence, R. I., 1964. MR **36** #5158.
6. W. G. Lister, *Ternary rings*, Trans. Amer. Math. Soc. **154** (1971), 37–55. MR **42** #7716.
7. O. Loos, *Assoziative Tripelsysteme*, Manuscripta Math. **7** (1972), 103–112. MR **46** #3581.
8. H. C. Myung, *A characterization of the Jacobson radical in ternary algebras*, Proc. Amer. Math. Soc. **38** (1973), 228–234. MR **49** #363.
9. P. Seibt, *Cohomology of algebras and triple systems*, Comm. Algebra **3** (1975), 1097–1120.
10. E. J. Taft, *Cleft algebras with operator groups*, Portugal. Math. **20** (1961), 195–198. MR **25** #2103.

UNIVERSITÄT HAMBURG, MATHEMATISCHES SEMINAR, BUNDESSTRASSE 55,2 HAMBURG 13,
FEDERAL REPUBLIC OF GERMANY