SHORTER NOTES

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is no other outlet.

ON THE COMMUTATION THEOREM FOR TENSOR PRODUCTS OF VON NEUMANN ALGEBRAS

R. ROUSSEAU, A. VAN DAELE AND L. VANHEESWIJCK

ABSTRACT. It is shown that the commutation theorem for tensor products of general von Neumann algebras follows trivially from the case of von Neumann algebras with a separating and cyclic vector.

Let M_1 and M_2 be von Neumann algebras acting in Hilbert spaces $\mathcal{H}_1$ and $\mathcal{H}_2$ respectively. Then the commutation theorem for tensor products of von Neumann algebras states

$$(M_1 \otimes M_2)' = M_1' \otimes M_2'.$$

The proof of this theorem in full generality was first obtained by Tomita in 1967 (see [3]). Later a number of simpler versions have been obtained (see e.g. [2]). Usually standard techniques are used to reduce the general case to the case of von Neumann algebras having a cyclic and separating vector. In this note we show that this reduction is almost a triviality.

Assume that the commutation theorem has been proved for von Neumann algebras with a separating and cyclic vector. To show that $(M_1 \otimes M_2)' = M_1' \otimes M_2'$ it is sufficient to show that $(M_1 \otimes M_2)'$ and $(M_1' \otimes M_2')'$ commute since obviously $M_1' \otimes M_2' \subseteq (M_1 \otimes M_2)'$.

If $\omega_1 \in \mathcal{H}_1$ and $\omega_2 \in \mathcal{H}_2$ we define $e_1 = [M_1' \omega]$, $e_1' = [M_1 \omega]$ and similarly $e_2 = [M_2' \omega]$, $e_2' = [M_2 \omega]$ where $[M\omega]$ denotes the projection onto the closed subspace generated by $x\omega$ with $x \in M$. It is well known that $e_1 \in M_1$, $e_1' \in M_1'$ and that the von Neumann algebras $e_1 M_1 e_1$ and $e_1' M_1' e_1'$ as acting on $e_1 e_1' \mathcal{H}$ are commutants of each other and have ω_1 as a separating and cyclic vector. A similar statement is true for M_2 .

Denote

$$\mathcal{K} = \mathcal{H}_1 \otimes \mathcal{H}_2, \quad M = M_1 \otimes M_2, \quad N = M_1' \otimes M_2', \\ e = e_1 \otimes e_2 \quad \text{and} \quad e' = e_1' \otimes e_2'.$$

From the commutation theorem for von Neumann algebras with a separating

Received by the editors March 9, 1975.

AMS (MOS) subject classifications (1970). Primary 46L10.

Copyright © 1977, American Mathematical Society

and cyclic vector applied to $e_1 M_1 e_1 e'_1$ and $e_2 M_2 e_2 e'_2$ we find that $e M e e'$ and $e' N e' e$ are commutants of each other on $ee'\mathcal{H}$. Now since $e \in M$ and $e' \in N \subseteq M'$ the commutant of $e M e e'$ on $ee'\mathcal{H}$ is $e' M' e' e$. Similarly the commutant of $e' N e' e$ on $ee'\mathcal{H}$ is $e N' e e'$. Hence $e' M' e' e$ and $e N' e e'$ commute.

Define $\omega = \omega_1 \otimes \omega_2$ and let $m' \in M'$ and $n' \in N'$. Using that $ee'\omega = \omega$ and that $e'm'e'e$ and $en'ee'$ commute, we obtain

$$\begin{aligned} \langle m'n'\omega, \omega \rangle &= \langle ee'm'n'ee'\omega, \omega \rangle = \langle (e'm'e)(e'n'e)\omega, \omega \rangle \\ &= \langle (e'm'e'e)(en'ee')\omega, \omega \rangle = \langle (en'ee')(e'm'e'e)\omega, \omega \rangle \\ &= \langle n'm'\omega, \omega \rangle. \end{aligned}$$

Using the polarization identity one can now obtain $m'n' = n'm'$. To see this denote $a = m'n' - n'm'$ so that $\langle a\omega_1 \otimes \omega_2, \omega_1 \otimes \omega_2 \rangle = 0$ for all $\omega_1 \in \mathcal{H}_1$ and $\omega_2 \in \mathcal{H}_2$. For a fixed ω_2 consider the sesquilinear form $\phi(\xi_1, \eta_1) = \langle a\xi_1 \otimes \omega_2, \eta_1 \otimes \omega_2 \rangle$ on $\mathcal{H}_1$. As $\phi(\omega_1, \omega_1) = 0$ for all $\omega_1 \in \mathcal{H}_1$ we get by polarization that $\phi = 0$, that means $\langle a\xi_1 \otimes \omega_2, \eta_1 \otimes \omega_2 \rangle = 0$ for all $\xi_1, \eta_1 \in \mathcal{H}_1$ and $\omega_2 \in \mathcal{H}_2$. Similarly for fixed $\xi_1, \eta_1 \in \mathcal{H}_1$ using the sesquilinear form $\psi(\xi_2, \eta_2) = \langle a\xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2 \rangle$ on $\mathcal{H}_2$ we get that $\langle a\xi_1 \otimes \xi_2, \eta_1 \otimes \eta_2 \rangle = 0$. But then $a = 0$. So $m'n' = n'm'$ for all $m' \in M'$ and $n' \in N'$. This completes the proof.

In [1] a simple proof of the commutation theorem is obtained already for the case of von Neumann algebras with a cyclic vector. Then of course the above simplification becomes even greater since it is sufficient to include only the projections e_1, e_2 and e and not e'_1, e'_2 and e' .

REFERENCES

1. M. A. Rieffel and A. Van Daele, *The commutation theorem for tensor products of von Neumann algebras*, Bull. London Math. Soc. 7 (1975), 257–260.
2. S. Sakai, *C*-algebras and W*-algebras*, Springer-Verlag, Berlin, 1971.
3. M. Takesaki, *Tomita's theory of modular Hilbert algebras and its applications*, Lecture Notes in Math., vol. 128, Springer-Verlag, Berlin and New York, 1970.

KATHOLIEKE UNIVERSITEIT TE LEUVEN, DEPARTEMENT WISKUNDE, CELESTIJNENLAAN, 200B, 3030 HEVERLEE, BELGIUM