A SUFFICIENT CONDITION FOR HYPERINVARIANCE

W. E. LONGSTAFF

ABSTRACT. A linear transformation on a finite-dimensional complex linear space has the property that all of its invariant subspaces are hyperinvariant if and only if its lattice of invariant subspaces is distributive [1]. It is shown that an operator on a complex Hilbert space has this property if its lattice of invariant subspaces satisfies a certain distributivity condition.

1. Preliminaries. Throughout this paper H will denote an arbitrary complex Hilbert space. All operators are bounded and all subspaces are closed. By a *subspace lattice* on H is meant a family of subspaces of H which is closed under the formation of arbitrary intersections and arbitrary closed linear spans and which contains the zero subspace (0) and H . The family of subspaces invariant under an operator T is denoted by $\text{Lat } T$. This is a subspace lattice as is the family of subspaces invariant under every operator commuting with T which we denote by $\text{Hyperlat } T$. The elements of $\text{Hyperlat } T$ are called the *hyperinvariant* subspaces of T . Clearly $\text{Hyperlat } T \subseteq \text{Lat } T$. A subspace lattice $\mathfrak{F}$ is called *commutative* if for every pair of subspaces $M, N \in \mathfrak{F}$ the corresponding projections P_M and P_N commute. Let L be an abstract lattice. We say that L is

(i) *distributive* if

$$a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c) \quad (a, b, c \in L)$$

and its dual statement holds identically;

(ii) *σ -infinitely meet distributive* if L is σ -complete and

$$a \vee \left\{ \bigwedge b_n : n \geq 1 \right\} = \bigwedge \{ a \vee b_n : n \geq 1 \} \quad (a, b_n \in L)$$

holds identically in L .

That the dual equation defining distributivity are equivalent to each other is an elementary result of lattice theory.

2. A sufficient condition for hyperinvariance. The key to the sufficient condition is a result of Sarason and the following lattice-theoretic result.

PROPOSITION 2.1. *If L is an abstract σ -infinitely meet distributive lattice and $\theta: L \rightarrow L$ is a lattice automorphism with the properties*

(I) $a \leq \theta(a) \vee \theta^{-1}(a)$ ($a \in L$);

(II) $a, \theta(a)$ comparable implies $a = \theta(a)$,

then θ is the identity automorphism.

PROOF. For every $n \geq 1$ let $\theta^n: L \rightarrow L$ be defined in the obvious way. Let a

Received by the editors March 27, 1975.

AMS (MOS) subject classifications (1970). Primary 47A15; Secondary 26A30.

Copyright © 1977, American Mathematical Society

be an arbitrary fixed element of L . For $n \geq 1$ put $a_n = a \wedge \theta(a) \wedge \theta^2(a) \wedge \cdots \wedge \theta^n(a)$. Then $a_n \wedge \theta(a_n) = a_{n+1}$ ($n \geq 1$). Using (I) and the fact that L is distributive the statement

$$\theta(x) = (x \wedge \theta(x)) \vee [\theta(x \wedge \theta(x))]$$

holds identically in L . Using this identity it is easily shown that $\theta(a) = a_1 \vee \theta(a_n)$ ($n \geq 1$). Thus

$$\theta(a) = \bigwedge \{a_1 \vee \theta(a_n) : n \geq 1\} = a_1 \vee (\bigwedge \{\theta(a_n) : n \geq 1\}).$$

If $c = \bigwedge \{a_n : n \geq 1\}$ then

$$\theta(c) = \bigwedge \{\theta(a_n) : n \geq 1\} \geq \bigwedge \{a_{n+1} : n \geq 1\} \geq c$$

and by (II), $\theta(c) = c$. Hence $\theta(a) = a_1 \vee \theta(c) = a_1 \vee c = a_1 \leq a$ and again by (II), $\theta(a) = a$. This completes the proof.

Let T be an operator on H . Notice that if S is an invertible operator commuting with T then $SM \in \text{Lat } T$ ($M \in \text{Lat } T$) and the mapping $M \rightarrow SM$ is a lattice automorphism with the mapping $M \rightarrow S^{-1}M$ as its inverse. If the operator A commutes with T and μ is a scalar with $|\mu| > \|A\|$, the operator $S = A - \mu$ is invertible, commutes with T and $\text{Lat } A = \text{Lat } S$. By a result of Sarason [4], $\text{Lat } S = \text{Lat } S^{-1}$. It readily follows that $\text{Hyperlat } T = \text{Lat } T$ if and only if for every invertible operator S commuting with T satisfying $\text{Lat } S = \text{Lat } S^{-1}$ the mapping $M \rightarrow SM$ ($M \in \text{Lat } T$) is the identity automorphism.

PROPOSITION 2.2. *If $\text{Lat } T$ is distributive and S is an invertible operator commuting with T , then $M \subseteq SM \vee S^{-1}M$ ($M \in \text{Lat } T$).*

PROOF. Choose λ with $0 < \lambda < 1/\|S\|$. The operator $C = 1 + \lambda S$ is invertible and commutes with T . Let $M \in \text{Lat } T$. It is readily verified that $CM \cap SM = C(M \cap SM)$ and $CM \cap M = C(M \cap S^{-1}M)$. Since $CM \subseteq SM \vee M$, by distributivity we have

$$\begin{aligned} CM &= (CM \cap SM) \vee (CM \cap M) = C(M \cap SM) \vee C(M \cap S^{-1}M) \\ &= C(M \cap [SM \vee S^{-1}M]) \end{aligned}$$

and the result follows.

THEOREM 2.3. *If $\text{Lat } T$ is σ -infinitely meet distributive $\text{Hyperlat } T = \text{Lat } T$.*

PROOF. By our earlier remarks it suffices to show that if S is an invertible operator commuting with T and satisfying $\text{Lat } S = \text{Lat } S^{-1}$ then the automorphism $M \rightarrow SM$ of $\text{Lat } T$ is the identity automorphism. Since $\text{Lat } T$ is distributive, this automorphism satisfies condition (I) of Proposition 2.1 by Proposition 2.2. Since $\text{Lat } S = \text{Lat } S^{-1}$, condition (II) is also satisfied. The result now follows from Proposition 2.1.

COROLLARY 2.3.1. *$\text{Hyperlat } T = \text{Lat } T$ if $\text{Lat } T$ is any one of the following:*

- (i) commutative;
- (ii) isomorphic to the direct product of complete chains;
- (iii) totally ordered.

PROOF. It is clear that in cases (ii) and (iii) $\text{Lat } T$ is σ -infinitely meet distributive. Suppose that $\text{Lat } T$ is commutative. Then it is also distributive. This follows from the fact that if P and Q are commuting projections then PQ is the projection onto the intersection of the ranges of P and Q and $P + Q - PQ$ is the projection onto the closed linear span of the ranges of P and Q . Thus if $K, L, M \in \text{Lat } T$ then

$$\begin{aligned} P_{K \cap (L \vee M)} &= P_K P_{L \vee M} = P_K (P_L + P_M - P_L P_M) \\ &= P_{K \cap L} + P_{K \cap M} - P_{K \cap L} P_{K \cap M} = P_{(K \cap L) \vee (K \cap M)}. \end{aligned}$$

Now let

$$K, L_n (n \geq 1) \in \text{Lat } T.$$

Then $P_{\cap \{L_n: n \geq 1\}}$, respectively $P_{\cap \{K \vee L_n: n \geq 1\}}$, is the strong limit of the sequence $\{P_{\cap \{L_n: 1 \leq n \leq k\}}: k \geq 1\}$, respectively $\{P_{\cap \{K \vee L_n: 1 \leq n \leq k\}}: k \geq 1\}$. But

$$\begin{aligned} P_{\cap \{K \vee L_n: 1 \leq n \leq k\}} &= P_{K \vee \cap \{L_n: 1 \leq n \leq k\}} \\ &= P_K + P_{\cap \{L_n: 1 \leq n \leq k\}} - P_K P_{\cap \{L_n: 1 \leq n \leq k\}}. \end{aligned}$$

Taking strong limits gives

$$P_{\cap \{K \vee L_n: n \geq 1\}} = P_K + P_{\cap \{L_n: n \geq 1\}} - P_K P_{\cap \{L_n: n \geq 1\}} = P_{K \vee \cap \{L_n: n \geq 1\}}.$$

Hence $\text{Lat } T$ is σ -infinitely meet distributive. The result follows by applying Theorem 2.3.

The results (i) and (iii) above are not new. The former was proved in [2] and the latter in [3].

3. Concluding remarks. It is a simple exercise to show for any linear transformation T on a finite-dimensional complex linear space that $\text{Hyperlat } T$ is distributive and finite, therefore σ -infinitely meet distributive. It seems an interesting question whether $\text{Hyperlat } T$ is always σ -infinitely meet distributive or even whether the converse of Theorem 2.3 holds. For a normal operator T , $\text{Hyperlat } T$ consists of the ranges of the spectral projections for T [2] and so is both commutative and a Boolean algebra and so is certainly σ -infinitely meet distributive.

REFERENCES

1. L. Brickman and P. A. Fillmore, *The invariant subspace lattice of a linear transformation*, *Canad. J. Math.* **19**(1967), 810–822. MR **35** #4242.
2. R. G. Douglas and Carl Pearcy, *On a topology for invariant subspaces*, *J. Functional Analysis* **2**(1968), 323–341. MR **38** #1547.
3. Peter Rosenthal, *A note on unicellular operators*, *Proc. Amer. Math. Soc.* **19**(1968), 505–506. MR **36** #5753.
4. Donald Sarason, *The H^p spaces of an annulus*, *Mem. Amer. Math. Soc. No. 56*(1965). MR **32** #6256.

SCHOOL OF MATHEMATICAL SCIENCES, FLINDERS UNIVERSITY, BEDFORD PARK, SOUTH AUSTRALIA 5042