DEPENDENCE ON DIMENSION OF A CONSTANT RELATED TO THE GRÖTZSCH RING

GLEN D. ANDERSON

ABSTRACT. For the constant $\lambda_n = \lim_{a \rightarrow 0} (\text{mod } R_{G,n}(a) + \log a)$ associated with the Grötzsch extremal ring $R_{G,n}$ in euclidean n -space, we obtain the limit $\lim_{n \rightarrow \infty} \lambda_n^{1/n} = e$.

1. Definitions and notation. By a *ring* R is meant a domain in euclidean n -space R^n whose complement consists of two components C_0 and C_1 , where C_0 is bounded. We let $B_0 = \partial C_0$ and $B_1 = \partial C_1$ be the boundary components of R . The *conformal capacity* (cf. [8]) of R is

$$\text{cap } R = \inf_{\varphi} \int_R |\nabla \varphi|^n d\omega,$$

where ∇ denotes the gradient, and where the infimum is taken over all real-valued C^1 functions φ in R with boundary values 0 on B_0 and 1 on B_1 . Then the *modulus* of the ring R is defined by

$$\text{mod } R = (\sigma_{n-1} / \text{cap } R)^{1/(n-1)},$$

where for each positive integer p we let σ_p denote the p -dimensional measure of the unit sphere $S^p = \{(x_1, \dots, x_{p+1}): \sum_{j=1}^{p+1} x_j^2 = 1\}$. Then

$$\sigma_p = 2\pi^{(p+1)/2} \Gamma((p+1)/2)^{-1}$$

(cf. [7], [9]), where Γ denotes the classical Gamma function; moreover, the relation

$$(1) \quad \int_0^{\pi/2} \cos^p u \, du = \frac{\sigma_{p+1}}{2\sigma_p}$$

holds for each positive integer p .

2. Background of problem and statement of result. Let $R_{G,n}(a)$ denote the n -dimensional Grötzsch ring, that is, the ring whose complementary components are

$$C_0 = \{(x_1, \dots, x_n): 0 \leq x_1 \leq a, x_j = 0, 2 \leq j \leq n\}$$

and

Received by the editors June 2, 1975.

AMS (MOS) subject classifications (1970). Primary 30A60.

Copyright © 1977, American Mathematical Society

$$C_1 = \left\{ (x_1, \dots, x_n) : \sum_{j=1}^n x_j^2 \geq 1 \right\}.$$

In [5] Gehring proved that $\text{mod } R_{G,3}(a) + \log a$ is monotone decreasing in the interval $0 < a < 1$ and he obtained bounds $4 \leq \lambda_3 \leq 12.4 \dots$, where $\log \lambda_3 = \lim_{a \rightarrow 0} (\text{mod } R_{G,3}(a) + \log a)$. Using analogous methods in higher dimensions, Caraman [3] and Ikoma [7] have shown that the limit

$$\log \lambda_n = \lim_{a \rightarrow 0} (\text{mod } R_{G,n}(a) + \log a)$$

exists for each $n \geq 3$. Bounds for λ_n have been obtained by Gehring [5], [6], Caraman [3], Ikoma [7], and Anderson [2] (cf. [1]).

In the present paper we determine the order of growth of λ_n for large n by establishing the following result.

THEOREM. $\lim_{n \rightarrow \infty} \lambda_n^{1/n} = e$, where e is the base of Napierian logarithms.

3. Proof of Theorem. First, by the work of Caraman [3] and Ikoma [7] we know that

$$(2) \quad \log \frac{\lambda_n}{4} \leq I_n = \int_1^\infty \left[\left(\frac{r^2 + 1}{r^2 - 1} \right)^{(n-2)/(n-1)} - 1 \right] \frac{dr}{r}.$$

Under the change of variable $t = (r^2 + 1)/(r^2 - 1)$ the integral I_n in (2) reduces to

$$(3) \quad I_n = \int_1^\infty \frac{t^{(n-2)/(n-1)} - 1}{t^2 - 1} dt.$$

Making use of the fact that $(t^2 - 1)^{-1} = t^{-2} + t^{-2}(t^2 - 1)^{-1}$, one may rewrite I_n as the sum of two integrals. The first of these is easy to evaluate exactly as $n - 2$, and by means of the Monotone Convergence Theorem it is easy to see that the second increases to the limit $1 - \log 2$. This procedure, together with (2), yields the result

$$(4) \quad \log \frac{\lambda_n}{4} - n \leq \lim_{n \rightarrow \infty} (I_n - n) = -1 - \log 2,$$

from which the upper bound $\lambda_n^{1/n} \leq e$ is easily derived (cf. [4, Lecture 9, Theorem 1]).

On the other hand, it was determined in [2] that

$$(5) \quad \log \frac{\lambda_n}{4} \geq \int_0^\infty \left[\left(\frac{2\sigma_{n-2}}{\sigma_{n-1}} \int_0^{\pi/2} (\sec^2 u + \text{csch}^2 v)^{(2-n)/2} du \right)^{1/(1-n)} - 1 \right] dv.$$

By making the change of variable $t = \coth v$ in (5) and combining the resulting inequality with (1), (2), and (3), we achieve

$$(6) \quad 0 \geq \log \frac{\lambda_n}{4} - I_n \geq \int_1^\infty \Phi_n(t) dt,$$

where

$$(7) \quad \Phi_n(t) = \frac{t^{(n-2)/(n-1)}}{t^2 - 1} \left[\left(\frac{2\sigma_{n-2}}{\sigma_{n-1}} \int_0^{\pi/2} \left(1 + \frac{\tan^2 u}{t^2} \right)^{(2-n)/2} du \right)^{1/(1-n)} - 1 \right].$$

We consider first the integral of $\Phi_n(t)$ over the finite interval $(1, M)$. First, using (1), it is convenient to rewrite (7) in the form

$$(8) \quad \Phi_n(t) = \frac{t^{(n-2)/(n-1)}}{t^2 - 1} \left[\left(\frac{\|\cos u\|_{n-2}}{\|(1 + t^{-2} \tan^2 u)^{-1/2}\|_{n-2}} \right)^{(n-2)/(n-1)} - 1 \right],$$

where $\|f\|_p$ denotes the $L_p[0, \pi/2]$ -norm of f . Since the functions $\cos u$ and $(1 + t^{-2} \tan^2 u)^{-1/2}$ are continuous on $[0, \pi/2]$, a well-known result in the theory of L_p spaces gives

$$(9) \quad \lim_{n \rightarrow \infty} \frac{\|\cos u\|_{n-2}}{\|(1 + t^{-2} \tan^2 u)^{-1/2}\|_{n-2}} = \frac{\max_{0 \leq u \leq \pi/2} |\cos u|}{\max_{0 \leq u \leq \pi/2} |1 + t^{-2} \tan^2 u|^{-1/2}} = 1.$$

From (8) and (9) it follows that for each fixed $t \in (1, M)$,

$$(10) \quad \lim_{n \rightarrow \infty} \Phi_n(t) = 0.$$

Next, in view of (8) it is easy to see that

$$0 > \Phi_n(t) \geq t^{(n-2)/(n-1)} (t^{(2-n)/(n-1)} - 1) / (t^2 - 1) \geq -1/(1 + t).$$

This and (10) allow us to invoke the Dominated Convergence Theorem to conclude that

$$(11) \quad \lim_{n \rightarrow \infty} \int_1^M \Phi_n(t) dt = 0$$

for each fixed $M \in (1, \infty)$.

Next, using the fact that $t^2/(t^2 - 1)$ is decreasing for $t > 1$ and that $\Phi_n(t) \leq 0$, we may write

$$\int_M^\infty \Phi_n(t) dt \geq \frac{M^2}{M^2 - 1} \left[\left(\frac{\|\cos u\|_{n-2}}{\|1\|_{n-2}} \right)^{(n-2)/(n-1)} - 1 \right] \int_M^\infty t^{(n-2)/(n-1)-2} dt.$$

Thus

$$\frac{1}{n} \int_M^\infty \Phi_n(t) dt \geq \frac{M^2}{M^2 - 1} \left(1 - \frac{1}{n} \right) M^{1/(1-n)} \left[\left(\frac{2}{\pi} \right)^{1/(n-1)} \|\cos u\|_{n-2}^{(n-2)/(n-1)} - 1 \right],$$

and the same argument as that leading to (10) shows that

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \int_M^\infty \Phi_n(t) dt \geq 0.$$

But because of (11), (6), and (4), this implies the statement

$$(12) \quad \liminf_{n \rightarrow \infty} \frac{1}{n} \log \lambda_n \geq 1.$$

Finally, the Theorem follows from (2) and (12).

REFERENCES

1. G. D. Anderson, *Symmetrization and extremal rings in space*, Ann. Acad. Sci. Fenn. Ser. AI No. 438 (1969), 1–24. MR 41 #465.
2. ———, *Extremal rings in n -space for fixed and varying n* , Ann. Acad. Sci. Fenn. Ser. AI No. 575 (1974), 1–21.
3. P. Caraman, *On the equivalence of the definitions of the n -dimensional quasiconformal homeomorphisms (QCfH)*, Rev. Roumaine Math. Pures Appl. 12 (1967), 889–943. MR 37 #409.
4. F. W. Gehring, *Lectures on quasiconformal mappings*, Institut Mittag-Leffler, 1972 (unpublished).
5. ———, *Symmetrization of rings in space*, Trans. Amer. Math. Soc. 101 (1961), 499–519. MR 24 #A2677.
6. ———, *Inequalities for condensers, hyperbolic capacity, and extremal lengths*, Michigan Math. J. 18 (1971), 1–20. MR 44 #2915.
7. K. Ikoma, *An estimate for the modulus of the Grötzsch ring in n -space*, Bull. Yamagata Univ. Natur. Sci. 6 (1967), no. 4, 395–400. MR 43 #508.
8. C. Loewner, *On the conformal capacity in space*, J. Math. Mech. 8 (1959), 411–414. MR 21 #3538.
9. J. Väisälä, *Lectures on n -dimensional quasiconformal mappings*, Lecture Notes in Math., vol. 229, Springer-Verlag, Berlin, 1971.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824