ON CERTAIN EXTREMAL PROBLEMS FOR FUNCTIONS WITH POSITIVE REAL PART

STEPHAN RUSCHEWEYH AND VIKRAMADITYA SINGH

ABSTRACT. For the class P of analytic functions $p(z)$, $p(0) = 1$, with positive real part in $|z| < 1$, a type of extremal problems is determined which can be solved already within the set $p(z) = (1 + \varepsilon z)/(1 - \varepsilon z)$, $|\varepsilon| = 1$. One problem of this kind is to find the largest number $\rho(s, \mu)$ such that

$$\operatorname{Re}\{p(z) + szp'(z)/(p(z) + \mu)\} > 0,$$

$|z| < \rho(s, \mu)$, for all $p \in P$, $-1 \neq \mu \in \mathbb{C}$, $s > 0$. Sharp upper bounds for two other functionals over P are also given.

1. Introduction. Let P be the class of analytic functions $p(z)$ in the unit disc $\Delta = \{z \mid |z| < 1\}$ with $p(0) = 1$, $\operatorname{Re} p(z) > 0$ in Δ . Robertson [2], [3] and Sakaguchi [6] obtained variational formulae for P and showed that, for fixed $z \in \Delta$, the extreme values of

$$(1) \quad \operatorname{Re} F(p(z), zp'(z)), \quad p \in P,$$

where $F(u, v)$ is analytic in $(u, v) \in \mathbb{C}^2$, $\operatorname{Re} u > 0$, are already attained by the functions

$$(2) \quad \lambda \left(\frac{1 + \varepsilon z}{1 - \varepsilon z} \right) + (1 - \lambda) \left(\frac{1 + \bar{\varepsilon} z}{1 - \bar{\varepsilon} z} \right), \quad \lambda \in [0, 1], |\varepsilon| = 1.$$

Zmorovič [10], [11] and others used this result to solve extremal problems in P . In many cases of interest, however, even this reduced problem is difficult to solve or appears insurmountable.

In the first part of this paper we present a theorem which characterises a class of extremal problems in P which can be solved in

$$P_0 := \{p \in P \mid p(z) = (1 + \varepsilon z)/(1 - \varepsilon z), |\varepsilon| = 1\}.$$

Our theorem rests on recent results of Suffridge [8] concerning the Hadamard product of starlike functions of order α (compare also Ruscheweyh and Sheil-Small [5] and Ruscheweyh [4]).

One problem which can be treated by our method is to find the largest number $\rho(s, \mu)$ such that for all $p \in P$

$$\operatorname{Re}\{p(z) + szp'(z)/(p(z) + \mu)\} > 0, \quad |z| < \rho(s, \mu),$$

holds, where $-1 \neq \mu \in \mathbb{C}$, $s > 0$. We shall use this result to solve a problem posed by Yoshikawa [9] and dealing with the 'radius of spiral-convexity' of spirallike functions in Δ .

In the last section we shall derive the exact upper bounds of the functionals

Received by the editors October 14, 1975.

AMS (MOS) subject classifications (1970). Primary 30A76, 30A40; Secondary 30A38, 30A32.

Key words and phrases. Functions with positive real part, Hadamard product, spiral-convexity.

© American Mathematical Society 1977

$(\operatorname{Re} p(z))/|p(z) + i \tan \alpha|$ and $|zp'(z)/(p(z) + i \tan \alpha)|$ for $p \in P$. Crude bounds, in particular of the second entity, have been used by several authors [1], [3], [7], but a sharp result has not so far been obtained.

2. Let

$$f(z) = \sum_{k=0}^{\infty} a_k z^k, \quad g(z) = \sum_{k=0}^{\infty} b_k z^k$$

be analytic in Δ . The function

$$(f * g)(z) := \sum_{k=0}^{\infty} a_k b_k z^k$$

is also analytic in Δ and is called the Hadamard product of f and g .

THEOREM 1. For $p \in P$ let

$$\hat{p}(z) := \exp \left(\int_0^z \frac{p(\xi) - 1}{\xi} d\xi \right).$$

Let $F(z)$ be analytic in Δ , $F(0) \neq 0$, and let $t \geq 0$. Then we have for fixed $z_0 \in \Delta$:

$$(3) \quad \min_{p \in P} \operatorname{Re} \frac{(F * (p\hat{p}'))(z_0)}{(F * \hat{p}')(z_0)} = \min_{p \in P_0} \operatorname{Re} \frac{(F * (p\hat{p}'))(z_0)}{(F * \hat{p}')(z_0)},$$

whenever the right-hand side of (3) is nonnegative.

It should be pointed out that

$$t \frac{(F * (p\hat{p}'))(z)}{(F * \hat{p}')(z)} = \frac{zg'(z)}{g(z)} + t - 1$$

where $g(z) = z(F * \hat{p}')(z)$.

Theorem 1 is almost trivial for $t = 0$. The case $t = 1$, however, contains the truth of the Pólya-Schoenberg conjecture for convex univalent functions in Δ .

REMARK 1. It is easy to see that the right-hand side of (3) is nonnegative if and only if

$$(4) \quad F * (1 - \varepsilon z)/(1 - z)^{1+2t} \neq 0, \quad |z| < |z_0|, \quad |\varepsilon| \leq 1.$$

This relation is useful for applications of Theorem 1.

Let $\alpha < 1$. By S_α^* we denote the class of functions starlike of order α in Δ , i.e. $f \in S_\alpha^*$ if and only if f is analytic in Δ , $f(0) = f'(0) - 1 = 0$, and fulfils $\operatorname{Re}(zf'(z)/f(z)) > \alpha$, $z \in \Delta$. In particular we have

$$s_\alpha := z/(1 - z)^{2-2\alpha} \in S_\alpha^*.$$

By s_α^{-1} we denote that well-defined analytic function which satisfies the relations

$$s_\alpha^{-1} * s_\alpha = z/(1 - z), \quad s_\alpha^{-1}(0) = 0.$$

We shall require the following lemmata.

LEMMA 1 (SUFFRIDGE [7, THEOREM 7]). For $\beta < 1$ we have $f \in S_\beta^*$ if and only if $s_\beta^{-1} * f * g \in S_\beta^*$ for all $g \in S_\beta^*$.

LEMMA 2 (SUFFRIDGE [7, THEOREM 10]). Let $\alpha \leq \beta < 1$. Then $f \in S_\alpha^*$ implies $s_\alpha^{-1} * s_\beta * f \in S_\beta^*$.

For $\alpha \leq \beta < 1$ we introduce the set $V_{\alpha\beta}$ of analytic functions $h(z)$ in Δ , $h(0) = 0$, such that $h * f \in S_\beta^*$ for each $f \in S_\alpha^*$.

LEMMA 3. Let $h(z)$ be analytic in Δ , $h(0) = 0$. Then $h \in V_{\alpha\beta}$ if and only if $h * s_\alpha \in S_\beta^*$.

PROOF. Let $g \in S_\beta^*$, $h := s_\alpha^{-1} * g$. Then for each $f \in S_\alpha^*$,

$$f * h = (s_\alpha^{-1} * s_\beta * f) * s_\beta^{-1} * g \in S_\beta^*$$

by Lemma 1, since $s_\alpha^{-1} * s_\beta * f \in S_\beta^*$ by Lemma 2. On the other hand, $h \in V_{\alpha\beta}$ implies $h * s_\alpha \in S_\beta^*$ since $s_\alpha \in S_\alpha^*$.

PROOF OF THEOREM 1. Lemma 3 shows that for $\alpha \leq \beta < 1$ and a certain $h(z)$ analytic in Δ , $h(0) = 0$, we have

$$(5) \quad \operatorname{Re} \frac{z(h * g)'(z)}{(h * g)(z)} > \beta \quad \text{for all } g \in S_\alpha^*, z \in \Delta,$$

if and only if

$$(6) \quad \operatorname{Re} \frac{z(h * s_\alpha)'(z)}{(h * s_\alpha)(z)} > \beta, \quad z \in \Delta.$$

Now let $p \in P$, $t > 0$. Obviously $z\hat{p}'(z) \in S_{1-t}^*$ and, in particular, for $p_0(z) = (1+z)/(1-z)$ we have $z\hat{p}'_0(z) = s_{1-t}(z)$. In this notation, (5) and (6) read ($1-t \leq \beta < 1$):

$$(7) \quad \operatorname{Re} t \frac{(h * (zp\hat{p}'))(z)}{(h * z\hat{p}')(z)} > \beta + t - 1 \quad \text{for all } p \in P, z \in \Delta,$$

if and only if

$$(8) \quad \operatorname{Re} t \frac{(h * (zp_0\hat{p}'_0))(z)}{(h * z\hat{p}'_0)(z)} > \beta + t - 1, \quad z \in \Delta.$$

Now assume

$$(9) \quad \min_{p \in P_0} \operatorname{Re} \frac{(F * p\hat{p}')(z_0)}{(F * \hat{p}')(z_0)} = v \geq 0$$

and put $h(z) := zF(z_0z)$, $v = (\beta + t - 1)/t$. Thus (9) implies (8) and, hence, (7), which in turn may be written as

$$(10) \quad \min_{p \in P} \operatorname{Re} \frac{(F * p\hat{p}')(z_0)}{(F * \hat{p}')(z_0)} = v.$$

The case $t = 0$ is well known and follows from the Herglotz representation of the functions in P .

3. In this section we give a simple application of Theorem 1. For $-1 \neq \mu \in \mathbb{C}$, $s > 0$, let

$$(11) \quad F(z) := \left(\frac{\mu + 1}{s} - 1 \right) \frac{1}{1-z} + \frac{1}{(1-z)^2}.$$

Then for $p \in P$ we obtain

$$\frac{(F * (p\hat{p}^{1/s}))(z)}{(F * \hat{p}^{1/s})(z)} = p(z) + s \frac{zp'(z)}{p(z) + \mu}.$$

COROLLARY 1. For $p \in P$ and

$$(12) \quad |z| < \frac{|\mu + 1|}{\left[A + (A^2 - |\mu^2 - 1|^2)^{1/2} \right]^{1/2}}, \quad A = 2(s+1)^2 + |\mu|^2 - 1,$$

we have

$$(13) \quad \operatorname{Re} \left(p(z) + s \frac{zp'(z)}{p(z) + \mu} \right) > 0.$$

The bound given by (12) is best possible.

PROOF. From Remark 1 we deduce that we only have to find the largest radius $\rho = \rho(s, \mu)$ such that for F , given by (11),

$$F * \frac{1 - \varepsilon z}{(1-z)^{1+2/s}} \neq 0, \quad |z| < \rho(s, \mu), \quad |\varepsilon| \leq 1.$$

This is equivalent to the condition

$$-s \frac{\varepsilon z}{1 - \varepsilon z} + (s+2) \frac{z}{1-z} \neq -\mu - 1, \quad |z| < \rho(s, \mu), \quad |\varepsilon| \leq 1,$$

and thus with $|2\rho^2 + (\mu + 1)(1 - \rho^2)| \geq 2\rho(s+1)$. This leads immediately to (12).

It should be mentioned that one can use Theorem 1 to compute

$$m(r) = \min_{p \in P} \min_{|z| \leq r} \operatorname{Re} \left(p(z) + \frac{szp'(z)}{p(z) + \mu} \right)$$

as long as $m(r)$ is nonnegative.

Let $f(z)$ be analytic in Δ , $f(0) = f'(0) - 1 = 0$. Yoshikawa [9] called $f(z)$ “ γ -spiral-convex” if it satisfies the condition

$$(14) \quad \operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} + i \tan \gamma \frac{zf'(z)}{f(z)} \right) > 0$$

in Δ and gave a geometrical interpretation of this condition. He proved that every γ -spiral-convex function is γ -spirallike, i.e. it fulfils

$$(15) \quad \operatorname{Re} e^{i\gamma} \frac{zf'(z)}{f(z)} > 0, \quad z \in \Delta.$$

He posed the problem to find the largest radius $\rho = \rho(\gamma)$, such that every γ -spirallike function is γ -spiral-convex in $|z| < \rho(\gamma)$, thereby generalizing the

radius of (0-spiral-)convexity of the starlike (= 0-spirallike) functions in Δ .

It is easily seen that this problem is equivalent to the following question: What is the greatest number $\rho(\gamma)$ such that

$$\operatorname{Re} \left(p(z) + \frac{zp'(z)}{p(z) + i \tan \gamma} \right) > 0, \quad |z| < \rho(\gamma),$$

for all $p \in P$. This question has been answered in Corollary 1 and thus we can state

THEOREM 2. *The greatest common radius $\rho(\gamma)$ of γ -spiral-convexity of γ -spirallike functions in Δ is given by*

$$(16) \quad \rho(\gamma) = \sqrt{1 + 3 \cos^2 \gamma} - \sqrt{3} \cos \gamma.$$

Yoshikawa himself was able to show that the value given in (16) is an upper bound for $\rho(\gamma)$.

4. In this last section we derive upper bounds of two special functionals on P . The method used is completely apart from those discussed in the previous chapters. The extremal functions, however, belong to P_0 too.

THEOREM 3. *For $p \in P$ and $\alpha \in (-\pi/2, \pi/2)$ the estimate*

$$(17) \quad \frac{\operatorname{Re} p(z)}{|p(z) + i \tan \alpha|} \leq \begin{cases} \frac{(1 - |z|^2) \cos \alpha}{1 - 2|z| |\sin \alpha| + |z|^2}, & |z| < \left| \tan \frac{\alpha}{2} \right|, \\ 1, & |z| \geq \left| \tan \frac{\alpha}{2} \right|, \end{cases}$$

is valid and sharp. Equality holds for certain functions in P_0 .

The estimate $|p'(z)| \leq (2 \operatorname{Re} p(z))/(1 - |z|^2)$ holds for every $p \in P$ and $z \in \Delta$ (see [3]). Equality holds only for every $p \in P_0$ and $z \in \Delta$. Thus Theorem 3 has the following corollary.

COROLLARY 2. *For $p \in P$ and $\alpha \in (-\pi/2, \pi/2)$ the estimate*

$$\left| \frac{zp'(z)}{p(z) + i \tan \alpha} \right| \leq \begin{cases} \frac{2|z| \cos \alpha}{1 - 2|z| |\sin \alpha| + |z|^2}, & |z| < \left| \tan \frac{\alpha}{2} \right|, \\ \frac{2|z|}{1 - |z|^2}, & |z| \geq \left| \tan \frac{\alpha}{2} \right|, \end{cases}$$

is valid and sharp. Equality holds only for certain functions $p \in P_0$.

We shall require the following well-known fact:

The range of the functional $w = p(z)$, z fixed, over P is the closed disc $K_{|z|}$ with center $(1 + |z|^2)/(1 - |z|^2)$ and radius $2|z|/(1 - |z|^2)$. Furthermore, the points on $\partial K_{|z|}$ correspond only to functions in P_0 .

$$\frac{\operatorname{Re} p(z)}{|p(z) + i \tan \alpha|} \leq \frac{\operatorname{Re} p(z)}{\operatorname{Re} p(z)} = 1,$$

and equality can hold if and only if there exists a $w_1 \in \partial K_{|z|}$ with $\operatorname{Im} w_1 = -\tan \alpha$. This is the case if and only if $|z| \geq |\tan(\alpha/2)|$ and equality holds for a certain $p \in P_0$. This proves the bottom half of (17).

Now let $z_0 \in \Delta$ be fixed, $|z_0| < |\tan(\alpha/2)|$. Let $p_0 \in P$ be an extremal

function for our functional at z_0 (P is a normal family). The function

$$p^*(z) = (p_0(z) + i\lambda) / (1 + i\lambda p_0(z))$$

belongs to P for arbitrary $\lambda \in \mathbf{R}$ and gives the variational formula

$$p^*(z) = p_0(z) + i\lambda(1 - p_0^2(z)) + o(\lambda).$$

Using the extremal property of p_0 and this variation we are led to the condition $\cos \alpha |p_0(z) + i \tan \alpha| = 1$. If we let $p_0(z_0) + i \tan \alpha = \sec \alpha e^{i\gamma}$ it follows for $p \in P$

$$\frac{\operatorname{Re} p(z_0)}{|p(z_0) + i \tan \alpha|} \leq \frac{\operatorname{Re} p_0(z_0)}{|p_0(z_0) + i \tan \alpha|} = \cos \gamma,$$

and we are left with the problem to maximize $\cos \gamma$ within the set of points $\sec \alpha e^{i\gamma} - i \tan \alpha \in K_{|z_0|}$. It is easily seen that the extremal value is

$$\frac{(1 - |z_0|^2) \cos \alpha}{1 - 2|z_0| |\sin \alpha| + |z_0|^2}$$

and corresponds to a boundary point of $K_{|z_0|}$. The theorem is proven.

ADDED IN PROOF. Theorem 2 has also been established by P. Eenigenburg and H. Yoshikawa in their paper *An application of the method of Zmorovič in geometric function theory*, J. Math. Anal. Appl. (to appear).

REFERENCES

1. S. D. Bernardi, *New distortion theorems for functions of positive real part and applications to the partial sums of univalent convex functions*, Proc. Amer. Math. Soc. **45** (1974), 113–118. MR **50** #10223.
2. M. S. Robertson, *Variational methods for functions with positive real part*, Trans. Amer. Math. Soc. **102** (1962), 82–93. MR **24** #A3288.
3. ———, *Extremal problems for analytic functions with positive real part and applications*, Trans. Amer. Math. Soc. **106** (1963), 236–253. MR **26** #325.
4. St. Ruscheweyh, *New criteria for univalent functions*, Proc. Amer. Math. Soc. **49** (1975), 109–115.
5. St. Ruscheweyh and T. Sheil-Small, *Hadamard products of schlicht functions and the Pólya-Schoenberg conjecture*, Comment. Math. Helv. **48** (1973), 119–135. MR **48** #6393.
6. K. Sakaguchi, *A variational method for functions with positive real part*, J. Math. Soc. Japan **16** (1964), 287–297. MR **31** #1375.
7. D. B. Shaffer, *Distortion theorems for a special class of analytic functions*, Notices Amer. Math. Soc. **19** (1972), A-519. Abstract #72T-B152.
8. T. J. Suffridge, *Starlike functions as limits of polynomials* (preprint).
9. H. Yoshikawa, *On a subclass of spiral-like functions*, Mem. Fac. Sci. Kyushu Univ. Ser. A **25** (1971), 271–279. MR **46** #9320.
10. V. A. Zmorovič, *On bounds of convexity for starlike functions of order α in the circle $|z| < 1$ and in the circular region $0 < |z| < 1$* , Mat. Sb. **68** (110) (1965), 518–526; English transl., Amer. Math. Soc. Transl. (2) **80** (1969), 203–213. MR **33** #5875.
11. ———, *On the bounds of starlikeness and univalence in certain classes of functions regular in the circle $|z| < 1$* , Ukrain. Mat. Ž. **18** (1966), 28–39; English transl., Amer. Math. Soc. Transl. (2) **80** (1969), 227–242. MR **33** #7525.

ABTEILUNG MATHEMATIK, UNIVERSITÄT DORTMUND, 4600 DORTMUND 50, WEST GERMANY

DEPARTMENT OF MATHEMATICS, PUNJABI UNIVERSITY, PATIALA (Pb), INDIA