THE EQUIVALENCE OF VARIOUS LIPSCHITZ CONDITIONS ON THE WEIGHTED MEAN OSCILLATION OF A FUNCTION

EDWARD P. LOTKOWSKI AND RICHARD L. WHEEDEN¹

ABSTRACT. The main result establishes the equivalence of various Lipschitz-like conditions on the weighted mean oscillation over cubes of a function. Previously known results are obtained as special cases.

Introduction. In [3], John and Nirenberg proved that if f is a function of bounded mean oscillation, i.e., if

$$\int_I |f(x) - f_I| dx \leq c|I|, \quad f_I = \frac{1}{|I|} \int_I f(t) dt,$$

for all cubes $I \subset R^n$, with c independent of I , then there exist positive constants c_1 and c_2 depending only on n such that

$$|\{x \in I: |f(x) - f_I| > \alpha\}| \leq c_1 e^{-c_2 \alpha/c} |I|$$

for all I and $\alpha > 0$. In [7] Muckenhoupt and Wheeden derived an analogous result for functions of weighted bounded mean oscillation, by which we mean those f with

$$\int_I |f(x) - f_I| dx \leq c \int_I w(x) dx$$

for all I , where w is nonnegative and satisfies appropriate conditions.

Meyers [5] and Campanato [1] showed independently that the condition

$$\int_I |f(x) - f_I| dx \leq c|I|^{1+\varepsilon}$$

for an ε satisfying $0 < \varepsilon \leq 1$ is equivalent (after redefining f in a set of measure 0) to Lipschitz continuity of order ε . Two different weighted versions of this result were then obtained by Cuerva [2] and Lotkowski [4]. Our main theorem generalizes and unifies these latter results, as we shall indicate below.

We first list some definitions. If μ is a Borel measure which is positive and

Received by the editors April 2, 1976.

AMS (MOS) subject classifications (1970). Primary 26A16.

¹ Research supported in part by NSF-MPS 75-07596.

finite on cubes and f is locally integrable with respect to μ , then $f_{I,d\mu}$ (or simply f_I) will denote the average of f over I with respect to μ :

$$f_I = f_{I,d\mu} = \frac{1}{\mu(I)} \int_I f(x) d\mu(x).$$

For $a > 0$, aI will denote the cube concentric with I whose edge length is a times that of I . We shall be concerned only with measures which satisfy the doubling condition

$$(1) \quad \mu(2I) \leq c\mu(I)$$

for every I , with c independent of I . It is an easy consequence of (1) that two cubes of equal size whose centers are at a distance bounded by a fixed multiple of their edge length have equivalent μ measure. For $1 < p < \infty$, a nonnegative function u is said to belong to $A_p(d\mu)$ if there is a constant c such that for all I

$$(2) \quad \left(\frac{1}{\mu(I)} \int_I u(x) d\mu(x) \right) \left(\frac{1}{\mu(I)} \int_I u(x)^{-1/(p-1)} d\mu(x) \right)^{p-1} \leq c.$$

Similarly, u is said to belong to $A_1(d\mu)$ if

$$(3) \quad u_\mu^*(x) \leq cu(x) \quad \text{a.e. } (d\mu),$$

with c independent of x , where u_μ^* is the Hardy-Littlewood maximal function of u with respect to $d\mu$:

$$u_\mu^*(x) = \sup \left\{ \frac{1}{\mu(I)} \int_I u(t) d\mu(t) : x \in I \right\}.$$

$F(I)$ will denote a positive function of cubes for which there exist constants α , β and c such that $1 < \alpha < \beta$, $c > 0$ and

$$(4) \quad F(I) \leq cF(J) \quad \text{if } I \subset J,$$

$$(5) \quad \alpha F(I) \leq F(2I) \leq \beta F(I).$$

It is crucial for what follows that α strictly exceed 1. An example of such an F is

$$F(I) = [\mu_1(I)]^{\epsilon_1} \cdots [\mu_m(I)]^{\epsilon_m}$$

with μ_j satisfying (1) and $\epsilon_j > 0$ for $1 \leq j \leq m$.

We shall often use the same letter c to denote different constants which are independent of x and I . For a function g which is nonzero a.e., g^{-1} will denote the function $1/g$.

Our main result is given by the following theorem.

THEOREM. *Suppose that g is nonnegative and locally integrable with respect to*

μ , that the measure $g d\mu$ satisfies (1), and that F satisfies (4) and (5). If $1 \leq p < \infty$ and $g^{-1} \in A_p(g d\mu)$, then the condition

$$(6) \quad \int_I |f(x) - f_I| g(x) d\mu(x) \leq c\mu(I)F(I), \quad f_I = f_{I, g d\mu},$$

is equivalent to

$$(7) \quad \int_I \{|f(x) - f_I| g(x)\}^{p'} d\mu(x) \leq c\mu(I)F(I)^{p'}, \quad 1/p + 1/p' = 1.$$

In case $p = 1$, we have $p' = \infty$, and (7) should be interpreted in the L^∞ sense:

$$(8) \quad \operatorname{ess\,sup}_I \{|f(x) - f_I| g(x)\} \leq cF(I),$$

the $\operatorname{ess\,sup}$ being taken with respect to $d\mu$, or equivalently $g d\mu$.

Before proceeding with the proof, we list two important special cases. First, given $\varepsilon > 0$ and a weight function w , we choose

$$\mu(I) = \int_I w(x) dx, \quad F(I) = \left[\int_I w(x) dx \right]^\varepsilon, \quad g(x) = w(x)^{-1},$$

obtaining that for $1 < p < \infty$, the condition

$$(9) \quad \int_I |f(x) - f_I| dx \leq c \left[\int_I w(x) dx \right]^{1+\varepsilon}, \quad f_I = f_{I, dx},$$

is equivalent to

$$\int_I |f(x) - f_I|^{p'} w(x)^{-1/(p-1)} dx \leq c \left[\int_I w(x) dx \right]^{1+\varepsilon p'}, \quad 1/p + 1/p' = 1,$$

provided that $w \in A_p(dx)$. If $w \in A_1(dx)$, (9) is equivalent to

$$\operatorname{ess\,sup}_I \{|f(x) - f_I| w(x)^{-1}\} \leq c \left[\int_I w(x) dx \right]^\varepsilon,$$

the $\operatorname{ess\,sup}$ being with respect to Lebesgue measure. Results of this type were proved in [2], using duality methods.

Taking μ and ν to satisfy (1), $F(I) = \nu(I)^\varepsilon$ and $g(x) \equiv 1$, one obtains that for any $p \geq 1$, the condition

$$\int_I |f(x) - f_I|^{p'} d\mu(x) \leq c\mu(I)\nu(I)^{\varepsilon p'}, \quad f_I = f_{I, d\mu}$$

is equivalent to

$$|f(x) - f_I| \leq c\nu(I)^\varepsilon \quad \text{for a.e. } (d\mu)x \in I,$$

since in this case g trivially satisfies the A_p condition for any $p \geq 1$. This may be found in [4].

Proof of the Theorem. The hard part of the proof is the implication (6) $\Rightarrow$ (7), the opposite one being an immediate corollary of Hölder's inequality. The proof we give is based on the methods used in [1] and [4]. We would like to thank Professor C. Fefferman for a useful comment concerning the proof of Lemma 1.

LEMMA 1. *Let μ and ν satisfy (1), and F satisfy (4) and (5). If*

$$(10) \quad \int_I |f(x) - f_I| d\nu(x) \leq c\mu(I)F(I), \quad f_I = f_{I, d\nu},$$

then for a.e. $(d\nu)x \in I$,

$$(11) \quad |f(x) - f_I| \leq cF(I) \sup\{(\mu(J)/\nu(J)): x \in J \subset 2I\}.$$

PROOF. For $x \in I$, let I_x denote the cube centered at x with the same edge length as I . We first claim that

$$(12) \quad |f_I - f_{I_x}| \leq cF(I) \frac{\mu(I)}{\nu(I)}.$$

In fact,

$$\begin{aligned} |f_I - f_{I_x}| &\leq |f_I - f_{2I}| + |f_{2I} - f_{I_x}| \\ &= \left| \frac{1}{\nu(I)} \int_I \{f(t) - f_{2I}\} d\nu(t) \right| + \left| \frac{1}{\nu(I_x)} \int_{I_x} \{f(t) - f_{2I}\} d\nu(t) \right|. \end{aligned}$$

Since both I and I_x lie in $2I$, and ν satisfies (1), each of the last two terms is majorized by

$$\frac{c}{\nu(I)} \int_{2I} |f(t) - f_{2I}| d\nu(t).$$

Using (10) to estimate the last integral, and then applying (1), (4) and (5), we obtain (12).

Next, we will show that for a.e. $(d\nu)x \in I$,

$$(13) \quad |f(x) - f_{I_x}| \leq cF(I) \sup\{(\mu(J)/\nu(J)): x \in J \subset 2I\}.$$

To see this, set $I_k = 2^{-k}I_x$, and note that for a.e. $(d\nu)x \in I$, differentiation of the integral gives

$$|f(x) - f_{I_x}| \leq \sum_{k=0}^{\infty} |f_{I_k} - f_{I_{k+1}}|.$$

However,

$$\begin{aligned} |f_{I_k} - f_{I_{k+1}}| &\leq \frac{1}{\nu(I_{k+1})} \int_{I_{k+1}} |f(t) - f_{I_k}| d\nu(t) \\ &\leq \frac{c}{\nu(I_k)} \int_{I_k} |f(t) - f_{I_k}| d\nu(t) \end{aligned}$$

by (1) and the fact that $I_{k+1} \subset I_k$. Hence, by (10),

$$|f_{I_k} - f_{I_{k+1}}| \leq \frac{c}{\nu(I_k)} \mu(I_k) F(I_k).$$

Since $2^k I_k = I_x$, (4) and (5) imply that $F(I_k) \leq c\alpha^{-k} F(I)$. Combining estimates, we obtain that for a.e. $(d\nu)x \in I$,

$$\begin{aligned} |f(x) - f_{I_x}| &\leq cF(I) \sum_{k=0}^{\infty} \alpha^{-k} \frac{\mu(I_k)}{\nu(I_k)} \\ &\leq cF(I) \left\{ \sup \frac{\mu(J)}{\nu(J)} \right\} \sum_{k=0}^{\infty} \alpha^{-k}, \end{aligned}$$

where the sup is taken over all cubes J such that $x \in J$ and $J \subset 2I$. Since $\alpha > 1$, the series converges and (13) follows. Combining (12) and (13) concludes the proof of the lemma.

We shall also use the following fact.

LEMMA 2. *Let ν be a Borel measure which satisfies (1), and let f_ν^* denote the Hardy-Littlewood maximal function of f with respect to ν :*

$$f_\nu^*(x) = \sup \left\{ \frac{1}{\nu(I)} \int_I |f(t)| d\nu(t) : x \in I \right\}.$$

If $u \in A_p(d\nu)$, $1 < p < \infty$, then there is a constant c , independent of f , such that

$$\int_{R^n} f_\nu^*(x)^p u(x) d\nu(x) \leq c \int_{R^n} |f(x)|^p u(x) d\nu(x).$$

This is a restatement of Theorem 7 of [6].

We now turn to the proof of the theorem, first showing that the assumptions on $g d\mu$ and g^{-1} imply that μ satisfies (1). Hölder's inequality and $g^{-1} \in A_p(g d\mu)$ imply

$$(14) \quad \left[\int_I g d\mu \right]^p \leq \mu(I) \left[\int_I g^{p/(p-1)} d\mu \right]^{p-1} \leq c \left[\int_I g d\mu \right]^p.$$

Since the same relation holds for $2I$,

$$\mu(2I) \left[\int_I g^{p/(p-1)} d\mu \right]^{p-1} \leq c \left[\int_{2I} g d\mu \right]^p \leq c \left[\int_I g d\mu \right]^p,$$

by (1) for $g d\mu$. Hence, by the first inequality (14), $\mu(2I) \leq c\mu(I)$.

If we set $d\nu = g d\mu$, then hypothesis (6) of the theorem is the same as (10) of Lemma 1. Hence, by Lemma 1, for a.e. $(d\nu)x \in I$,

$$|f(x) - f_J| \leq cF(I) \sup \{ (\mu(J)/\nu(J)) : x \in J \subset 2I \},$$

where $f_I = f_{I,d\nu}$. If χ_{2I} denotes the characteristic function of $2I$, it follows from the formula $\mu(J) = \int_J g^{-1} d\nu$ that

$$(15) \quad |f(x) - f_I| \leq cF(I)[g^{-1}\chi_{2I}]_\nu^*(x) \quad \text{for a.e. } (d\nu)x \in I.$$

For $1 < p < \infty$, multiply both sides of (15) by $g(x)$, raise the result to the power p' , and then integrate over I with respect to $d\mu$, obtaining

$$(16) \quad \int_I \{|f - f_I|g\}^{p'} d\mu \leq cF(I)^{p'} \int_{R^n} \{[g^{-1}\chi_{2I}]_\nu^*\}^{p'} g^{p'-1} d\nu.$$

It is easy to see that the assumption $g^{-1} \in A_p(d\nu)$ is equivalent to $g^{p'-1} \in A_p(d\nu)$. Hence, Lemma 2 shows that the expression on the right in (16) is at most

$$cF(I)^{p'} \int_{2I} g^{-p'} g^{p'-1} d\nu = cF(I)^{p'} \mu(2I).$$

Since μ satisfies (1), we obtain

$$\int_I \{|f - f_I|g\}^{p'} d\mu \leq cF(I)^{p'} \mu(I),$$

as desired. If $p = 1$, the hypothesis $g^{-1} \in A_1(d\nu)$ gives

$$[g^{-1}]_\nu^* \leq cg^{-1} \quad \text{a.e.}$$

Combining this with (15) proves (8). Thus, the proof of the theorem is complete.

As a final remark, we note that the proof given above does not yield the results on bounded mean oscillation mentioned previously, since these correspond to $F(I) \equiv 1$, which fails to satisfy (5) (α must strictly exceed 1 there, in order that the series in Lemma 1 converge).

REFERENCES

1. S. Campanato, *Proprietà di hölderianità di alcune classi di funzioni*, Ann. Scuola Norm Sup. Pisa (3) **17** (1963), 175–188. MR **27** #6119.
2. J.-G. Cuerva, *Weighted H^p spaces*, Ph.D. dissertation, Washington Univ., St. Louis, Mo., 1975.
3. F. John and L. Nirenberg, *On functions of bounded mean oscillation*, Comm. Pure Appl. Math. **14** (1961), 415–426. MR **24** #A1348.
4. E. P. Lotkowski, *Lipschitz spaces with weights*, Ph.D. dissertation, Rutgers Univ., New Brunswick, N.J., 1975.
5. N. G. Meyers, *Mean oscillation over cubes and Hölder continuity*, Proc. Amer. Math. Soc. **15** (1964), 717–721. MR **29** #5969.
6. B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165** (1972), 207–226. MR **45** #2461.
7. B. Muckenhoupt and R. L. Wheeden, *Weighted bounded mean oscillation and the Hilbert transform*, Studia Math. **54** (1976), 221–237.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903
(Current address of R. L. Wheeden)

Current address (E. P. Lotkowski): Prudential Property and Casualty Insurance Company, One Woodbridge Center, Woodbridge, New Jersey 07095