REMARK ON MEASURABLE GRAPH THEOREMS

JÜRGEN LEHN

ABSTRACT. A theorem generalizing some known measurable graph theorems is proved.

1. Notations. Two measurable spaces $(X_1, \mathcal{Q}_1)$ and $(X_2, \mathcal{Q}_2)$ are isomorphic if there exists a bimeasurable one-to-one function from X_1 onto X_2 . A measurable space is called a standard measurable space if it is isomorphic with $(I, \mathcal{B}(I))$, the unit interval and its Borel σ -field. We call it a Blackwell space if it is isomorphic with some space $(A, \mathcal{B}(A))$, where A is an analytic subset of I and $\mathcal{B}(A)$ is the system of all intersections of A with sets in $\mathcal{B}(I)$. We claim that $(X, \mathcal{B}(X))$ is a standard measurable space if X is any nondenumerable complete separable metric space and $\mathcal{B}(X)$ its Borel σ -field. If $(X, \mathcal{Q}, \mu)$ is a measure space, then we write throughout the paper $\mathcal{Q}_\mu$ for the completion of $\mathcal{Q}$ with respect to μ .

2. Main results. Let $(X, \mathcal{Q})$ and $(Y, \mathcal{B})$ be measurable spaces, f a function $f: X \rightarrow Y$, $\text{gr}(f)$ the graph of f and $\mathcal{Q} \otimes \mathcal{B}$ the product σ -field. It is well known that $(\mathcal{Q}, \mathcal{B})$ -measurability of the function implies $\text{gr}(f) \in \mathcal{Q} \otimes \mathcal{B}$ if $\mathcal{B}$ is countably separated (see e.g. [5, I.3.4 and 5]). Concerning the reverse implication we shall prove in this paper two notes:

(A) Let $(X, \mathcal{Q})$ and $(Y, \mathcal{B})$ be Blackwell spaces and μ be a measure on $\mathcal{Q}$. If $\text{gr}(f) \in \mathcal{Q}_\mu \otimes \mathcal{B}$, then f is $(\mathcal{Q}_\mu, \mathcal{B})$ -measurable.

(B) Let $(X, \mathcal{Q}, \mu)$ be a σ -finite measure space and $(Y, \mathcal{B})$ a Blackwell space. If $\text{gr}(f) \in \mathcal{Q}_\mu \otimes \mathcal{B}$, then f is $(\mathcal{Q}_\mu, \mathcal{B})$ -measurable.

In [3] Buckley has proved a result corresponding to (A) on the more restrictive assumption that the spaces $(X, \mathcal{Q})$ and $(Y, \mathcal{B})$ are complete separable metric spaces with their Borel σ -fields. In the most important case of a σ -finite measure, Buckley's result is known from Bierlein's measurable graph theorem in [2], where $(Y, \mathcal{B})$ is assumed to be a standard measurable space and $(X, \mathcal{Q}, \mu)$ is any σ -finite measure space. In (B) Bierlein's assumption on $(Y, \mathcal{B})$ is generalized to the assumption of a Blackwell space.

Now (A) and (B) may be applied to give a more general version of Buckley's theorem, of the main part of Bierlein's measurable graph theorem and of the known fact that $\text{gr}(f) \in \mathcal{Q} \otimes \mathcal{B}$ is equivalent to $(\mathcal{Q}, \mathcal{B})$ -measurability.

Received by the editors May 27, 1976.

AMS (MOS) subject classifications (1970). Primary 28A20.

Key words and phrases. Measurable functions, measurable graph theorems, analytic sets, Blackwell spaces.

© American Mathematical Society 1977

bility of f if both spaces $(X, \mathcal{A})$ and $(Y, \mathfrak{B})$ are Blackwell spaces (cf. [5, II.4.3]). Since $\mathfrak{B}$ is countably separated if $(Y, \mathfrak{B})$ is a Blackwell space, the following theorem is a consequence of (A) and (B).

MEASURABLE GRAPH THEOREM. *Let $(X, \mathcal{A}, \mu)$ be a measure space, $(Y, \mathfrak{B})$ a Blackwell space and f a function $f: X \rightarrow Y$. If either the measure μ is σ -finite or the space $(X, \mathcal{A})$ is a Blackwell space, then the equivalence holds:*

f is $(\mathcal{A}_\mu, \mathfrak{B})$ -measurable if and only if $\text{gr}(f) \in \mathcal{A}_\mu \otimes \mathfrak{B}$.

3. Proofs. The proof of (A) is based on the following lemma. The proof of the lemma is omitted as a standard one.

LEMMA. *Let $(X, \mathcal{A}, \mu)$ be a measure space and $(Y, \mathfrak{B})$ a measurable space. If $G \in \mathcal{A}_\mu \otimes \mathfrak{B}$, then there exists a set $N_0 \in \mathcal{A}$ such that $\mu(N_0) = 0$ and $G \setminus (N_0 \times Y) \in \mathcal{A} \otimes \mathfrak{B}$.*

PROOF OF (A). Let $(X, \mathcal{A})$ and $(Y, \mathfrak{B})$ be Blackwell spaces and $\text{gr}(f) \in \mathcal{A}_\mu \otimes \mathfrak{B}$. Then, by the Lemma, X is the union of two disjoint sets $X_0 \in \mathcal{A}$ and $N_0 \in \mathcal{A}$ such that

$$\mu(N_0) = 0 \quad \text{and} \quad \text{gr}(f) \setminus (N_0 \times Y) \in \mathcal{A} \otimes \mathfrak{B}.$$

Let $\mathcal{A} \cap X_0$ be the σ -field of all intersections of X_0 with sets of $\mathcal{A}$. Since, by definition, $(X_0, \mathcal{A} \cap X_0)$ is a Blackwell space, too, we can apply the measurable graph theorem for Blackwell spaces (see [5, II.4.3]) to the function $f_0: X_0 \rightarrow Y$ defined as the restriction of f on X_0 . Since $\text{gr}(f_0) = \text{gr}(f) \setminus (N_0 \times Y)$ is an element of $(\mathcal{A} \cap X_0) \otimes \mathfrak{B}$, by this theorem, f_0 is an $(\mathcal{A} \cap X_0, \mathfrak{B})$ -measurable function. Then we have for any $B \in \mathfrak{B}$,

$$\begin{aligned} f^{-1}(B) &= (f^{-1}(B) \cap X_0) \cup (f^{-1}(B) \cap N_0) \\ &= f_0^{-1}(B) \cup (f^{-1}(B) \cap N_0) \in \mathcal{A}_\mu, \end{aligned}$$

because $f_0^{-1}(B) \in \mathcal{A}$, $f^{-1}(B) \cap N_0$ is a subset of N_0 and $\mu(N_0) = 0$. This shows that f is $(\mathcal{A}_\mu, \mathfrak{B})$ -measurable, and (A) is proved.

To prove (B) it is possible to apply a projection theorem (II.2.2 in [5] is a suitable one) and Choquet's theorem on capacities (see [4, Theorem 1]) as done by Bierlein [2] in the case of a standard measurable space $(Y, \mathfrak{B})$. However, to give a short proof, we claim that (B) is an immediate consequence of the following generalization of Aumann's selection theorem (see [1, p. 17]) proved in [6] (cf. Theorem 3, and for the difference in formulation see [5, III.2.3]):

SELECTION THEOREM. *Let $(X, \mathcal{A})$ be a measurable space having the property that $\mathcal{A}$ equals its universal completion, $(Y, \mathfrak{B})$ be a Blackwell space and $C \in \mathcal{A} \otimes \mathfrak{B}$. If the projection of C on X is all of X , then there exists an $(\mathcal{A}, \mathfrak{B})$ -measurable function $s: X \rightarrow Y$ such that $(x, s(x)) \in C$ for all $x \in X$.*

REMARKS. 1. That our theorem generalizes Buckley's theorem [3] and the main part of Bierlein's measurable graph theorem [2] is evident. But our

theorem generalizes also the known result that for Blackwell spaces $(X, \mathcal{Q})$ and $(Y, \mathcal{B})$, $\text{gr}(f) \in \mathcal{Q} \otimes \mathcal{B}$ is equivalent to $(\mathcal{Q}, \mathcal{B})$ -measurability of the function f . This can be seen if we define the measure μ on $\mathcal{Q}$ by $\mu = 0$ for the void set and $\mu = \infty$ for all other sets in $\mathcal{Q}$.

2. That neither in (A) nor in (B) the assumption on $(Y, \mathcal{B})$ to be a Blackwell space is superfluous can be seen by the following simple example: $X = Y = I$ (the unit interval), $\mathcal{Q}$ the Borel sets in I , $\mathcal{B}$ the σ -field generated by the Borel sets and one set V being nonmeasurable with respect to Lebesgue measure, and μ the Lebesgue measure on $\mathcal{Q}$. If f is the identity on I , then $\text{gr}(f) \in \mathcal{Q}_\mu \otimes \mathcal{B}$, but the function is not $(\mathcal{Q}_\mu, \mathcal{B})$ -measurable, since $f^{-1}(V) \notin \mathcal{Q}_\mu$.

ACKNOWLEDGMENTS. The author thanks the referee, Professor Tibor Neubrunn, for suggestions that led to an improvement of the paper.

REFERENCES

1. R. J. Aumann, *Measurable utility and measurable choice theorem* (Colloq. Internat., 1967), Centre Nat. Recherche Sci., Paris, 1969, pp. 15–26. MR 41 #6548.
2. D. Bierlein, *Der Graph messbarer Funktionen mit abstraktem Definitionsbereich*, Math. Z. 76 (1961), 468–471. MR 23 #A3817.
3. J. J. Buckley, *Graphs of measurable functions*, Proc. Amer. Math. Soc. 44 (1974), 78–80. MR 49 #508.
4. G. Choquet, *Forme abstraite du théorème de capacitabilité*, Ann. Inst. Fourier Grenoble 9 (1959), 83–89. MR 22 #3692b.
5. J. Hoffman-Jørgensen, *The theory of analytic sets*, Various Publication Series 10, Aarhus, 1970.
6. M.-F. Sainte-Beuve, *On the extension of von Neumann-Aumann's theorem*, J. Functional Analysis 17 (1974), 112–129. MR 51 #10564.

MATHEMATISCHES INSTITUT, UNIVERSITÄT KARLSRUHE, D-7500 KARLSRUHE, FEDERAL REPUBLIC OF GERMANY