COINCIDENT PAIRS OF CONTINUOUS SECTIONS IN PROFINITE GROUPS

H. APPELGATE AND H. ONISHI

ABSTRACT. Given a profinite group G and a closed subgroup H , we show there exist continuous sections $s: (G/H)_l \rightarrow G$ and $s': (G/H)_r \rightarrow G$ of the projections $G \rightarrow (G/H)_l$ and $G \rightarrow (G/H)_r$ onto the left and right coset spaces, respectively, such that $\text{im}(s) = \text{im}(s')$.

1. Coincident pairs of sections. Given a group G and a subgroup H , let $(G/H)_l$ and $(G/H)_r$ denote, respectively, the left and right coset spaces of H in G . A pair of sections

$$(G/H)_l \xrightarrow{s} G \xleftarrow{s'} (G/H)_r$$

will be called *coincident* if $\text{im}(s) = \text{im}(s')$. Having a coincident pair of sections is equivalent to having a subset S of G such that

$$G = \coprod_{x \in S} xH = \coprod_{x \in S} Hx \quad (\text{disjoint union})$$

($S = \text{im}(s) = \text{im}(s')$) or to having a 1-1 correspondence $aH \leftrightarrow Hb$ between $(G/H)_l$ and $(G/H)_r$ such that $aH \cap H_b \neq \emptyset$.

The following criterion is easy to establish (see [1]).

LEMMA 1. *There exists a coincident pair of sections for $G \bmod H$ if and only if for every a in G , $(H: H \cap aHa^{-1}) = (H: H \cap a^{-1}Ha)$ where, in general, $(G: H)$ denotes the cardinality of G/H (left or right coset space).*

Not every $G \supset H$ will admit a coincident pair of sections. We thank our colleague Peter Stebe for the following example. Let $G = \langle a, b \rangle$ be the free group on two generators a, b and H the subgroup generated by $a^n b a^{-n}$ for all $n \geq 0$. Then $aHa^{-1} < H$ (proper subgroup), $H < a^{-1}Ha$ and $(H: H \cap a^{-1}Ha) > 1$ but $(H: H \cap aHa^{-1}) = 1$.

When $(G: H) < \infty$ there is always a coincident pair of sections. More generally, if $N(H)$ is the normalizer of H and G is torsion mod $N(H)$, i.e. for every $x \in G$, $x^n \in N(H)$ for some $n > 0$, then there exists a coincident pair of sections. This can be seen using the following criterion. Given an inner automorphism $f: G \rightarrow G$ let $H_0 = H$, $H_n = f^{-n}(H) \cap H_{n-1}$ for $n > 0$. If for

Received by the editors December 30, 1974.

AMS (MOS) subject classifications (1970). Primary 20F20.

© American Mathematical Society 1977

every f there exists $n > 0$ such that $(H_{n-1} : H_n) = (H_{n-1} : f(H_n))$, then there exists a coincident pair of sections for $G \bmod H$. This criterion follows from Lemma 1 and the observation that for any subgroups K, L of H , $(K : K \cap L) = (L : K \cap L)$ implies $(H : K) = (H : L)$.

For finite groups G , one can prove the existence of coincident pairs of sections by applying the following well-known result from combinatorics. For a proof see, e.g., [2, pp. 50–51].

LEMMA 2. *If A is a nonempty finite set and $A = \coprod_{i \in I} B_i = \coprod_{i \in I} C_i$ are two partitions of A with $|B_i| = |C_i| = r$ ($|B|$ denotes number of elements) for all i in the finite index set I , then there is a bijection $i \mapsto j$ of I onto itself such that $B_i \cap C_j \neq \emptyset$.*

We will make use of this result later on.

2. Coincident pairs of sections in profinite groups. Let G be a profinite group and let U range over the open normal subgroups of G (for definitions and basic properties see, e.g., [3, Chapter I]). Then for any closed subgroups H, K of G , $H = \lim_U HU/U$ and $H \cap K = \lim_U (HU \cap KU)/U$ (lim denotes projective limit). Using these facts, it is not hard to prove $(H : H \cap aHa^{-1}) = (H : H \cap a^{-1}Ha)$ for any a in G . Thus, by Lemma 1, there exists a coincident pair of sections for $G \bmod H$.

On the other hand, it is well known that for any closed subgroup H of G there exists a continuous section $G/H \rightarrow G$. Our objective is to prove the following

THEOREM. *If G is a profinite group, then for any closed subgroup H of G there exists a coincident pair of continuous sections for $G \bmod H$.*

PROOF. Let $\mathfrak{S}$ be the set of all triples (N, s, s') where N is a closed normal subgroup of G and

$$(G/HN)_1 \xrightarrow{s} G/N \xleftarrow{s'} (G/HN)_r$$

is a coincident pair of continuous sections. By taking $N = G$ we see that $\mathfrak{S} \neq \emptyset$.

Given (N, s, s') and (K, t, t') in $\mathfrak{S}$, let $(N, s, s') \leq (K, t, t')$ if $N \supset K$ and the two squares

$$\begin{array}{ccccc} (G/HK)_1 & \xrightarrow{t} & G/K & \xleftarrow{t'} & (G/HK)_r \\ \downarrow & & \downarrow & & \downarrow \\ (G/HN)_1 & \xrightarrow{s} & G/N & \xleftarrow{s'} & (G/HN)_r \end{array}$$

are commutative; the vertical arrows are projections. Clearly $\leq$ is a partial order on $\mathfrak{S}$. Let (N_i, s_i, s'_i) be a chain in $\mathfrak{S}$. Put $N = \bigcap_i N_i$. Then $HN = \bigcap_i HN_i$

and $G/N = \lim_i G/N_i$. From these facts we easily get a pair (s, s') such that $(N, s, s') \in \mathfrak{S}$ and $(N_i, s_i, s'_i) \leq (N, s, s')$ for all i . We can now apply Zorn's lemma.

Let (N, s, s') be a maximal element of $\mathfrak{S}$. We want to show $N = 1$. Suppose $N \neq 1$. Then there exists an open normal subgroup V of G such that $N \not\subseteq V$. Let $N_1 = N \cap V$. Then $N_1 < N$ and $(N : N_1) < \infty$. We will produce a pair (t, t') such that $(N_1, t, t') \in \mathfrak{S}$ and $(N, s, s') < (N_1, t, t')$. Replacing G/N_1 by G , we may assume $N_1 = 1$. Then N is finite. We have the following diagram:

$$\begin{array}{ccccc}
 (G/H)_1 & \xleftarrow{\pi} & G & \xrightarrow{\pi'} & (G/H)_r \\
 q \downarrow & & p \downarrow & & q' \downarrow \\
 (G/HN)_1 & \xleftarrow{\varphi} & G/N & \xrightarrow{\varphi'} & (G/HN)_r \\
 \parallel & & \parallel & & \parallel \\
 (G/HN)_1 & \xrightarrow{s} & G/N & \xleftarrow{s'} & (G/HN)_r
 \end{array}$$

where $\text{im}(s) = \text{im}(s') = S$, say, and $p, q, q', \pi, \pi', \varphi, \varphi'$ are projections, and s, s' are continuous sections of φ, φ' , respectively.

Choose an open normal subgroup U of G such that $N \cap U = 1$ and $HN \cap U \subset H$. With some abuse of notation we put $aUN = p(aU)$, $aUH = \pi(aU)$, $HUa = \pi'(Ua)$, $aUHN = q\pi(aU)$ and $NHUA = q'\pi'(Ua)$ for $a \in G$. Since $N \cap U = 1$, p restricted to aU is a homeomorphism of aU onto aUN . Also, since $HN \cap U \subset H$, q is 1-1 on aUH and q' is 1-1 on HUa . Let $K = H \cap N$. Since N is finite, there are $n_i \in N, i \in I$ finite, such that $N = \coprod_i n_i K = \coprod_i K n_i$. Then for each $A = aN \in G/N, A = \coprod_i a n_i K = \coprod_i K n_i a$. Since A is finite and $|a n_i K| = |K n_i a| = |K|$, by Lemma 2, there is a bijection $i \mapsto j$ of I onto I such that $a n_i K \cap K n_j a \neq \emptyset$.

Now choose $A_1, \dots, A_m \in S = \text{im}(s) = \text{im}(s')$ such that $S \subset \coprod_k A_k \bar{U}$, where $\bar{U} = UN/N$. Then choose $a_k \in G$ such that $A_k = a_k N$ and choose a bijection $i \mapsto j$ of I onto I depending on k such that $a_k n_i K \cap K n_j a_k \neq \emptyset$ and finally choose an element b_{kij} in $a_k n_i K \cap K n_j a_k$. Since s and s' are continuous, $s^{-1}(A_k \bar{U})$ and $s'^{-1}(A_k \bar{U})$ are open and clearly

$$\begin{aligned}
 s^{-1}(A_k \bar{U}) &\subset a_k UHN, & (G/HN)_r &= \coprod_k s^{-1}(A_k \bar{U}), \\
 s'^{-1}(A_k \bar{U}) &\subset NHUa_k, & (G/HN)_r &= \coprod_k s'^{-1}(A_k \bar{U}).
 \end{aligned}$$

Now let

$$\begin{aligned}
 W_{ki} &= a_k n_i UH \cap q^{-1} s^{-1}(A_k \bar{U}), \\
 W'_{kj} &= HUn_j a_k \cap q'^{-1} s'^{-1}(A_k \bar{U}).
 \end{aligned}$$

Then these are open sets and

$$(G/H)_1 = \coprod_{k,i} W_{ki}, \quad (G/H)_r = \coprod_{k,j} W'_{kj}.$$

Now let $p_{kij}^{-1}: A_k \bar{U} = a_k UN \rightarrow b_{kij} U$ be the inverse of p on $b_{kij} U$ and define functions

$$(G/H)_1 \xrightarrow{t} G \xleftarrow{t'} (G/H)_r$$

by $t = p_{kij}^{-1}sq$ on W_{ki} and $t' = p_{kij}^{-1}s'q'$ on W'_{kj} . Clearly t and t' are continuous, $pt = sq$, $pt' = s'q'$. On each W_{ki} , $q\pi t = \varphi p p_{kij}^{-1}sq = \varphi sq = q$. Since $\pi t(W_{ki}) \subset a_k n_i UH$ and q is 1-1 on $a_k n_i UH$, we conclude that $\pi t = \text{id}$, i.e. that t is a section. Similarly t' is a section.

Finally, we claim that $\text{im}(t) = \text{im}(t')$, in fact $t(W_{ki}) = t'(W'_{kj})$ when $i \mapsto j$ for the given k . Let $a = t(aH)$, $aH \in W_{ki}$. Then $a \in bU$, where $b = b_{kij}$, say $a = bu$, $u \in U$. Write $b = a_k n_i x = x' n_j a_k$, $x, x' \in K$. Since $N \cap U = 1$, $nu = un$ for $n \in N$. Thus $a = bu = a_k u n_i x = x' n_j a_k u$. Thus $sq(aH) = pt(aH) = p(a) = a_k uN$ and hence $a = t(aH) = p_{kij}^{-1}sq(aH) = p_{kij}^{-1}(a_k uN)$. But since $a_k uN \in \text{im}(s) = \text{im}(s')$, $a_k uN = s'(NH a_k u) = s'q'(Ha) \in A_k \bar{U}$. Since $Ha \in HUn_j a$, this means that $Ha \in W'_{kj}$. Therefore $t'(Ha) = p_{kij}^{-1}s'q'(Ha) = p_{kij}^{-1}(a_k uN) = a$. This completes the proof of the theorem.

COROLLARY. *There exists a homeomorphism between $(G/H)_1$ and $(G/H)_r$, such that the corresponding cosets intersect.*

REFERENCES

1. O. Ore, *On coset representatives in groups*, Proc. Amer. Math. Soc. **9** (1958), 665–670. MR **20** #7068.
2. H. Ryser, *Combinatorial mathematics*, Carus Math. Monographs, no. 14, Wiley, New York, 1963. MR **27** #51.
3. J.-P. Serre, *Cohomologie galoisienne*, Lecture Notes in Math., vol. 5, Springer-Verlag, Berlin and New York, 1964. MR **31** #4785.

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF NEW YORK, CITY COLLEGE, NEW YORK, NEW YORK 10031