LOWER BOUNDS FOR THE ZEROS OF BESSEL FUNCTIONS

ROGER C. MCCANN

ABSTRACT. Let $j_{p,n}$ denote the n th positive zero of J_p , $p > 0$. Then

$$j_{p,n} > (j_{0,n}^2 + p^2)^{1/2}.$$

We begin by considering the eigenvalue problem

$$(1) \quad -(xy')' + x^{-1}y = \lambda^2 x^{2p-1}y, \quad \lambda, p > 0,$$

$$(2) \quad y(a) = y(1) = 0, \quad 0 < a < 1.$$

For simplicity of notation we will set $q = p^{-1}$. It is easily verified that the general solution of (1) is

$$y(x) = C_1 J_q(\lambda q x^{1/q}) + C_2 Y_q(\lambda q x^{1/q})$$

and that the eigenvalues are given by

$$J_q(\lambda q) Y_q(\lambda q a^{1/q}) - J_q(\lambda q a^{1/q}) Y_q(\lambda q) = 0.$$

If $z_n(a, r)$ denotes the n th positive zero of $J_r(z) Y_r(z a^{1/q}) - J_r(z a^{1/q}) Y_r(z) = 0$, then the n th eigenvalue, $\lambda_n^2(a)$, of (1), (2) is given by

$$(3) \quad \lambda_n^2(a) = (z_n(a, q)/q)^2.$$

Let $j_{r,n}$ denote the n th positive zero of J_r . On p. 38 of [4] it is shown that $z_n(a, r) \rightarrow j_{r,n}$ as $a \rightarrow 0^+$ whenever r is a positive integer. The restriction on r is extrinsic so that

$$(4) \quad \lim_{a \rightarrow 0^+} z_n(a, r) = j_{r,n}, \quad r \geq 0.$$

Let $R[p, y]$ denote the Rayleigh quotient

$$R[p, y] = \int_a^1 (-(xy')' + x^{-1}y)y \, dx / \int_a^1 x^{2p-1}y^2 \, dx.$$

It is well known that the eigenvalues $\{\lambda_n^2(p)\}$ of (1), (2) can be obtained from the Rayleigh quotient [5]. Let V denote the linear space of all functions in $C^2((a, 1))$ which satisfy the boundary conditions (2). Then

$$\lambda_1^2(p) = \min_{y \in V, y \neq 0} R[p, y].$$

Let $y_1, y_2, \dots, y_n$ be n functions in V , A denote the subspace of V spanned by $y_1, y_2, \dots, y_n$ and $A^\perp$ denote the orthogonal complement of A relative to V . Then

Received by the editors January 5, 1976 and, in revised form, September 13, 1976.

AMS (MOS) subject classifications (1970). Primary 33A40.

© American Mathematical Society 1977

$$\lambda_{n+1}^2(p) = \max_A \min_{y \in A^\perp, y \neq 0} R[p, y]$$

where the maximum is taken over all sets of n functions in V .

Whenever $p > 0$ we have that $x^{2p-1} < x^{-1}$ for all $x \in (0, 1)$. Then

$$(5) \quad R[p, y] = \frac{\int_a^1 - (xy')' y \, dx}{\int_a^1 x^{2p-1} y^2 \, dx} + \frac{\int_a^1 x^{-1} y^2 \, dx}{\int_a^1 x^{2p-1} y^2 \, dx} \geq Q[p, y] + 1,$$

where

$$Q[p, y] = \int_a^1 - (xy')' y \, dx / \int_a^1 x^{2p-1} y^2 \, dx$$

is the Rayleigh quotient for the eigenvalue problem

$$(6) \quad - (xy')' = \mu^2 x^{2p-1} y,$$

$$(7) \quad y(a) = y(1) = 0,$$

Equation (6) is equivalent to

$$(8) \quad x^2 y'' + yx' + \mu^2 x^{2p} y = 0.$$

It is easily checked that the general solution of (8) and, hence, of (6) is (recall that $q = p^{-1}$)

$$y(x) = C_1 J_0(\mu q x^{1/q}) + C_2 Y_0(\mu q x^{1/q})$$

and that the eigenvalues are given by

$$J_0(\mu q) Y_0(\mu q a^{1/q}) - J_0(\mu q a^{1/q}) Y_0(\mu q) = 0.$$

In particular the n th eigenvalue, $\mu_n^2(a)$, of (6), (7) is given by

$$(9) \quad \mu_n^2(a) = (z_n(a, 0)/q)^2.$$

From (3), (5), and (9) we obtain

$$(10) \quad (z_n(a, q)/q)^2 \geq (z_n(a, 0)/q)^2 + 1.$$

If we now replace q by p , let $a \rightarrow 0^+$ in (10), and using (4) we find that $(j_{p,n}/p)^2 \geq (j_{0,n}/q)^2 + 1$.

THEOREM. $j_{p,n} \geq ((j_{0,n})^2 + p^2)^{1/2}$ whenever $p \geq 0$.

COROLLARY. $j_{p,n} > ((n - \frac{1}{4})^2 \pi^2 + p^2)^{1/2}$ whenever $p \geq 0$.

PROOF. It is known (see [9, p. 489]) that the positive zeros of J_0 lie in the intervals $(m\pi + \frac{3}{4}\pi, m\pi + \frac{7}{8}\pi)$ for $m = 0, 1, 2, \dots$. Hence, $j_{0,n} > (n-1)\pi + \frac{3}{4}\pi = (n - \frac{1}{4})\pi$. The desired result follows.

In [8] it is shown that

$$j_{p,n} = p + a_n p^{1/3} + b_n p^{-1/3} + O(p^{-1}) \quad (n = 1, 2, \dots),$$

where a_n and b_n are independent of p . Hence,

$$j_{p,n}^2 = p^2 + c_n p^{4/3} + O(p^{2/3}) \quad (n = 1, 2, \dots),$$

where c_n is independent of p . This shows that the second term of the lower

bound for $j_{p,n}$ given in the Theorem is of the wrong order. Other asymptotic expansions for $j_{p,n}$ may be found in [1], [2], and [6].

In [3] it is shown that for $0 \leq p \leq \frac{1}{2}$

$$(11) \quad p\pi/2 + \left(n - \frac{1}{4}\right)\pi \leq j_{p,n}.$$

For $p = 0$ the result of the Theorem is exact, while the expression in (11) has a strict inequality. Hence, our result is stronger than (11) whenever p is sufficiently small. However, when $p = \frac{1}{2}$, the result in (11) is exact. Hence, for $0 \leq p \leq \frac{1}{2}$ neither result implies the other. It should be emphasized that the Theorem is valid for all $p \geq 0$, while (11) is valid only for $0 \leq p \leq \frac{1}{2}$.

I would like to thank the referee for his helpful suggestions and for bringing [2], [3], and [4] to my attention.

REFERENCES

1. T.H. Boyer, *Concerning the zeros of some functions related to Bessel functions*, J. Mathematical Phys. **10** (1969), 1729–1744. MR **41** #2074.
2. H.W. Hethcote, *Error bounds for asymptotic approximations of zeros of transcendental functions*, SIAM J. Math. Anal. **1** (1970), 147–152. MR **42** #744.
3. ———, *Bounds for zeros of some special functions*, Proc. Amer. Math. Soc. **25** (1970), 72–74. MR **41** #569.
4. M. Kline, *Some Bessel equations and their application to guide and cavity theory*, J. Mathematical Phys. **27** (1948), 37–48. MR **9**, 431.
5. S.G. Mikhlin, *Variational methods in mathematical physics*, GITTL, Moscow, 1957; English transl., Macmillan, New York, 1964. MR **22** #1981; **30** #2712.
6. F.W.J. Olver, *The asymptotic expansion of Bessel functions of large order*, Philos. Trans. Roy. Soc. London Ser. A **247** (1954), 328–368. MR **16**, 696.
7. ——— (Editor), *Bessel functions*. Part III, Roy. Soc. Math. Tables, Vol. 7, Cambridge Univ. Press, Cambridge, 1960.
8. F. Tricomi, *Sulle funzioni di Bessel di ordine e argomento pressochè uguali*, Atti. Accad. Sci. Torino Cl. Sci. Fis. Mat. Nat. **83** (1949), 3–20. MR **11**, 594.
9. G.N. Watson, *A treatise on the theory of Bessel functions*, Cambridge Univ. Press, Cambridge, 1958.

DEPARTMENT OF MATHEMATICS, CASE WESTERN RESERVE UNIVERSITY, CLEVELAND, OHIO 44106