RELATIONSHIPS BETWEEN CONTINUUM NEIGHBORHOODS IN INVERSE LIMIT SPACES AND SEPARATIONS IN INVERSE LIMIT SEQUENCES

HARVEY S. DAVIS

ABSTRACT. The main result of this paper is the following theorem. Let $\{X_\alpha, f_{\alpha\beta}, \alpha, \beta \in I\}$ be an inverse system of compact Hausdorff spaces and continuous onto maps with inverse limit X . Let $p \in X$ and A be closed in X . There exists a continuum neighborhood of p disjoint from A if and only if there exists $\alpha \in I$ and disjoint sets U and V open in X_α , neighborhoods respectively of p_α and A_α such that for all $\beta \geq \alpha$, $f_{\alpha\beta}^{-1}(U)$ lies in a single component of $X_\beta - f_{\alpha\beta}^{-1}(V)$. This is Theorem B of the text.

Introduction. Throughout this paper the conventions and elementary results on inverse limits of topological spaces are those given in Bourbaki [3] except when noted otherwise. $(I, \leq)$ is a directed set, $\{X_\alpha | \alpha \in I\}$ is a collection of nonvoid compact Hausdorff spaces and, for every pair (α, β) where $\alpha \leq \beta$, $f_{\alpha\beta}: X_\beta \rightarrow X_\alpha$ is continuous. The maps $f_{\alpha\beta}$ are not necessarily onto. $(X_\alpha, f_{\alpha\beta})$ is an inverse system of topological spaces with inverse limit X and canonical maps $f_\alpha: X \rightarrow X_\alpha$, for all α . If $A \subseteq X$, $p \in X$ then denote $f_\alpha(A) = A_\alpha$ and $f_\alpha(p) = p_\alpha$.

If A is any subset of a topological space S then $\mathcal{T}(A)$ will denote the collection of all open (in S) subsets of A and $\mathcal{F}(A)$ will denote the collection of all closed (in S) subsets of A . $\text{Int}(A)$ will denote the interior of A and $\text{Clo}(A)$ will denote the closure of A .

W is a *subcontinuum* of a topological space S if and only if W is compact, Hausdorff and connected. W is a *continuum neighborhood* of a point p of S if and only if p is an element of the interior of W . If A is any subset of S then $T(A)$ will denote the set of those points for which there does not exist a continuum neighborhood disjoint from A .

Let $\alpha \in I$. $\{W_\beta | \beta \geq \alpha\}$ denotes an *inverse system of subcontinua* if and only if for all $\beta \geq \alpha$, W_β is a subcontinuum of X_β , and $(W_\beta, f_{\alpha\beta} | W_\beta)$ is an inverse system.

Let S be a compact Hausdorff space. The following dictionary provides translations of some common properties of S into properties of the set function T defined on $\{A | A \subseteq S\}$.

Received by the editors September 29, 1975 and, in revised form, July 26, 1976.

AMS (MOS) subject classifications (1970). Primary 54F15; Secondary 54B25.

Key words and phrases. Continuum neighborhood, set function T , inverse limit, compact Hausdorff space.

© American Mathematical Society 1977

S is *connected im kleinem* at $p \in S$ if and only if for every $A \in \mathfrak{F}(S)$, if $p \notin A$ then $p \notin T(A)$.

S is *semi-locally connected* at $p \in S$ if and only if $T(p) = p$.

S is *aprosyndetic* at q with respect to p if and only if $q \notin T(p)$.

S is an *indecomposable continuum* if and only if for every $p \in S$, $T(p) = S$.

Other material on the set function T can be found in [1], [2], [4], [5].

A. Fundamental theorems. Theorems A1 and A2 establish necessary and sufficient conditions on the inverse limit sequence that a given point p is not an element of $T(A)$ for a given closed set A .

LEMMA A1. *Let $p \in X$ and $A \in \mathfrak{F}(X)$. If there exists $\alpha \in I$, $U \in \mathfrak{T}(X_\alpha)$, $V \in \mathfrak{T}(X_\alpha)$ and an inverse system of subcontinua $\{W_\beta | \beta \geq \alpha\}$ such that $p_\alpha \in U$, $A_\alpha \subseteq V$ and such that for all $\beta \geq \alpha$, $f_{\alpha\beta}^{-1}(U) \subseteq W_\beta \subseteq X_\beta - f_{\alpha\beta}^{-1}(V)$, then $p \notin T(A)$.*

PROOF. Let W be the canonical image of $\text{inv lim } W_\beta$ in X . Since each W_β is a continuum, W is a continuum. Since, for each $\beta \geq \alpha$, $f_\alpha^{-1}(U) \subseteq \text{Clo}(f_\alpha^{-1}(U)) \subseteq \text{Clo}(f_\beta^{-1}f_{\alpha\beta}^{-1}(U)) \subseteq f_\beta^{-1}(W_\beta)$ and since $W = \bigcap \{f_\beta^{-1}(W_\beta) | \beta \geq \alpha\}$, $f_\alpha^{-1}(U) \subseteq W$. Since $p \in f_\alpha^{-1}(U)$, $p \in \text{Int}(W)$. Since, for all $\beta \geq \alpha$,

$$f_\beta^{-1}(W_\beta) \subseteq X - f_\beta^{-1}f_{\alpha\beta}^{-1}(V) = X - f_\alpha^{-1}(V) = f_\alpha^{-1}(X_\alpha - V) \subseteq X - A,$$

$\bigcap \{f_\beta^{-1}(W_\beta) | \beta \geq \alpha\} \subseteq X - A$, so $W \cap A = \emptyset$. Therefore $p \notin T(A)$.

LEMMA A2. *Let $p \in X$ and $A \in \mathfrak{F}(X)$. If $p \notin T(A)$ then there exists $\alpha \in I$, $U \in \mathfrak{T}(X_\alpha)$, $V \in \mathfrak{T}(X_\alpha)$ and an inverse system of subcontinua $\{W_\beta | \beta \geq \alpha\}$ such that $p_\alpha \in U$, $A_\alpha \subseteq V$ and for all $\beta \geq \alpha$,*

$$f_{\alpha\beta}^{-1}(U) \cap f_\beta(X) \subseteq W_\beta \subseteq f_\beta(X) - f_{\alpha\beta}^{-1}(V).$$

PROOF. Since $p \notin T(A)$, there exists a subcontinuum W of X such that $p \in \text{Int}(W)$ and $W \cap A = \emptyset$. There exists $\alpha(p) \in I$, $U(p) \in \mathfrak{T}(X_{\alpha(p)})$ such that $p \in f_{\alpha(p)}^{-1}(U(p)) \subseteq \text{Int}(W)$ since $\{f_\alpha^{-1}(U) | \alpha \in I, U \in \mathfrak{T}(X_\alpha)\}$ is a basis for $\mathfrak{T}(X)$. Similarly, for each $x \in A$, there exists $\alpha(x) \in I$, $U(x) \in \mathfrak{T}(X_{\alpha(x)})$ such that $x \in f_{\alpha(x)}^{-1}(U(x)) \subseteq X - W$. Let $\{f_{\alpha(i)}^{-1}(U_i) | i = 1, \dots, n\}$ be a finite subcover of the open cover $\{f_{\alpha(x)}^{-1}(U(x)) | x \in A\}$ of the compact set A . Since I is a directed set, there exists $\alpha \in I$ such that $\alpha \geq \alpha(p)$ and for $i = 1, \dots, n$, $\alpha \geq \alpha(i)$. Let $U = f_{\alpha(p)\alpha}^{-1}(U(p))$ and $V = f_{\alpha(1)\alpha}^{-1}(U_1) \cup \dots \cup f_{\alpha(n)\alpha}^{-1}(U_n)$. For $\beta \geq \alpha$ let $W_\beta = f_\beta(W)$.

Let $\beta \geq \alpha$ and let $z \in f_{\alpha\beta}^{-1}(U) \cap f_\beta(X)$. Since $z \in f_\beta(X)$, there exists $x \in X$ such that $x_\beta = z$. Since $f_{\alpha(p)\beta}(z) \in U(p)$, $x_{\alpha(p)} \in U(p)$. Therefore $x \in f_{\alpha(p)}^{-1}(U(p)) \subseteq \text{Int}(W) \subseteq W$. Thus $z = x_\beta \in f_\beta(W) = W_\beta$. So $f_{\alpha\beta}^{-1}(U) \cap f_\beta(X) \subseteq W_\beta$. Now let $z \in W_\beta = f_\beta(W)$. Let $z = x_\beta$ for some $x \in W$. Clearly $z \in f_\beta(X)$. Suppose $z \in f_{\alpha\beta}^{-1}(V)$. There exists $i \leq n$ such that $f_{\alpha(i)\beta}(z) \in U_i$ so $x_{\alpha(i)} \in U_i$. Therefore $x \notin W$. This contradiction establishes that $W_\beta \subseteq f_\beta(X) - f_{\alpha\beta}^{-1}(V)$.

THEOREM A1. Let $p \in X$ and $A \in \mathcal{F}(X)$. The following are equivalent.

(a) $p \notin T(A)$.

(b) There exists $\alpha \in I$, $U \in \mathcal{T}(X_\alpha)$, $V \in \mathcal{T}(X_\alpha)$ and an inverse system of subcontinua $\{W_\beta \mid \beta \geq \alpha\}$ such that $p_\alpha \in U$, $A_\alpha \subseteq V$ and for all $\beta \geq \alpha$,

$$f_{\alpha\beta}^{-1}(U) \cap f_\beta(X) \subseteq W_\beta \subseteq f_\beta(X) - f_{\alpha\beta}^{-1}(V).$$

PROOF. This follows directly from Lemmas A1 and A2 once it is noted that if W is the canonical image of $\text{inv lim } W_\beta$ in X then for all β , $f_\beta(W) = f_\beta(X) \cap W_\beta$, $f_\beta(W)$ is a continuum and W is the canonical image of $\text{inv lim } f_\beta(W)$. The relations in Lemma A1 imply that $f_{\alpha\beta}^{-1}(U) \cap f_\beta(X) \subseteq f_\beta(W) \subseteq f_\beta(X) - f_{\alpha\beta}^{-1}(V)$.

THEOREM A2. Suppose that each $f_{\alpha\beta}$ is onto. Let $p \in X$ and $A \in \mathcal{F}(X)$. The following are equivalent.

(a) $p \notin T(A)$.

(b) There exists $\alpha \in I$, $U \in \mathcal{T}(X_\alpha)$, $V \in \mathcal{T}(X_\alpha)$ and an inverse system of subcontinua $\{W_\beta \mid \beta \geq \alpha\}$ such that $p_\alpha \in U$, $A_\alpha \subseteq V$ and such that for all $\beta \geq \alpha$,

$$f_{\alpha\beta}^{-1}(U) \subseteq W_\beta \subseteq X_\beta - f_{\alpha\beta}^{-1}(V).$$

PROOF. This follows directly from Theorem A1 once it is noted that, since each X_α is compact, the hypothesis that each $f_{\alpha\beta}$ is onto forces each f_α to be onto.

COROLLARY A1. Suppose that each $f_{\alpha\beta}$ is onto. Let J be a cofinal subset of I and let $A \in \mathcal{F}(X)$. Then

$$\cap \{f_\alpha^{-1}(T(A_\alpha)) \mid \alpha \in J\} \subseteq T(A).$$

PROOF. Suppose $p \in X - T(A)$. Let α , U , V and $\{W_\beta\}$ be as in Theorem A2. Since J is cofinal, there exists $\beta \in J$ such that $\beta \geq \alpha$. But then $p_\beta \in f_{\alpha\beta}^{-1}(U) \subseteq W_\beta \subseteq X_\beta - f_{\alpha\beta}^{-1}(V) \subseteq X_\beta - A_\beta$. Therefore $p_\beta \notin T(A_\beta)$ so $p \notin f_\beta^{-1}(T(A_\beta))$. Thus $p \notin \cap \{f_\gamma^{-1}(T(A_\gamma)) \mid \gamma \in J\}$.

COROLLARY A2. Suppose that each $f_{\alpha\beta}$ is onto. Let J be a cofinal subset of I and let $A \in \mathcal{F}(X)$. If for all $\gamma, \delta \in I$ where $\gamma \leq \delta$, $f_{\gamma\delta}$ is monotone, then

$$T(A) = \cap \{f_\gamma^{-1}(T(A_\gamma)) \mid \gamma \in J\}.$$

PROOF. Suppose $p \in X - \cap \{f_\gamma^{-1}(T(A_\gamma)) \mid \gamma \in J\}$. There exists α such that $p_\alpha \notin T(A_\alpha)$ so there exist U , $V \in \mathcal{T}(X_\alpha)$ and a continuum W such that $p_\alpha \in U \subseteq W \subseteq X_\alpha - V \subseteq X_\alpha - A_\alpha$. Now let $\beta \geq \alpha$ and let $W_\beta = f_{\alpha\beta}^{-1}(W)$. Since $f_{\alpha\beta}$ is monotone, W_β is a continuum. It is clear that $\{W_\beta \mid \beta \geq \alpha\}$ is an inverse system of subcontinua and also that $f_{\alpha\beta}^{-1}(U) \subseteq W_\beta \subseteq X_\beta - f_{\alpha\beta}^{-1}(V)$. Hence, by Theorem A2, $p \notin T(A)$ so $T(A) \subseteq \cap \{f_\gamma^{-1}(T(A_\gamma)) \mid \gamma \in J\}$. Com-

binning this with the above corollary gives the equality.

B. Separations in inverse limit sequences.

DEFINITION. Let M and N be subsets of a topological space S . M separates N in S if and only if there exist sets P and Q such that $S - M = P \cup Q$, $P \cap N \neq \emptyset \neq Q \cap N$ and $(P \cap \text{Clo}(Q)) \cup (Q \cap \text{Clo}(P)) = \emptyset$.

REMARKS. If $N \subseteq M$ then M does not separate N in S . If M does not separate N in S and $L \subseteq M$ then L does not separate N in S .

DEFINITIONS. Let $A \in \mathfrak{F}(X)$ and $p \in X$. $p \notin S(A)$ if and only if there exists $\alpha \in I$, $U \in \mathfrak{T}(X_\alpha)$ and $V \in \mathfrak{T}(X_\alpha)$ such that $p_\alpha \in U$, $A_\alpha \subseteq V$, $U \cap V = \emptyset$, and, for all $\beta \geq \alpha$, $f_{\alpha\beta}^{-1}(V)$ does not separate $f_{\alpha\beta}^{-1}(U)$ in X_β .

REMARK. Let $A \in \mathfrak{F}(X)$. It is immediate from the definitions that $A \subseteq S(A)$.

THEOREM B1. Let $A \in \mathfrak{F}(X)$. $T(A) \subseteq S(A)$.

PROOF. Suppose $p \notin S(A)$. Let α , U , V be as in the above definition of S . Now let $\beta \geq \alpha$. Since X_β is compact Hausdorff and $f_{\alpha\beta}^{-1}(V)$ does not separate $f_{\alpha\beta}^{-1}(U)$ in X_β , $f_{\alpha\beta}^{-1}(U)$ lies entirely in a single component W_β of $X_\beta - f_{\alpha\beta}^{-1}(V)$. It is clear that W_β is a subcontinuum of X_β , that $f_{\alpha\beta}^{-1}(U) \subseteq W_\beta$ and that $W_\beta \subseteq X_\beta - f_{\alpha\beta}^{-1}(V)$. For all $\beta \geq \alpha$, choose such a W_β . Now let $\gamma \geq \beta \geq \alpha$. Note that $f_{\beta\gamma}(W_\gamma) \subseteq f_{\beta\gamma}(X_\gamma - f_{\alpha\gamma}^{-1}(V)) \subseteq X_\beta - f_{\alpha\beta}^{-1}(V)$. Also note that $f_{\alpha\beta}^{-1}(U) = f_{\beta\gamma}(f_{\alpha\gamma}^{-1}(U)) \subseteq f_{\beta\gamma}(W_\gamma)$ so that $f_{\alpha\beta}^{-1}(U) \subseteq f_{\beta\gamma}(W_\gamma) \cap W_\beta$. Since W_γ is connected and $f_{\beta\gamma}(W_\gamma) \cap W_\beta \neq \emptyset$, $f_{\beta\gamma}(W_\gamma) \subseteq W_\beta$ so $\{W_\beta | f_{\beta\gamma} | W_\beta\}$ is an inverse of subcontinua. Therefore by Lemma A1, $p \notin T(A)$. Thus $T(A) \subseteq S(A)$.

THEOREM B2. Suppose that each $f_{\alpha\beta}$ is onto. Let $A \in \mathfrak{F}(X)$. $S(A) = T(A)$.

PROOF. Suppose $p \notin T(A)$. Let α , U , V and $\{W_\beta\}$ be as in Theorem A2. Note that $A_\alpha \subseteq V$, $p_\alpha \in U$ and that $U \cap V = \emptyset$. Now let $\beta \geq \alpha$. Since $f_{\alpha\beta}^{-1}(U) \subseteq W_\beta$ and $W_\beta \cap f_{\alpha\beta}^{-1}(V) = \emptyset$, it is clear that $f_{\alpha\beta}^{-1}(V)$ does not separate $f_{\alpha\beta}^{-1}(U)$ in X_β . Hence $p \notin S(A)$ and thus $S(A) \subseteq T(A)$. By Theorem B1, $S(A) = T(A)$.

The main result can now be stated.

THEOREM B. Suppose that each $f_{\alpha\beta}$ is onto. Let $A \in \mathfrak{F}(X)$, and $p \in X$. There exists a continuum neighborhood of p disjoint from A if and only if there exists $\alpha \in I$ and disjoint sets U and V open in X_α , neighborhoods respectively of p_α and A_α such that for all $\beta \geq \alpha$, $f_{\alpha\beta}^{-1}(U)$ lies in a single component of $X_\beta - f_{\alpha\beta}^{-1}(V)$.

PROOF. Once it is noted that in a compact Hausdorff space the failure of an open set G to separate an open set D is equivalent to the existence of a component K of $S - G$ that contains D the theorem follows as a corollary to Theorem B2.

BIBLIOGRAPHY

1. D. P. Bellamy and H. S. Davis, *Continuum neighborhoods and filterbases*, Proc. Amer. Math. Soc. **27** (1971), 371–374. MR **43** #2653.
2. D. E. Bennett, *A characterization of locally connectedness by means of the set function T* , Fund. Math. **86** (1974), 137–141. MR **50** #11189.
3. N. Bourbaki, *General topology*. Parts 1 and 2, Addison-Wesley, Reading, Mass., 1966. MR **34** #5044a, 5044b.
4. H. S. Davis, D. P. Stadtlander and P. M. Swingle, *Properties of the set function T^n* , Portugal. Math. **21** (1962), 113–133. MR **25** #5501.
5. H. S. Davis, *A note on connectedness im kleinen*, Proc. Amer. Math. Soc. **19** (1968), 1237–1241. MR **40** #8021.

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48824