OPERATORS IN THE COMMUTANT OF A REDUCTIVE ALGEBRA

ROBERT L. MOORE

ABSTRACT. Let $\mathcal{Q}$ be a reductive algebra. It is shown that there is a subspace $\mathfrak{M}$ that reduces $\mathcal{Q}$ and such that the commutant of $\mathcal{Q}|_{\mathfrak{M}}$ is selfadjoint and the commutant of $\mathcal{Q}|_{\mathfrak{M}^\perp}$ consists of hyporeductive operators. It is then shown that under a variety of conditions, if an operator T is in $\mathcal{Q}'$, then T^* is in $\mathcal{Q}'$.

In [5], we made use of a decomposition of reductive operators on Hilbert space to deduce some results concerning the selfadjointness of the commutants of such operators. In that paper we also observed that a similar decomposition of a reductive operator algebra is possible. In this paper we produce such a decomposition and use it to answer a number of questions first raised by Rosenthal [9].

By an algebra, we will mean a weakly closed subalgebra (with identity) of the algebra of all (bounded) operators on a separable Hilbert space. If $\mathcal{Q}$ is an algebra and $\mathfrak{M}$ a subspace of $\mathcal{H}$ then $\mathcal{Q}\mathfrak{M} = \{Af: A \in \mathcal{Q} \text{ and } f \in \mathfrak{M}\}$, and $\mathfrak{M}$ is *invariant* for $\mathcal{Q}$ if $\mathcal{Q}\mathfrak{M} \subseteq \mathfrak{M}$. The lattice of subspaces invariant for $\mathcal{Q}$ is denoted by $\text{Lat } \mathcal{Q}$. We also denote by $\mathcal{Q}^*$ the algebra $\{A^*: A \in \mathcal{Q}\}$, and by $\mathcal{Q}'$ the algebra $\{B: AB = BA \text{ for all } A \in \mathcal{Q}\}$. Finally, $\mathcal{Q}$ is *reductive* if $\mathfrak{M} \in \text{Lat } \mathcal{Q}$ implies $\mathfrak{M}^\perp \in \text{Lat } \mathcal{Q}$, or equivalently, if $\text{Lat } \mathcal{Q} = \text{Lat } \mathcal{Q}^*$.

The results for which we are aiming are Theorems 2 and 3, which state that for every reductive algebra $\mathcal{Q}$ there is a subspace $\mathfrak{M}_0 \in \text{Lat } \mathcal{Q} \cap \text{Lat } \mathcal{Q}'$, such that $\mathcal{Q}'|_{\mathfrak{M}_0}$ is selfadjoint and $\text{Lat}(\mathcal{Q}|_{\mathfrak{M}_0^\perp}) \subseteq \text{Lat}(\mathcal{Q}'|_{\mathfrak{M}_0^\perp})$. Before we begin, we remark that certain techniques developed by Hoover [4] will also yield Theorems 2 and 3.

THEOREM 1. *Let $\mathcal{Q}$ be a reductive algebra and $\mathfrak{M} \in \text{Lat } \mathcal{Q}$. Let $X: \mathfrak{M}^\perp \rightarrow \mathfrak{M}$ and suppose that $T \in \mathcal{Q}'$, where T has the form*

$$T = \begin{pmatrix} 0 & X \\ 0 & 0 \end{pmatrix}$$

according to the decomposition $\mathcal{H} = \mathfrak{M} \oplus \mathfrak{M}^\perp$. Then

- (1) $T^* \in \mathcal{Q}'$.
- (2) $(\text{ran } T)^\perp \in \text{Lat } \mathcal{Q}$ and $(\mathcal{Q}|_{(\text{ran } T)^\perp})'$ is selfadjoint.
- (3) $\ker T \in \text{Lat } \mathcal{Q}$ and $(\mathcal{Q}|_{\ker^\perp T})'$ is selfadjoint.

Received by the editors February 8, 1977.

AMS (MOS) subject classifications (1970). Primary 47C05; Secondary 47D99.

Key words and phrases. Reductive algebra, hyperinvariant subspace, hyporeductive operator.

© American Mathematical Society 1977

PROOF. (1) All decompositions of vectors and operators are with respect to $\mathfrak{M}$ and $\mathfrak{M}^\perp$. Let $\mathfrak{N}$ be the subspace $\{\langle Xf, f \rangle : f \in \mathfrak{M}^\perp\}$. Since $T \in \mathcal{Q}'$ it is easy to check that $\mathfrak{N} \in \text{Lat } \mathcal{Q}$, and thus the subspace $\mathfrak{N}^\perp = \{\langle g, -X^*g \rangle : g \in \mathfrak{N}\}$ also lies in $\text{Lat } \mathcal{Q}$.

If $A \in \mathcal{Q}$ we can decompose A as $A_1 \oplus A_2$, since $\mathcal{Q}$ is *reductive*. If $g \in \mathfrak{N}$ we have

$$(A_1 \oplus A_2)\langle g, -X^*g \rangle = \langle A_1g, -A_2X^*g \rangle.$$

Since this vector must lie in $\mathfrak{N}^\perp$ it follows that $-X^*A_1g = -A_2X^*g$ for any $g \in \mathfrak{N}$, and hence that $X^*A_1 = A_2X^*$, which implies that $T^* \in \mathcal{Q}'$.

(2) Since $T \in \mathcal{Q}'$, $(\text{ran } T)^- \in \text{Lat } \mathcal{Q}$. Let C be an operator on $(\text{ran } T)^-$ that lies in the commutant of $\mathcal{Q}|(\text{ran } T)^-$; we want to show that $C^* \in (\mathcal{Q}|(\text{ran } T)^-)'$. Let C_1 be the operator on $\mathcal{H}$ defined by

$$C_1f = \begin{cases} Cf, & f \in (\text{ran } T)^-, \\ 0, & f \in \text{ran}^\perp T. \end{cases}$$

Then $C_1 \in \mathcal{Q}'$ and it will suffice to show that $C_1^* \in \mathcal{Q}'$.

Let $T_1 = C_1T$; then $T_1 \in \mathcal{Q}'$. Moreover, since $\mathfrak{M}$ is invariant under C_1 , it is easy to see that T_1 has the same form as T , namely

$$T_1 = \begin{pmatrix} 0 & * \\ 0 & 0 \end{pmatrix}.$$

It follows by part (1) that $T_1^* \in \mathcal{Q}'$, that is, for any $A \in \mathcal{Q}$, $T^*C_1^*A = AT^*C_1^*$. Since $T^* \in \mathcal{Q}'$, $T^*(AC_1^* - C_1^*A) = 0$, that is, $\text{ran}(AC_1^* - C_1^*A) \subseteq \ker T^*$. On the other hand $\text{ran } C_1^* \subseteq (\text{ran } T)^-$ and $(\text{ran } T)^-$ reduces A , so that

$$\text{ran}(AC_1^* - C_1^*A) \subseteq (\text{ran } T)^- = \ker^\perp T^*.$$

We conclude that $AC_1^* - C_1^*A = 0$, and $C_1^* \in \mathcal{Q}'$.

(3) Let U be the unitary operator $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Since $T^* \in \mathcal{Q}'$, we have

$$\begin{pmatrix} 0 & X^* \\ 0 & 0 \end{pmatrix} = U^*T^*U \in (U^*\mathcal{Q}U)'.$$

By part (2), $[(U^*\mathcal{Q}U)|(\text{ran } U^*T^*U)^-]'$ is selfadjoint. Thus $[\mathcal{Q}|(\text{ran } T^*)^-]'$ is selfadjoint and (3) follows because $(\text{ran } T^*)^- = \ker^\perp T$.

COROLLARY. Let $\mathcal{Q}$ be *reductive* and $\mathfrak{M} \in \text{Lat } \mathcal{Q}$. Let $Y: \mathfrak{M} \rightarrow \mathfrak{M}^\perp$ and suppose that $S \in \mathcal{Q}'$, where $S = \begin{pmatrix} 0 & Y \\ Y^* & 0 \end{pmatrix}$. Then

- (1) $S^* \in \mathcal{Q}'$.
- (2) $\ker S \in \text{Lat } \mathcal{Q}$ and $(\mathcal{Q}|\ker^\perp S)'$ is selfadjoint.
- (3) $(\text{ran } S)^- \in \text{Lat } \mathcal{Q}$ and $(\mathcal{Q}|\text{ran } S)'$ is selfadjoint.

PROOF. Consider adjoints and apply the theorem.

THEOREM 2. Let $\mathcal{Q}$ be a *reductive algebra*. There is a subspace $\mathfrak{M}_0$ such that

- (1) $\mathfrak{M}_0 \in \text{Lat } \mathcal{Q}$;
- (2) $(\mathcal{Q}|\mathfrak{M}_0)'$ is selfadjoint;
- (3) there is no nonzero subspace $\mathfrak{N} \subseteq \mathfrak{M}_0^\perp$ with properties (1) and (2).

Moreover, this subspace $\mathfrak{M}_0$ reduces $\mathcal{Q}'$ as well.

PROOF. Let $\mathcal{F} = \{\mathfrak{M} \in \text{Lat } \mathcal{Q}: (\mathcal{Q}|\mathfrak{M})' \text{ is selfadjoint}\}$. The family $\mathcal{F}$ is nonempty since it contains the zero subspace. Suppose that $\{\mathfrak{M}_\alpha\}_{\alpha \in B}$ is a chain in $\mathcal{F}$; in order to apply Zorn's lemma we would like to show that $\mathfrak{M} = \bigvee \{\mathfrak{M}_\alpha: \alpha \in B\}$ is also in $\mathcal{F}$, for which it suffices to show that $(\mathcal{Q}|\mathfrak{M})'$ is selfadjoint.

Let T be an operator on $\mathfrak{M}$ such that $T \in (\mathcal{Q}|\mathfrak{M})'$. To show that $T^* \in (\mathcal{Q}|\mathfrak{M})'$ we must show that for all $A \in \mathcal{Q}|\mathfrak{M}$ and for all $f \in \mathfrak{M}$ we have $T^*Af = AT^*f$, and in fact it will be enough to show this equality for all $f \in \bigcup \{\mathfrak{M}_\alpha: \alpha \in B\}$, because this set is dense in $\mathfrak{M}$. On the other hand, in this case $f \in \mathfrak{M}_\beta$ for some $\beta \in B$.

Decompose $\mathfrak{M}$ as $\mathfrak{M}_\beta \oplus (\mathfrak{M} \ominus \mathfrak{M}_\beta)$; then (since $\mathfrak{M}_\beta \in \text{Lat } \mathcal{Q}$) if $A \in \mathcal{Q}|\mathfrak{M}$ we have

$$A = \begin{pmatrix} A_1 & 0 \\ 0 & A_2 \end{pmatrix} \quad \text{and} \quad T = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix}.$$

Because $f \in \mathfrak{M}_\beta$ we have

$$T^*Af = \langle T_{11}^*A_1f, T_{12}^*A_1f \rangle \quad \text{and} \quad AT^*f = \langle A_1T_{11}^*f, A_2T_{12}^*f \rangle.$$

Since $T_{11} \in (\mathcal{Q}|\mathfrak{M}_\beta)'$ and $(\mathcal{Q}|\mathfrak{M}_\beta)'$ is selfadjoint, we have $T_{11}^*A_1 = A_1T_{11}^*$. Furthermore, the operator $\begin{pmatrix} 0 & T_{12} \\ 0 & 0 \end{pmatrix}$ lies in $(\mathcal{Q}|\mathfrak{M})'$ and by Theorem 1, so does its adjoint $\begin{pmatrix} 0 & 0 \\ T_{21}^* & 0 \end{pmatrix}$ and it follows that $A_2T_{12}^* = T_{21}^*A_1$. Thus $T^*Af = AT^*f$, $T^* \in (\mathcal{Q}|\mathfrak{M})'$, and $\mathfrak{M} \in \mathcal{F}$.

By Zorn's lemma there exists a maximal element $\mathfrak{M}_0$ of $\mathcal{F}$. $\mathfrak{M}_0$ automatically satisfies requirements (1) and (2) of the theorem. Suppose there is a nonzero subspace $\mathfrak{M}_1$ of $\mathfrak{M}_0^\perp$ for which (1) and (2) hold. We assert that $\mathfrak{M}_0 \oplus \mathfrak{M}_1$ lies in $\mathcal{F}$, a fact which contradicts the maximality of $\mathfrak{M}_0$. We must show that $(\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$ is selfadjoint. If S is an operator on $\mathfrak{M}_0 \oplus \mathfrak{M}_1$ such that $S \in (\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$ then we decompose S as

$$\begin{pmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix},$$

where $S_{11} \in (\mathcal{Q}|\mathfrak{M}_0)'$ and $S_{22} \in (\mathcal{Q}|\mathfrak{M}_1)'$. By assumption $S_{11}^* \in (\mathcal{Q}|\mathfrak{M}_0)'$ and $S_{22}^* \in (\mathcal{Q}|\mathfrak{M}_1)'$. Moreover, the operator

$$\begin{pmatrix} 0 & S_{12} \\ 0 & 0 \end{pmatrix}$$

satisfies the hypotheses of Theorem 1 for the reductive algebra $(\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)$. Thus by that theorem the operator

$$\begin{pmatrix} 0 & 0 \\ S_{12}^* & 0 \end{pmatrix}$$

lies in $(\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$. Similarly we use the corollary to Theorem 1 to show that $\begin{pmatrix} 0 & S_{21}^* \\ 0 & 0 \end{pmatrix}$ lies in $(\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$. Since

$$S^* = \begin{pmatrix} S_{11}^* & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & S_{22}^* \end{pmatrix} + \begin{pmatrix} 0 & S_{21}^* \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ S_{12}^* & 0 \end{pmatrix}$$

we see that $S^* \in (\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$, and thus that $(\mathcal{Q}|\mathfrak{M}_0 \oplus \mathfrak{M}_1)'$ is selfadjoint.

Finally, to show that $\mathfrak{M}_0 \in \text{Lat } \mathcal{Q}'$, suppose that $T \in \mathcal{Q}'$ and write

$$T = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix}$$

according to the decomposition $\mathcal{H} = \mathfrak{M}_0 \oplus \mathfrak{M}_0^\perp$.

Since $\mathfrak{M}_0$ reduces $\mathcal{Q}$ the operator

$$S = \begin{pmatrix} 0 & T_{12} \\ 0 & 0 \end{pmatrix}$$

also lies in $\mathcal{Q}'$, and by Theorem 1, $\ker S$ reduces $\mathcal{Q}$ and $(\mathcal{Q}|\ker^\perp S)'$ is selfadjoint. However, $\ker^\perp S$ is a subspace of $\mathfrak{M}_0^\perp$ so by the maximality of $\mathfrak{M}_0$ it must be that $\ker^\perp S = \{0\}$, that is, $T_{12} = 0$. Similarly we can show that $T_{21} = 0$ and thus $\mathfrak{M}_0$ reduces T .

We use the notation $\text{Red } \mathfrak{B}$ to mean $\text{Lat } \mathfrak{B} \cap \text{Lat } \mathfrak{B}^*$, where $\mathfrak{B}$ is any algebra.

THEOREM 3. *Let $\mathcal{Q}$ be reductive and suppose that for no nonzero subspace $\mathfrak{M}$ in $\text{Lat } \mathcal{Q}$ is it true that $(\mathcal{Q}|\mathfrak{M})'$ is selfadjoint. Then $\text{Lat } \mathcal{Q} \subseteq \text{Red } \mathcal{Q}'$.*

PROOF. Suppose $\mathfrak{M} \in \text{Lat } \mathcal{Q}$ and $T \in \mathcal{Q}'$. Decompose $\mathcal{H}$ as $\mathfrak{M} \oplus \mathfrak{M}^\perp$ and T as

$$T = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix};$$

let

$$S = \begin{pmatrix} 0 & T_{12} \\ 0 & 0 \end{pmatrix}.$$

By Theorem 1, $(\text{ran } S)^\perp \in \text{Lat } \mathcal{Q}$ and $(\mathcal{Q}|(\text{ran } S)^\perp)'$ is selfadjoint. Hence, by the hypothesis of the theorem, $\text{ran } S = \{0\}$, that is, $T_{12} = 0$. Similarly $T_{21} = 0$ and $\mathfrak{M} \in \text{Red } \mathcal{Q}'$.

C. K. Fong [1] has used the word *hyporeductive* to refer to an operator T such that every hyperinvariant subspace of T reduces T .

COROLLARY. *Let $\mathcal{Q}$ be as in Theorem 3. If $T \in \mathcal{Q}'$ and $\mathfrak{M}$ is hyperinvariant for T then $\mathfrak{M} \in \text{Red } \mathcal{Q}'$. In particular, T is hyporeductive.*

It follows from all the above that if $\mathcal{Q}$ is any reductive algebra and $T \in \mathcal{Q}'$, then $T = T_1 \oplus T_2$ where $T_1^* \in (\mathcal{Q}|\mathfrak{M}_0)'$ and T_2 is hyporeductive. Thus if we desire to show that $T^* \in \mathcal{Q}'$, it suffices to show that $T_2^* \in (\mathcal{Q}|\mathfrak{M}_0^\perp)'$.

In [9], P. Rosenthal introduced the following property which an operator T may have in connection with reductive algebras:

(P) If $\mathcal{Q}$ is any reductive algebra such that $\mathcal{Q}'$ contains T , then $\mathcal{Q}'$ contains T^* .

Rosenthal then asked if T has property (P) under each of the following conditions:

- (1) T is polynomially compact,
- (2) $1 - T^*T$ is in some C_p class,
- (3) $T^* - T$ is in some C_p class,
- (4) T is a part of a finite-multiplicity backward shift.

We will show that each of the above conditions implies (P), but we need a preliminary result (Lemma 2).

In [5], the following lemma is proved:

LEMMA 1. *Let C be a nonzero compact operator, and suppose that B is an operator such that every subspace that reduces both B and C and has dimension greater than 1 properly contains a nonzero subspace that reduces B and C . Then B and C have a common reducing eigenvector.*

The argument used to prove this lemma, with minor (and obvious) modifications will yield the following fact:

LEMMA 2. *Let B be an operator and C a nonzero compact operator. Suppose that*

- (1) *Every hyperinvariant subspace of B reduces B and C .*
- (2) *Every hyperinvariant subspace of B of dimension greater than 1 properly contains a nonzero hyperinvariant subspace of B .*

Then B and C have a common reducing eigenvector.

PROOF. See [5, p. 230].

We are now ready to answer Rosenthal's questions. It should be remarked that C. K. Fong [2] has proved part (1) of the following theorem.

THEOREM 4. *An operator T has property (P) under any one of the following conditions:*

- (1) *T is polynomially compact,*
- (2) *$T^* - T$ is in some C_p class,*
- (3) *$1 - T^*T$ is in some C_p class,*
- (4) *T is a part of some finite-multiplicity backward shift.*

PROOF. (4) follows from (3), because the multiplicity of the shift of which T is a part is the rank of $\sqrt{1 - T^*T}$. (See [3, p. 278].) If this rank is finite then so is the rank of $1 - T^*T$.

Next we remark that each of the conditions (1), (2), (3) is inherited by direct summands, and that each condition guarantees the existence of hyperinvariant subspaces [8, Corollaries 6.13, 6.15, 6.16] and [7, Theorem 1.1].

We will prove in detail that (2) implies that (P) holds; the proofs for (1) and (3) are analogous. Let $C = T^* - T$ and suppose that C is in some C_p class. We also suppose that $\mathcal{Q}$ is a reductive algebra and that $T \in \mathcal{Q}'$. Let $\mathfrak{M}_0$ be

the subspace of Theorem 2; note that $\mathfrak{N}_0$ reduces $\mathcal{Q}$, T , and C , and that $(T^*|\mathfrak{N}_0) \in (\mathcal{Q}|\mathfrak{N}_0)'$. Thus it suffices to consider the case where T is hyporeductive (by the remark following the Corollary to Theorem 3).

Since T is hyporeductive, the space $\mathfrak{N}_1$ spanned by all the eigenvectors of T reduces $\mathcal{Q}$, T , and C , and $(T^*|\mathfrak{N}_1) \in (\mathcal{Q}|\mathfrak{N}_1)'$ (the last statement follows by Lemma 5 of [1]); thus it suffices to consider the restriction of $\mathcal{Q}$, T , and C to $\mathfrak{N}_1^\perp$; i.e., we consider the case where T has no eigenvalues and is hyporeductive.

After these reductions suppose C is nonzero. Because T is hyporeductive, every hyperinvariant subspace of T reduces T and C . Further, suppose $\mathfrak{N}$ is a hyperinvariant subspace of T , of dimension greater than 1. Then $\mathfrak{N}$ reduces T and C , and $(T|\mathfrak{N})^* - (T|\mathfrak{N})$ lies in some C_p class. Thus there is a hyperinvariant subspace of T properly contained in $\mathfrak{N}$. It now follows from the assumption that C is nonzero, and from Lemma 2, that T has a reducing eigenvector; however, we reduced to the case where T has no eigenvectors. Thus it must be that $C = 0$, which means that $T^* = T$ and $T^* \in \mathcal{Q}'$. The proof is complete.

To show that (1) implies (P) let p be a polynomial such that $p(T) = C$ is compact and proceed as above. It is necessary to know that an algebraic hyporeductive operator is normal [1, Theorem 4].

To show that (3) implies (P) let $C = 1 - T^*T$ and proceed as above.

We remark that the proof of Theorem 4 also establishes the following fact :

COROLLARY. *If T is hyporeductive and any one of conditions (1) through (4) holds, then T is normal.*

REFERENCES

1. C. K. Fong, *On operators with reducing hyperinvariant subspaces* (preprint).
2. ———, *On commutants of reductive algebras*, Proc. Amer. Math. Soc. **63** (1977), 111–114.
3. P. R. Halmos, *A Hilbert space problem book*, Van Nostrand, Princeton, N.J., 1967.
4. T. B. Hoover, *Operator algebras with reducing invariant subspaces*, Pacific J. Math. **44** (1973), 173–179.
5. R. L. Moore, *Reductive operators that commute with a compact operator*, Michigan Math. J. **22** (1975), 229–233.
6. ———, *Hyperinvariant subspaces of reductive operators*, Proc. Amer. Math. Soc. **63** (1977), 91–94.
7. C. Pearcy, J. R. Ringrose and N. Salinas, *Remarks on the invariant subspace problem*, Michigan Math. J. **21** (1974), 163–166.
8. H. Radjavi and P. Rosenthal, *Invariant subspaces*, Springer-Verlag, Berlin and New York, 1973.
9. P. Rosenthal, *On commutants of reductive operator algebras*, Duke Math. J. **41** (1974), 829–834.

DEPARTMENT OF MATHEMATICS, BUCKNELL UNIVERSITY, LEWISBURG, PENNSYLVANIA 17837