A HELLY TYPE THEOREM ON THE SPHERE

MEIR KATCHALSKI

ABSTRACT. This paper establishes a Helly type theorem for convex sets in S_n , the n -dimensional unit sphere.

1. Introduction. Let S_n denote the n -dimensional unit sphere in Euclidean $(n + 1)$ -dimensional space, R^{n+1} .

A subset of S_n will be called *convex* if it is the intersection of S_n with a convex cone with apex 0 in R^{n+1} .

A subset of S_n will be called *strongly convex* if it is convex and does not contain antipodal points.

Thus a set consisting of two antipodal points is convex but not strongly convex.

For other definitions of convex sets in S_n and for a survey of Helly type theorems see [2]. For standard notation and terminology see [3].

M. J. C. Baker has proven in [1] the following

THEOREM. *For all positive integers n and t , if the intersection of each $n + 1$ members of a family of at least $n + 1 + 2t$ strongly convex sets in S_n is nonempty then the intersection of some $n + 1 + t$ members of the family is nonempty.*

The purpose of this paper is to generalize Baker's result by proving

THEOREM A. *Let n and t be positive integers and let $\mathcal{Q}$ be a family of $n + 1 + t$ convex sets in S_n .*

If every $n + 1$ members of $\mathcal{Q}$ have nonempty intersection then the intersection of some $n + 1 + [t/2]$ members of $\mathcal{Q}$ is nonempty.

Moreover, $n + 1 + [t/2]$ cannot be replaced by $n + [t/2] + 2$.

It is possible to improve Theorem A for strongly convex sets and $t = 2$ by establishing

THEOREM B. *Let n be an integer and let $\mathcal{Q}$ be a family of $n + 3$ strongly convex sets in S_n . Suppose that every $n + 1$ members of $\mathcal{Q}$ have nonempty intersection.*

If k is the number of different $n + 2$ membered subfamilies $\mathcal{B}$ of $\mathcal{Q}$ such that $\bigcap \mathcal{B} = \emptyset$ then $k \leq 2$.

Received by the editors October 14, 1976.

AMS (MOS) subject classifications (1970). Primary 52A20, 52A35.

Key words and phrases. Convex sets, n -dimensional unit sphere, Helly type theorems.

© American Mathematical Society 1977

The proof of Theorem A is based on

(1) Properties of certain families of convex cones called *nondegenerate families* (*N.D.F.'s*), discussed in [4] and [5].

(2) The existence, or the nonexistence of certain k -neighborly polytopes in R^n ; see [3].

All the necessary definitions and properties of N.D.F.'s and of neighborly polytopes are stated in §2.

A proof of Theorem A is given in §3.

The proof of Theorem B is similar to the proof of Theorem A and is therefore omitted.

ACKNOWLEDGEMENT. The author wishes to express his thanks to Professor Grünbaum for his help and advice.

2. Nondegenerate families and neighborly polytopes. Let $\mathcal{Q}$ be a nonempty finite family of convex cones with apex 0 in R^n .

The family $\mathcal{Q}$ will be called a *nondegenerate family* (a *N.D.F.*) if each member of the family is of dimension n , the intersection of any two members of the family is at least of dimension $(n - 1)$, . . . , the intersection of any n -members is at least of dimension 1 and the intersection of all members is $\{0\}$.

A subset $\mathfrak{B}$ of $\mathcal{Q}$ will be called a *face* of $\mathcal{Q}$ if $\bigcap (\mathcal{Q} \setminus \mathfrak{B})$ is a subspace.

A subset $\mathfrak{B}$ of $\mathcal{Q}$ is a k -face of $\mathcal{Q}$ if it is a face of $\mathcal{Q}$ and if $\dim \bigcap (\mathcal{Q} \setminus \mathfrak{B}) = |\mathfrak{B}| - k$. ($|\mathfrak{B}|$ means the cardinality of $\mathfrak{B}$.)

A relationship between N.D.F.'s and polytopes is given by

THEOREM C. (i) If $\mathcal{Q}$ is a N.D.F. in R^n and $|\mathcal{Q}| = n + t$ then there exists a $(t - 1)$ -polytope P in R^{t-1} such that

(1) The lattice of faces of P and the lattice of faces of $\mathcal{Q}$, both ordered by the inclusion relation, are isomorphic. The isomorphism carries k -faces of P to $(k + 1)$ -faces of $\mathcal{Q}$.

(ii) For any $(t - 1)$ -polytope P with $n + t$ vertices there exists a N.D.F. $\mathcal{Q}$ in R^n with $|\mathcal{Q}| = n + t$ such that (1) is satisfied.

A proof of Theorem C is given in [5].

NEIGHBORLY POLYTOPES. A polytope P is k -neighborly if every subset of k vertices of P is the set of vertices of a proper face of P .

Proofs of the following two theorems may be found in Chapter 7 of [3].

THEOREM D. If P is a k -neighborly d -polytope then:

(i) All k vertices of P are affinely independent and P is k' neighborly for $1 \leq k' \leq k$.

(ii) if $k > \lfloor \frac{1}{2} d \rfloor$ then P is a simplex.

THEOREM E. For every d and $v > d$ there exist d -polytopes with v vertices which are $\lfloor \frac{1}{2} d \rfloor$ neighborly.

3. Proof of Theorem A. In order to prove Theorem A it is sufficient to prove

THEOREM F. *Let n and t be nonnegative integers. Let $\mathcal{Q}$ be a family of $n + t$ convex cones with apex 0 in R^n such that*

$$(2) \quad \bigcap \mathcal{Q} = \{0\}$$

and

$$(3) \quad \bigcap \mathcal{B} \neq \{0\} \quad \text{for any } n \text{ membered subfamily } \mathcal{B} \text{ of } \mathcal{Q}.$$

Then there exists an $n + [t/2]$ membered subfamily $\mathcal{C}$ of $\mathcal{Q}$ such that $\bigcap \mathcal{C} \neq \{0\}$.

Moreover, $n + [t/2]$ in the last statement cannot be replaced by $n + [t/2] + 1$.

PROOF OF THEOREM F. The proof is by induction on n . The cases $n = 0$ and $t \leq 1$ are trivial, so assume that $n > 0$ and that $t \geq 2$.

Let $\mathcal{Q}$ be a family of convex cones with apex 0 on R^n such that $|\mathcal{Q}| = n + t$ and assume that (2) and (3) are satisfied.

Suppose that A is not a N.D.F. let $\mathcal{N}$ be a maximal subset of $\mathcal{Q}$ such that

$$(4) \quad n' = \dim \bigcap \mathcal{N} \leq n - |\mathcal{N}|.$$

Clearly ϕ satisfies (4) since $\dim \bigcap \phi = n = n - |\phi|$.

It is not difficult to verify that $\dim \bigcap \mathcal{N} = n - |\mathcal{N}|$ and that $\mathcal{N} \neq \phi$ since $\mathcal{Q}$ is not a N.D.F.

Define a family $\mathcal{Q}' = \{\bigcap \mathcal{N} \cap \{A\} : A \in \mathcal{Q} \setminus \mathcal{N}\}$ of convex cones with apex 0 in $R^{n'} = \text{span } \bigcap \mathcal{N}$.

Since $|\mathcal{Q}'| = n + t - |\mathcal{N}| = n' + t$ and since $\mathcal{Q}'$ satisfies (2) and (3) we can use the induction hypothesis to obtain a $\mathcal{C}' \subset \mathcal{Q} \setminus \mathcal{N}$ such that $|\mathcal{C}'| \geq n' + [t/2]$ and $\bigcap \mathcal{C}' \neq \{0\}$.

Define $\mathcal{C} = \mathcal{C}' \cup \mathcal{N}$. The subfamily $\mathcal{C}$ has the desired properties since

$$|\mathcal{C}| = |\mathcal{C}'| + |\mathcal{N}| \geq n' + [t/2] + |\mathcal{N}| = n + [t/2]$$

and

$$\bigcap \mathcal{C} = \bigcap \mathcal{C}' \neq \{0\}.$$

Suppose that A is a N.D.F. Assume by contradiction that $\bigcap \mathcal{B} = \{0\}$ for any $n + [t/2]$ membered subfamily $\mathcal{B}$ of $\mathcal{Q}$. Thus for any $\mathcal{C} \subset \mathcal{Q}$ such that

$$|\mathcal{C}| \leq n + t - (n + [t/2]) = t - [t/2],$$

$\mathcal{C}$ is a $|\mathcal{C}|$ -face of $\mathcal{Q}$.

Let P be the $(t - 1)$ -polytope described in Theorem C. It follows from (1) of Theorem C that P is a $t - [t/2]$ neighborholy polytope with $n + t$ vertices. Since $t - [t/2] > [(t - 1)/2]$, P is by Theorem D a $(t - 1)$ -simplex and therefore $n + t = (t - 1) + 1$, so that $n = 0$, a contradiction. This completes the proof of the first part of Theorem F.

To complete the proof of Theorem F let P be a $[(t - 1)/2]$ neighborly $(t - 1)$ -polytope with $n + t$ vertices (see Theorem E).

Let $\mathcal{Q}$ be the N.D.F. described in (ii) of Theorem C. The family satisfies (2) and (3) since any N.D.F. in R^n satisfies (2) and also (3) if $n > 0$.

For any $\mathcal{B} \subset \mathcal{Q}$ such that $\mathcal{B} \geq n + [t/2] + 1$ we have that

$$|\mathcal{Q} \setminus \mathcal{B}| \leq t - [t/2] - 1 \leq [(t-1)/2].$$

By Theorem C(i), for any such $\mathcal{B}$, $\mathcal{Q} \setminus \mathcal{B}$ is a $|\mathcal{Q} \setminus \mathcal{B}|$ -face of $\mathcal{Q}$.

Therefore $\dim \cap \mathcal{B} = \dim \cap (\mathcal{Q} \setminus (\mathcal{Q} \setminus \mathcal{B})) = |\mathcal{Q} \setminus \mathcal{B}| - |\mathcal{Q} \setminus \mathcal{B}| = 0$ so that $\cap \mathcal{B} = \{0\}$. The proof of Theorem F is now complete.

REFERENCES

1. M. J. C. Baker, *A spherical Helly type theorem*, Pacific J. Math. **23** (1967), 1-3. MR 36 #5820.
2. L. Danzer, B. Grünbaum and V. Klee, *Helly's theorem and its relatives*, Proc. Sympos. Pure Math., vol. 7, Amer. Math. Soc., Providence, R. I., 1963, pp. 101-180. MR 28 #524.
3. B. Grünbaum, *Convex polytopes*, Interscience, New York, 1967. MR 37 #2085.
4. M. Katchalski, *Some properties of families of convex cones*, Trans. Amer. Math. Soc. (to appear).
5. ———, *Non-degenerate families of convex cones and convex polytopes*, Discrete Math. (to appear).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA T6G 2G1, CANADA