STICKY ARCS IN E^n ($n \geq 4$)

DAVID G. WRIGHT

ABSTRACT. Let A and B be arcs in E^3 , Euclidean 3-space. Then A can be "slipped" off B ; i.e., there exists a homeomorphism of E^3 onto itself, arbitrarily close to the identity, such that $h(A) \cap B = \emptyset$. The purpose of this note is to show that arcs in E^n ($n \geq 4$) do not always enjoy this property. The examples depend heavily on a recent result of McMillan.

1. Introduction. If X and Y are subsets of E^n , Euclidean n -space, we say that X can be *slipped* off Y in E^n if for each $\varepsilon > 0$ there is an ε -homeomorphism $h: E^n \rightarrow E^n$ such that $h(X) \cap Y = \emptyset$; otherwise, we say X *cannot be slipped off* Y . Results of Armentrout [1] and McMillan [5] show that if A and B are arcs in E^3 , then A can be slipped off B . We show that this is false in higher dimensions by proving the following

THEOREM. *There exist cellular arcs A and B in E^n ($n \geq 4$) such that A cannot be slipped off B .*

Existence of A and B . McMillan [4] has shown, for each $n \geq 4$, there exists an arc C in an n -manifold M^n which has no neighborhood in M^n which embeds in E^n . Without loss of generality, we may assume that C is the union of arcs A' and B' whose intersection is a common endpoint and such that A' and B' have neighborhoods U and V , respectively, which embed in E^n . Let $f: U \rightarrow E^n$ and $g: V \rightarrow E^n$ be embeddings. By [2] and a correct choice of U and V , we may assume that f and g agree on $U \cap V$. We now let $A = f(A')$ and $B = g(B')$.

PROOF OF THEOREM. Let B_1 and B_2 be concentric n -balls centered at $f(A' \cap B')$ such that $B_1 \subset \text{Int } B_2 \subset f(U \cap V) = g(U \cap V)$. We suppose that A can be slipped off B and choose a homeomorphism $h: E^n \rightarrow E^n$ such that $h(A) \cap B = \emptyset$. Furthermore, h is chosen so close to the identity that the following conditions are satisfied.

- (1) $h(\text{Bd } B_2) \cap B_1 = \emptyset$.
- (2) There is an embedding $e: B_2 - \text{Int } B_1 \rightarrow E^n$ such that $e|_{\text{Bd } B_1} = \text{identity}$ and $e|_{\text{Bd } B_2} = h|_{\text{Bd } B_2}$.
- (3) The embedding e is so close to the identity that $e(A \cap (B_2 - \text{Int } B_1)) \cap B = \emptyset$.

The existence of the embedding e is a corollary of the Kirby-Edwards local contractibility theorems [3] and is used explicitly in [6].

Received by the editors April 4, 1977 and, in revised form, May 30, 1977.

AMS (MOS) subject classifications (1970). Primary 57A15; Secondary 57A35, 57A60.

© American Mathematical Society 1977

Consider the homeomorphism $H: E^n \rightarrow E^n$ given by $H|_{B_1} = \text{identity}$, $H|(B_2 - \text{Int } B_1) = e$, and $H|(E^n - B_2) = h|(E^n - B_2)$. Thus, $H \circ f: U \rightarrow E^n$ and $g: V \rightarrow E^n$ are embeddings which agree on a neighborhood of $A' \cap B'$ in M^n and such that $H \circ f(A') \cap g(B')$ is a single point. By choosing even smaller neighborhoods U' and V' of A' and B' , respectively, we may assume that $f' = H \circ f|_{U'}$ and $g' = g|_{V'}$ agree on $U' \cap V'$ and that $f'(U') \cap g'(V') = f'(U' \cap V') = g'(U' \cap V')$. Thus, f' and g' can be used to give an embedding of $U' \cup V'$ into E^n which yields a contradiction.

An examination of McMillan's arc C shows that the interior of C has a neighborhood in M^n which is homeomorphic with the suspension of a set X , ΣX , the homeomorphism sending C to $\Sigma\{x\}$, $x \in X$. The set X is obtained from a piecewise-linear $(n - 1)$ -manifold by identifying a cell-like set in its interior to the point x . This neighborhood may be used to show that any proper subarc of C can be pushed arbitrarily close to one of its endpoints by a homeomorphism of M^n which has support in an arbitrary neighborhood in M^n of the subarc. This fact implies that any subarc of C is cellular. Therefore we may assume that A and B are cellular.

REFERENCES

1. S. Armentrout, *Decompositions of E^3 with a compact 0-dimensional set of nondegenerate elements*, Trans. Amer. Math. Soc. **123** (1966), 165–177.
2. M. Brown, *A proof of the generalized Schoenflies theorem*, Bull. Amer. Math. Soc. **66** (1960), 74–76.
3. R. D. Edwards and R. C. Kirby, *Deformations of spaces of imbeddings*, Ann. of Math. (2) **93** (1971), 63–88.
4. D. R. McMillan, Jr., *An arc in a PL n -manifold with no neighborhood that embeds in S^n , $n > 4$* , Michigan Math. J. (to appear).
5. ———, *A criterion for cellularity in a manifold. II*, Trans. Amer. Math. Soc. **126** (1967), 217–224.
6. C. L. Seebeck, III, *Collaring an $(n - 1)$ -manifold in an n -manifold*, Trans. Amer. Math. Soc. **148** (1970), 63–68.

DEPARTMENT OF MATHEMATICS, UTAH STATE UNIVERSITY, LOGAN, UTAH 84322