A TYPE OF NEAREST POINT SET IN A COMPLETE l -GROUP¹

MICHAEL KEISLER

ABSTRACT. A theorem by W. D. L. Appling (Riv. Mat. Univ. Parma, (3) 2 (1973), 251–276) demonstrates that a C -set is a nearest point set in $\text{ba}(S, \Sigma)$ with respect to the variation norm. This paper demonstrates an analogous result for a generalized form of C -set in a complete l -group with distance with respect to the norm being replaced by a stronger property definable in a complete l -group (distance between elements x and y of a complete l -group G is taken to be $|x - y|$). The result is then shown to be a characterization of sets possessing the stronger property in the case of a complete vector lattice, but not a characterization in the case of a complete l -group.

1. Introduction. There are obvious analogies between a norm on a space and the absolute value of elements of an l -group. Using $|x - y|$ as our concept of “distance”, we will characterize the “nearest point” sets in a complete vector lattice, and demonstrate that the characterization fails in the case of a complete l -group. It is noted in [3, pp. 366, 371] that in a uniformly monotone Banach lattice (UMB-lattice) $|||x||| = \|x\|$, and if $0 \leq x < y$, then $\|x\| < \|y\|$. Therefore, in a UMB-lattice, every nearest point set of the type described above is a nearest point set with respect to the norm. The converse is not true, however, as a closed circular disc in the plane with $\|(x, y)\| = |x| + |y|$ readily demonstrates.

The general properties of l -groups, as presented in [3], are assumed throughout. Necessary properties not specifically proven in [3] are presented in the lemma below.

LEMMA 1. *Let G be a complete l -group and $\{a, b, c\} \subseteq G$,*

- (i) *if $|a| \wedge |b| = 0$, then $|a + b| = |a| + |b|$,*
- (ii) *if $a \wedge b \wedge c \geq 0$, then $(a + b) \wedge c \leq a \wedge c + b \wedge c$,*
- (iii) *$||a| - |b|| \leq |a - b| \wedge |a + b|$.*

PROOF. The triangle inequality implies $|a| \leq |a - b| + |b|$ and $|b| \leq |a -$

Presented to the Society, January 23, 1975 under the title *A characterization of a type of nearest point set in complete l -groups*; received by the editors June 18, 1976.

AMS (MOS) subject classifications (1970). Primary 06A60, 06A65, 26A51, 46A40; Secondary 28A10, 47B55.

Key words and phrases. Complete l -group, complete vector lattice, complex, l -nearest point set, normal subgroup, projection operator.

¹This article contains, in part, research undertaken to fulfill requirements for the Ph. D. at North Texas State University, Denton, Texas, in 1974.

© American Mathematical Society 1978

$b| + |a|$, from which it follows that

$$||a| - |b|| = (|a| - |b|) \vee (|b| - |a|) \leq |a - b|.$$

Since $|b| = |-b|$, we have (iii). (ii) follows easily from the inequality $(a + b) \wedge c \leq (a + b) \wedge (c + b) \wedge (a + c) \wedge (c + c)$. Finally, (i) can be proven by applying (iii);

$$\begin{aligned} ||a| - |b|| &\leq |a + b| \leq |a| + |b| = |a| \vee |b| - |a| \wedge |b| \\ &= (|a| - |b|) \vee 0 + |b| + (-|a|) \vee (-|b|) \\ &= (|a| - |b|) \vee 0 + (|b| - |a|) \vee 0 = ||a| - |b||. \end{aligned}$$

DEFINITION. Let G be an l -group. $S \subseteq G$ is an l -nearest point set (l -nps) iff for $x \in G$, there is $x' \in S$ such that if $s \in S$ and $s \neq x'$, then $|x - x'| < |x - s|$. (For an l -nps S , denote the nearest point map from G to S by f_S ; i.e., for x in G , the nearest point in S is $f_S(x)$.)

Some basic properties of l -nearest point sets are given by the lemma below. Although the main results of this paper deal with complete l -groups, this lemma takes G to be an arbitrary l -group.

LEMMA 2. Let G be an l -group with l -nps S . For $x \in G$, $S - x$ is an l -nps (with $f_{S-x}(z) = f_S(z + x) - x$, for $z \in G$), $-S$ is an l -nps if G is abelian (with $f_{-S}(z) = -f_S(-z)$, for z in G), and if $S' \subseteq S$ such that $x = \sup S'$ exists, then $x \in S$.

PROOF. For $z \in G$,

$$|z - [f_S(z + x) - x]| = |z + x - f_S(z + x)|,$$

and the latter is $< |z + x - s| = |z - (s - x)|$, for $s - x$ in $S - x$, unless $s - x = f_S(z + x) - x$; hence $S - x$ is an l -nps. For $z \in G$, $|z - (-f_S(-z))| = |(-z) - f_S(-z)|$, and the latter is less than $|(-z) - s| = |z - (-s)|$, for $-s \in -S$, unless $-s = -f_S(-z)$; $-S$ is an l -nps. Finally, given $x = \sup S'$,

$$\begin{aligned} |x - f_S(x)| &\leq \inf\{|x - s| | s \in S'\} = \inf\{x - s | s \in S'\} \\ &= x - \sup S'; \end{aligned}$$

hence $|x - f_S(x)| = 0$ and $x = f_S(x) \in S$.

2. Some l -nearest point sets. The study of projection operators has a very long history. Bochner and Phillips noted in [4] that the function $p_Y(x) = \sup\{x \wedge y | y \in Y\}$, for $x \geq 0$ and Y a normal subspace of a complete vector lattice, provided a decomposition (Lebesgue decomposition for $\text{ba}(S, \Sigma)$, where Y is a principal normal subspace) of an element x into $x - p_Y(x)$ and $p_Y(x)$ such that $|x - p_Y(x)| \wedge |p_Y(x)| = 0$, where p_Y has been extended to the entire vector lattice as follows; $p_Y(x) = p_Y(x^+) - p_Y(x^-)$, for x a member of the vector lattice, with notation as in [4]. Kakutani also studied principal projection operators in [5], using the form $p_Y(x) = \sup\{x \wedge ny | n = 1, 2, 3, \dots\}$. Applying Lemma 1, we have that

$$|x - p_Y(z)| = |x - p_Y(x)| + |p_Y(x - z)| > |x - p_Y(x)|,$$

if $p_Y(z) \neq p_Y(x)$, since projection operators are additive and the singularity property mentioned above applies to $x - p_Y(x)$ and any member of Y . Hence every normal subspace is an l -nps. Appling [1] observed that the Lebesgue decomposition, using Kakutani's representation for the projection operator, provided a proof of a nearest point property in $\text{ba}(S, \Sigma)$ with respect to the variation norm. As noted in the introduction, every l -nps in a UMB-lattice, which $\text{ba}(S, \Sigma)$ with variation norm is, is a nearest point set, so the observation about normal subspaces above implies Appling's result.

Pinsker studied projection operators as well in [7] and [8] (the material contained herein is taken from a summary published later in English [6]). He established the properties of projection operators mentioned above for complete l -groups, but more to the point of this paper, he examined subsets of the positive cone called complexes.

DEFINITION. Let G be a complete l -group with positive cone P . A subset A of P is a *complex* provided (1) $x \in A$ implies $[0, x] \subseteq A$ (where $[0, x] = \{y | 0 \leq y \leq x\}$), (2) $E \subseteq A$ and E bounded implies $\sup E \in A$, and (3) for any $x > 0$, there is a positive integer n such that $nx \notin A$.

Appling studied subsets of $\text{ba}(S, \Sigma)$ which were, as we shall show, direct sums of sets of the form $A + (-A)$, where A is a complex, and a normal subspace. He termed the subsets *C-sets* (see [2]); their definition is essentially the same as that of Pinsker for a complex except that (1) becomes (1'), $x \in A$ implies $[-|x|, |x|] \subseteq A$, and (3) is omitted. The approach taken below owes much to that taken by Appling.

Henceforth G is assumed to be a complete l -group.

DEFINITION. $I = \{\alpha | \alpha: P \rightarrow P, \alpha \leq \geq, \alpha \text{ is isotone, idempotent and subadditive}\}$.

LEMMA 3. If $\alpha \in I$ and $\{x, y\} \subseteq P$, then $y \leq \alpha(x)$ implies $y = \alpha(y)$.

PROOF. $\alpha(x) = \alpha(\alpha(x)) \leq \alpha(\alpha(x) - y) + \alpha(y) \leq \alpha(x) - y + \alpha(y)$.

LEMMA 4. If $x, y \in P$ and $\alpha \in I$, then $x \wedge \alpha(y) = \alpha(x \wedge y) = \alpha(x) \wedge y$.

PROOF. Since α is isotone and a subset of $\geq$, $\alpha(x \wedge y) \leq x \wedge \alpha(y) \leq x \wedge y$. Hence $\alpha^2 = \alpha$ implies $\alpha(x \wedge y) \leq \alpha(\alpha(x \wedge y)) \leq \alpha(x \wedge y)$. By Lemma 3, $\alpha(x \wedge \alpha(y)) = x \wedge \alpha(y)$ and thus $\alpha(x \wedge y) = x \wedge \alpha(y)$. Similarly for $\alpha(x) \wedge y$.

LEMMA 5. $\alpha \in I$ iff $\alpha(x) = \sup\{x \wedge y | y \in S\}$, for some $S \subseteq P$.

PROOF. By Lemma 4, if $\alpha \in I$, then $\alpha(x \wedge \alpha(z)) = x \wedge \alpha(z)$, for $x, z \in P$. Thus

$$\alpha(x) \geq \sup\{x \wedge \alpha(z) | z \in P\} \geq x \wedge \alpha(x) = \alpha(x),$$

and the theorem is proven for $S = \alpha(P)$. If $S \subseteq P$ and $\alpha(x) = \sup\{x \wedge y | y \in S\}$, then clearly $\alpha \leq \geq$ and α is isotone. And

$$\begin{aligned}\alpha(\alpha(x)) &= \sup\{\alpha(x) \wedge y \mid y \in S\} \\ &\geq \sup\{(x \wedge y) \wedge y \mid y \in S\} = \alpha(x) \geq \alpha(\alpha(x))\end{aligned}$$

implies $\alpha^2 = \alpha$. Finally, Lemma 1(ii) implies $\alpha(x + y) \leq \alpha(x) + \alpha(y)$.

The last lemma allows us to note that both the notion of complex (by defining $\alpha_A(x) = \sup\{x \wedge a \mid a \in A\}$, for each complex A) and projection operator is included in the class I . If we define $(\alpha + \beta)(x) = \alpha(x) + \beta(x - \alpha(x))$, $(\alpha \wedge \beta)(x) = \alpha(x) \wedge \beta(x)$ and $(\alpha \vee \beta)(x) = \alpha(x) \vee \beta(x)$, for $x \in P$, and $\alpha \leq \beta$ iff $\alpha(x) \leq \beta(x)$, for $x \in P$, then $+$, $\wedge$ and $\vee$ are binary operations on I and $\leq$ is a partial order on I compatible with $+$. Furthermore, $0(x) = 0$, and $1(x) = x$, for $x \in P$, are lower and upper bounds in I .

If we use the notation $[-\alpha, \beta] = -\alpha(P) + \beta(P)$ (which, in turn, $= \{x \mid x^+ = \beta(x^+)$ and $x^- = \alpha(x^-)\}$), then we have a convenient way of describing a large class of l -nearest point sets. Intervals of the form $[-\alpha, \alpha]$, for $\alpha \in I$, are C -sets. Appling [2] showed that such intervals in $\text{ba}(S, \Sigma)$ form nearest point sets with respect to the variation norm. The result below extends Appling's to l -nearest point sets in a complete l -group.

THEOREM 1. *Let G be a complete l -group. If $\alpha, \beta \in I$, then $[-\alpha, \beta]$ is an l -nps, and hence $x + [-\alpha, \beta]$ is an l -nps, for $x \in G$.*

PROOF. For $x \in G$, let $f(x) = \beta(x^+) - \alpha(x^-)$, where $x^+ = x \vee 0$ and $x^- = (-x) \vee 0$. Then for $x \in G$ and $z \in [-\alpha, \beta]$, such that $z \neq f(x)$, we know $\alpha(x^-) \neq z^-$ or $\beta(x^+) \neq z^+$, from which it follows that

$$\begin{aligned}|x - f(x)| &= |(x^+ - \beta(x^+)) - (x^- - \alpha(x^-))| \\ &= x^+ - \beta(x^+) + x^- - \alpha(x^-) \\ &< x^+ - x^+ \wedge z^+ + z^+ - x^+ \wedge z^+ + x^- - x^- \wedge z^- + z^- - x^- \wedge z^- \\ &= |(x^+ - x^+ \wedge z^+ + z^+ - x^- \wedge z^-) \\ &\quad - (x^- - x^- \wedge z^- + z^- - x^+ \wedge z^+)| = |x - z|,\end{aligned}$$

by noting that

$$\begin{aligned}0 &= x^+ \wedge x^- = z^+ \wedge z^- \\ &= (x^+ - x^+ \wedge z^+) \wedge (z^+ - x^+ \wedge z^+) \\ &= (x^- - x^- \wedge z^-) \wedge (z^- - x^- \wedge z^-),\end{aligned}$$

and applying (i) and (ii) of Lemma 1.

For $x \in P$, let $\tau_x(z) = z \wedge x$, for $z \in P$, then $\tau_x \in I$. If $x, y \in P$, then $[-\tau_x, \tau_y] = [-x, y]$, using the usual interval notation for a partially ordered set, and we have the following.

COROLLARY. *Let G be a complete l -group. If $x, y \in G$, and $x \leq y$, then $[x, y]$ is an l -nps.*

Not all l -nearest point sets in a complete l -group are necessarily of the form given in Theorem 1. For instance, the set of integers forms a complete

l -group, and $\{0, 3\}$ is an l -nps. Since $0 < 2 < 3$, and $2 \notin \{0, 3\}$, we can conclude that we have an l -nps other than the type described above. In §4 we shall show that the description of Theorem 1 does characterize l -nearest point sets in complete vector lattices. First we will investigate some further properties of members of I .

3. Decompositions of members of I . It was pointed out in the last section that C -sets could be written as the sum of a set of the form $A + (-A)$ and a normal subspace. This corresponds to demonstrating that each element of I can be decomposed into a function whose image is a complex, and an additive member of I , because it is clear from Lemma 5 that if A is a C -set, then $A \cap P$ defines a member of I , α , whose image is $A \cap P$. Since $A = A \cap P + (-A) \cap P$, by applying (1'), we have $A = [-\alpha, \alpha]$. The decomposition is achieved below.

THEOREM 2. *If $\alpha \in I$, then there is $\alpha', \alpha^* \in I$ such that $\alpha' \wedge \alpha^* = 0$, α' is additive, $\alpha^*(P)$ is a complex and $\alpha = \alpha^* + \alpha'$.*

PROOF. Define $\alpha'(x) = \sup\{\mu(x) \mid \mu \in I, \mu \text{ is additive and } \mu \leq \alpha\}$, for $x \in P$. It is clear that $\alpha': P \rightarrow P$, $\alpha' \subseteq \geq$, α' is isotone and α' is idempotent (the latter since $\mu(\alpha'(x)) = \mu(x) \wedge \alpha'(x) = \mu(x)$, for μ such that $\mu(x) \leq \alpha'(x)$, by Lemma 4). A member γ of I is additive iff $\gamma(P) + (-\gamma(P))$ is a normal subgroup of G ; i.e., iff γ is a projection operator. Referring to the property $(x - \gamma(x)) \wedge \gamma(x) = 0$, for $x \in P$, then if each of μ and β are additive and $\leq \alpha$, then

$$\begin{aligned} (\mu + \beta)(x) &= \mu(x) + \beta(x - \mu(x)) = \mu(x) \vee \beta(x - \mu(x)) \\ &\leq \alpha(x) \vee \alpha(x - \mu(x)) = \alpha(x). \end{aligned}$$

The latter inequality implies α' is additive since it yields $\alpha'(x) + \alpha'(y) \leq \alpha'(x + y)$. Thus α' is an additive member of I . Let $\alpha^*(x) = \alpha(x) - \alpha'(x)$, for $x \in P$. If $0 \leq x \leq y$, then

$$\begin{aligned} \alpha'(y) &= \alpha'(\alpha(y)) = \alpha'(\alpha(y) - \alpha(x)) + \alpha'(\alpha(x)) \\ &\leq \alpha(y) - \alpha(x) + \alpha'(x), \end{aligned}$$

which implies α^* is isotone. Since the other properties of members of I are clearly possessed by α^* , then α^* is in I . For $x \in P$,

$$\begin{aligned} (\alpha^* \wedge \alpha')(x) &= (\alpha(x) - \alpha'(x)) \wedge \alpha'(x) \\ &\leq (x - \alpha'(x)) \wedge \alpha'(x) = 0, \end{aligned}$$

and we have $\alpha^* \wedge \alpha' = 0$. If $x > 0$, then $nx \in \alpha^*(P)$ for every positive integer n implies $p_x \leq \alpha^* \leq \alpha$, which leads, by definition of α' , to $p_x \leq \alpha'$. The last statement is a contradiction. Therefore, for $x > 0$, there is a positive integer n such that $nx \notin \alpha^*(P)$; i.e., $\alpha^*(P)$ is a complex.

For $\alpha, \beta \in I$ and $x, y \in P$,

$$\beta(y) \geq \beta(y) - (\alpha(\alpha(x) + \beta(y)) - \alpha(x)),$$

and, by Lemma 3,

$$\begin{aligned} (\alpha + \beta)(\alpha(x) + \beta(y)) &= \alpha(\alpha(x) + \beta(y)) \\ &\quad + (\alpha(x) + \beta(y) - \alpha(\alpha(x) + \beta(y))) \\ &= \alpha(x) + \beta(y), \end{aligned}$$

from which it follows that $\alpha(P) + \beta(P) \subseteq (\alpha + \beta)(P)$. The reverse containment is immediate, and we have $(\alpha + \beta)(P) = \alpha(P) + \beta(P)$. Pinsker [6, pp. 65–67] proved that the collection of complexes with set addition as the operation is isomorphic to the positive cone of a complete l -group, and hence one in which P can be embedded; $x \rightarrow \tau_x(P)$ is easily seen to be a morphism (for $z \in P$,

$$\begin{aligned} (\tau_x + \tau_y)(z) &= \tau_x(z) + \tau_y(z - \tau_x(z)) \\ &= x \wedge z + y \wedge (z - x \wedge z) = (x \wedge z + y) \wedge z \\ &= [(x + y) \wedge (z + y)] \wedge z \\ &= (x + y) \wedge z = \tau_{x+y}(z). \end{aligned}$$

The latter will be useful in Theorem 3.

This theorem demonstrates that each complex can be decomposed as the direct sum of a collection of disjoint intervals of the form $[0, a]$. The decomposition suggests a different description of the positive cone P^* of G^* as a collection of equivalence classes containing disjoint subsets of P .

THEOREM 3. *Let $\alpha \in I$ such that $\alpha(P)$ is a complex. There is a disjoint subset A of P such that $\alpha(x) = \sup\{x \wedge a \mid a \in A\}$, for $x \in P$.*

PROOF. Define $\beta(x) = \sup\{p_{z-\alpha(z)}(x) \mid z \in P\}$, for $x \in P$. As in the proof of Theorem 2, β is an additive member of I .

$$(\alpha + \beta)(x) = \alpha(x) + \beta(x - \alpha(x)) = \alpha(x) + x - \alpha(x) = x,$$

for $x \in P$. Since $x > 0$ implies there is a positive integer n such that $nx \neq \alpha(nx)$, then the above implies

$$0 < \beta(nx - \alpha(nx)) \leq \beta(nx) = n\beta(x).$$

Thus, $x > 0$ leads to $\beta(x) > 0$. Since $\beta(x - \beta(x)) = 0$, for $x \in P$, we have $\beta(x) = x$, for $x \in P$. A straightforward application of Kuratowski's Lemma will provide a disjoint subset E of P with each e in $E \leq x_e - \alpha(x_e)$, for some $x_e \in P$, and $\sup\{p_e(x) \mid e \in E\} = x$, for $x \in P$. Let $\bar{e} = x_e - \alpha(x_e)$, for $e \in E$.

$$\begin{aligned} (\alpha \wedge \tau_{\bar{e}})(z) &= \alpha(z) \wedge \tau_{\bar{e}}(z) = \alpha(z) \wedge \bar{e} \\ &= z \wedge \alpha(\bar{e}) = \tau_{\alpha(\bar{e})}(z) \leq \tau_{\alpha(x_e)}(z), \end{aligned}$$

for $z \in P$, by application of Lemma 4. If n is a positive integer such that $\alpha \wedge n\tau_{\bar{e}} \leq \tau_{\alpha(x_e)}$, where $n\tau_{\bar{e}}$ is defined to be $\tau_{\bar{e}} + \tau_{\bar{e}} + \cdots + \tau_{\bar{e}}$, then

$$\begin{aligned}\alpha \wedge (n+1)\tau_{\bar{e}} &\leq \alpha \wedge (\alpha \wedge \tau_{\bar{e}} + \alpha \wedge n\tau_{\bar{e}}) \\ &\leq \alpha \wedge (\tau_{\bar{e}} + \tau_{\alpha(x_e)}) = \alpha \wedge \tau_{(\bar{e} + \alpha(x_e))} = \tau_{\alpha(x_e)},\end{aligned}$$

and by induction we have $\alpha \wedge \tau_{n\bar{e}} = \alpha \wedge n\tau_{\bar{e}} \leq \tau_{\alpha(x_e)}$, for every positive integer n , from which it follows that $\alpha \wedge p_{\bar{e}} \leq \tau_{\alpha(x_e)}$. Let $a_e = (\alpha \wedge p_e)(x_e)$, for $e \in E$; then $A = \{a_e | e \in E\}$ forms a disjoint subset of P such that

$$\begin{aligned}\sup\{x \wedge a | a \in A\} &= \sup\{x \wedge (\alpha \wedge p_e)(x_e) | e \in E\} \\ &= \sup\{(\alpha \wedge p_e)(x) \wedge x_e | e \in E\} = \sup\{p_e(x) \wedge \alpha(x) \wedge \alpha(x_e) | e \in E\} \\ &= \sup\{p_e(x) \wedge \alpha(x) | e \in E\} = \sup\{p_e(\alpha(x)) | e \in E\} = \alpha(x).\end{aligned}$$

4. l -nearest point sets in a complete vector lattice. In this section we shall return to a study of l -nearest point sets. In order to arrive at the theorem characterizing l -nearest point sets, we must first prove technical lemmas about such sets. Again G is taken to be a complete l -group.

LEMMA 6. *Let S be an l -nps; then for $x \in G$, $f_S(x \vee f_S(x)) = f_S(x)$.*

PROOF.

$$\begin{aligned}|x \vee f_S(x) - f_S(x)| &= (x - f_S(x)) \vee 0 \\ &= |x - f_S(x)| + (x - f_S(x)) \wedge 0 \\ &\leq |x - f_S(x \vee f_S(x))| - (f_S(x) - x) \vee 0 \\ &= ||x - f_S(x \vee f_S(x))| - |(f_S(x) - x) \vee 0|| \\ &\leq |x - f_S(x \vee f_S(x)) + (f_S(x) - x) \vee 0| \\ &= |x \vee f_S(x) - f_S(x \vee f_S(x))|,\end{aligned}$$

by applying properties of complete l -groups along with Lemma 1(iii).

LEMMA 7. *Let S be an l -nps, $\{0, t\} \subseteq S \cap P$ and $m \in [0, t]$ such that $m \wedge (t - m) = 0$; then $m \in S$.*

PROOF. $|m - f_S(m)| \leq |m - 0| \wedge |m - t| = m \wedge (t - m) = 0$.

In the remainder of the paper G will be taken to be a complete vector lattice (see [3] for definitions). The main difficulty in demonstrating the converse of Theorem 1 is, as the example of §2 suggests, in establishing the convexity of an l -nps. The lemma below provides an essential tool for doing so.

LEMMA 8. *If S is an l -nps, $\{0, t\} \subseteq S \cap P$ and $x \in [0, t]$ such that $f_S(x) = 0$, then $x = 0$.*

PROOF. Since $x = |x - f_S(x)| < |x - t| = t - x$, we have

$$T = \{y | 2x \leq y \text{ and } y \in S \cap [0, t]\} \neq \Phi,$$

and thus $w = \inf T$ exists and is in $S \cap [0, t]$.

$$w/2 - f_S(w/2) \leq |w/2 - f_S(w/2)| \leq |w/2 - 0| = w/2,$$

implies $f_S(w/2) \geq 0$. Let $r = (x - f_S(w/2))^+$, and note that $p_r(f_S(w/2)) \in S$ by Lemma 7. Since $r \wedge (x - f_S(w/2))^- = 0$,

$$\begin{aligned} 0 &= p_r(f_S(w/2)) \wedge ((f_S(w/2) - x) \vee 0) \\ &= [p_r(f_S(w/2)) \wedge (f_S(w/2) - x)] \vee [p_r(f_S(w/2)) \wedge 0] \\ &= [(p_r(f_S(w/2)) + x) \wedge f_S(w/2)] - x \vee 0, \end{aligned}$$

from which it follows that

$$p_r(f_S(w/2)) \leq (p_r(f_S(w/2)) + x) \wedge f_S(w/2) \leq x.$$

Thus

$$|x - p_r(f_S(w/2))| \leq x = |x - f_S(x)|,$$

and we have $p_r(f_S(w/2)) = 0$. Since

$$0 \leq p_r(w) \wedge f_S(w/2) = w \wedge p_r(f_S(w/2)) = 0,$$

by Lemma 4, then we can apply Lemma 1(i), (ii), and note that

$$\begin{aligned} |w/2 - p_r(w) - f_S(w/2)| &= |(w/2 - p_r(w/2) - f_S(w/2)) - p_r(w/2)| \\ &= |w/2 - p_r(w/2) - f_S(w/2) + p_r(w/2)| = |w/2 - f_S(w/2)|. \end{aligned}$$

The fact that $p_r(w) + f_S(w/2) = p_r(w) \vee f_S(w/2) \in S$ implies that $f_S(w/2) = p_r(w) + f_S(w/2)$, and hence $p_r(w) = 0$. But $0 \leq r \leq x \leq w$ implies $r = p_r(r) = 0$, and hence $x - f_S(w/2) \leq 0$. Therefore $x = |x - f_S(w/2)| = f_S(w/2) - x$, and we have that $f_S(w/2) \in T$, from which it follows that $w \leq f_S(w/2)$. The latter tells us that

$$\begin{aligned} |w/2 - f_S(w/2)| &= f_S(w/2) - w/2 \geq w - w/2 \\ &= |w/2 - w| = |w/2 - 0|, \end{aligned}$$

from which we may conclude $0 = f_S(w/2) = w$. The result follows.

With the lemma above we can proceed to prove the convexity of an l -nps S as follows: Suppose $a, b \in S$ and $a \leq b$. Then x in $[a, b]$ implies that $f_S(x) \leq x \vee f_S(x) \leq b \vee f_S(x) \in S$. Define $T = S - f_S(x)$; then T is an l -nps, $t = b \vee f_S(x) - f_S(x) \in T$, and $x \vee f_S(x) - f_S(x) \in [0, t]$ such that $f_T(x \vee f_S(x) - f_S(x)) = 0$. By Lemma 8, $x \vee f_S(x) - f_S(x) = 0$, or $f_S(x) \geq x$. Since $-S$ is an l -nps, with $-b \leq -x \leq -a$, the same argument produces $f_{-S}(-x) \geq -x$; i.e., $f_S(x) \leq x$. Thus we have proven the convexity of S .

LEMMA 9. If S is an l -nps and $a, b \in S$ such that $a \leq b$, then $[a, b] \subseteq S$.

We can now demonstrate the converse to Theorem 1.

THEOREM 4. Let G be a complete vector lattice. S is an l -nps iff $S = x + [-\alpha, \beta]$, for some $x \in G$ and $\alpha, \beta \in I$.

PROOF. Suppose S is an l -nps and $x \in S$. Then $T = S - x$ is an l -nps and $0 \in T$. Let $\beta(x) = \sup\{x \wedge y | y \in T\}$, and $\alpha(x) = \sup\{x \wedge y | y \in -T\}$, for $x \in P$. It is clear that $\alpha, \beta \in I$, by Lemma 5, and since $x \wedge y \in [0, y]$, for $y \in T \cap P$, then $x \wedge y \in T \cap P$, by Lemma 9, and we have $\beta(x) = \sup\{y | y \in T \text{ and } y \leq x\}$, which is in T by Lemma 2. Similarly for α . Thus $[0, \beta] \subseteq T \cap P$ and $[-\alpha, 0] \subseteq T \cap (-P)$. Since the reverse containment is clear, we have

$$[-\alpha, \beta] = [-\alpha, 0] + [0, \beta] = T \cap (-P) + T \cap P = T.$$

The converse implication follows from Theorem 1.

BIBLIOGRAPHY

1. W. D. L. Appling, *Set functions and an extremal problem in absolute continuity*, Rend. Circ. Mat. Palermo (2) **14** (1965), 256–266.
2. ———, *A generalization of absolute continuity and of an analogue of the Lebesgue decomposition theorem*, Riv. Mat. Univ. Parma (3) **2** (1973), 251–276.
3. G. Birkhoff, *Lattice theory*, Amer. Math. Soc. Colloq. Publ., vol. 25, Amer. Math. Soc., Providence, R. I., 1967.
4. S. Bochner and R. S. Phillips, *Additive set functions and vector lattices*, Ann. of Math. (2) **42** (1941), 316–324.
5. S. Kakutani, *Concrete representation of abstract (L)-spaces and the mean ergodic theorem*, Ann. of Math. (2) **42** (1941), 523–537.
6. L. V. Kantorovič, B. Z. Vulih and A. G. Pinsker, *Partially ordered groups and partially ordered linear spaces*, Amer. Math. Soc. Transl. (2) **27** (1963), 51–124.
7. A. G. Pinsker, *On the extension of partially ordered spaces*, Dokl. Akad. Nauk SSSR **21** (1938), 6–10. (Russian)
8. ———, *On some properties of extended K-spaces*, Dokl. Akad. Nauk SSSR **22** (1939), 220–224. (Russian)

DEPARTMENT OF MATHEMATICS, ARKANSAS TECH UNIVERSITY, RUSSELLVILLE, ARKANSAS 72801