CONVERGENCE OF PROBABILITY MEASURES ON SEPARABLE BANACH SPACES

L. Š. GRINBLAT

ABSTRACT. The following result follows immediately from a general theorem on the convergence of probability measures on separable Banach spaces: On the space $C[0, 1]$ there exists a norm $p(x)$ equivalent to the ordinary norm such that if $\xi_1(t), \dots, \xi_n(t), \dots$ and $\xi(t)$ are continuous random processes ($0 < t < 1$) and for any finite set of points $t_1, \dots, t_k \subset [0, 1]$ the joint distribution of $p(\xi_n), \xi_n(t_1), \dots, \xi_n(t_k)$ converges to the joint distribution of $p(\xi), \xi(t_1), \dots, \xi(t_k)$, then $\xi_n(t)$ converges weakly to $\xi(t)$.

1. Let X be a Banach space and X^* be the dual space of X .

DEFINITION [1, §2]. We say that X has the H -property relative to a set $\Gamma \subset X^*$ if from the conditions $\|x_n\| = \|x\| = 1$, $\lim_{n \rightarrow \infty} f(x_n) = f(x)$ for all $f \in \Gamma$ it follows that $\lim_{n \rightarrow \infty} \|x_n - x\| = 0$.

In what follows we replace the phrase, " X has the property H relative to Γ " by the symbol $X \in H_\Gamma$.

THEOREM 1 (KADETS, SEE [1, §2]). *Let X be a separable Banach space, and Γ be a subset of X^* such that*

$$\inf_{x \in X} \sup_{f \in \Gamma} \frac{|f(x)|}{\|f\| \cdot \|x\|} = \eta(\Gamma) > 0 \quad (x \neq 0, f \neq 0).$$

Then we can introduce in X an equivalent norm $h(x)$ relative to which $X \in H_\Gamma$.

It is easy to see that if $\eta(\Gamma) > 0$, then there exists a countable set $\Gamma_0 \subset \Gamma$ such that $\eta(\Gamma_0) > 0$.

In what follows in this section, we suppose that $X, \Gamma, h(x)$ satisfy the conditions of Theorem 1 and Γ is a countable set.

Let

$$S = \{x \in X: h(x) = 1\}, \quad B = \{x \in X: h(x) \leq 1\}.$$

We denote by $L_\Gamma(S)$ the Banach algebra¹ of bounded continuous functionals,² defined on S , which is generated by all functionals from Γ , restricted to S .

Received by the editors September 9, 1976 and, in revised form, March 23, 1977.

AMS (MOS) subject classifications (1970). Primary 60B05, 60B10; Secondary 46B99, 54D35.

¹Throughout this paper, an algebra will mean an algebra with identity.

²Throughout this paper, a functional defined on a set A will mean a real function (without supplementary conditions) on A .

We denote by $M_\Gamma(B)$ the Banach algebra of bounded continuous functionals, defined on B , which is generated by all functionals from Γ , restricted to B , and the functional $h(x)$, restricted to B .

LEMMA 1. *For any point $x_0 \in S$ and for any closed set $F \subset S$ ($x_0 \notin F$) there exists $f \in L_\Gamma(S)$ such that $f(x_0) = 0$ and $f(x) \geq 1$ if $x \in F$.*

PROOF. Let $\Gamma = \{f_1, \dots, f_k, \dots\}$. Consider numbers $\varepsilon_1 > \varepsilon_2 > \dots > \varepsilon_k \dots$, such that $\varepsilon_k \rightarrow 0$. Let $U_k = \{x \in S: |f_i(x_0) - f_i(x)| < \varepsilon_k, i = 1, \dots, k\}$. There exists k such that $U_k \cap F = \emptyset$. Define the functional f by

$$f(x) = \frac{1}{\varepsilon_k^2} \sum_{i=1}^k (f_i(x_0) - f_i(x))^2. \quad \text{Q.E.D.}$$

LEMMA 2. *For any point $x_0 \in B$ and for any closed set $F \subset B$ ($x_0 \notin F$) there exists $f \in M_\Gamma(B)$ such that $f(x_0) = 0$ and $f(x) \geq 1$ if $x \in F$.*

PROOF. Let $h(x_0) = a$. There exists a positive number $\varepsilon < a$ such that

$$U = \{x \in B: |h(x) - a| < \varepsilon, h(xa/h(x) - x_0) < \varepsilon\}$$

is contained in $B \setminus F$. Let

$$\hat{F} = \{x \in B \setminus U: h(x) = a\}.$$

By Lemma 1 applied to $\hat{F}$ there exists $\hat{f} \in M_\Gamma(B)$ such that $\hat{f}(x_0) = 0$ and $\hat{f} \geq 1$ on $\hat{F}$. Since functionals in $M_\Gamma(B)$ are uniformly continuous, there exists a positive number $\delta < \min(\varepsilon, \frac{1}{2})$ such that $\hat{f}(x) > \frac{1}{2}$ if $|h(x) - a| < \delta$ and $xa/h(x) \in B \setminus U$. If $x \in F$, either $|h(x) - a| > \delta$, or $|h(x) - a| < \delta$ and $xa/h(x) \in B \setminus U$. Therefore we may take f to be

$$f(x) = \frac{1}{\delta^2} [\hat{f}(x)^2 + (h(x) - a)^2]. \quad \text{Q.E.D.}$$

In [2, §5] and in [3, §2] the following was proved:

LEMMA 3. *Let the probability measures $\mu_1, \dots, \mu_n, \dots$ and μ be defined on the Borel sets of the separable metric space Ω and let $G(\Omega)$ be a Banach algebra of bounded continuous functions defined on Ω . Suppose that for any point $\omega_0 \in \Omega$ and for any closed set $F \subset \Omega$ ($\omega_0 \notin F$) there exists $g \in G(\Omega)$ such that $g(\omega_0) = 0$ and $g(\omega) \geq 1$ for all $\omega \in F$. Suppose, finally, that for every function $g \in G(\Omega)$ the distribution of $g(\omega)$ with respect to the measure μ_n converges to the distribution of $g(\omega)$ with respect to the measure μ . Then μ_n converges weakly to μ .*

Lemmas 2 and 3 imply the following:

THEOREM 2. *Let the probability measures $\mu_1, \dots, \mu_n, \dots$ and μ be defined on the Borel sets of X . If for any finite set of functionals $f_1, \dots, f_k \in \Gamma$ the joint distribution of $h(x), f_1(x), \dots, f_k(x)$ with respect to the measure μ_n converges to the joint distribution $h(x), f_1(x), \dots, f_k(x)$ with respect to the measure μ , then μ_n converges weakly to μ .*

2. Consider the space $C[0, 1]$ with the ordinary norm $\|x(t)\| = \max_{0 \leq t \leq 1} |x(t)|$. Let Γ be the set of functionals $f(x) = x(\tau)$, where τ is rational, $0 \leq \tau \leq 1$, $x \in C[0, 1]$. It is obvious that $\eta(\Gamma) = 1$. By virtue of Theorem 1 on the space $C[0, 1]$ there exists a norm $p(x)$, equivalent to the ordinary norm, relative to which $C[0, 1] \in H_\Gamma$. Theorem 2 implies the following:

THEOREM 3. *Let $\xi_1(t), \dots, \xi_n(t), \dots$, and $\xi(t)$ be continuous random processes. If for any finite set of points $t_1, \dots, t_k \subset [0, 1]$ the joint distribution of $p(\xi_n), \xi_n(t_1), \dots, \xi_n(t_k)$ converges to the joint distribution of $p(\xi), \xi(t_1), \dots, \xi(t_k)$, then for any continuous functional f on $C[0, 1]$ the distribution of $f(\xi_n)$ converges to the distribution of $f(\xi)$.*

One can give an explicit formula for the norm $p(x)$. For example

$$p(x) = \max_{0 \leq t \leq 1} |x(t)| + \sum_{n=1}^{\infty} \frac{1}{2^n} \cdot \max_{|t-s| \leq 1/n} |x(t) - x(s)|.$$

3. It is obvious that if we weaken the hypothesis of Theorem 3, and demand only that the joint distribution of $\xi_n(t_1), \dots, \xi_n(t_k)$ converges to the joint distribution of $\xi(t_1), \dots, \xi(t_k)$, then, in general, the conclusion no longer holds. Therefore, it is interesting to compare Theorem 3 to the following results:

THEOREM 4 [4]. *Let $\xi_1(t), \dots, \xi_n(t), \dots$, and $\xi(t)$ be measurable random processes ($0 \leq t \leq 1$). Suppose that there exist C and $p \geq 1$ such that for all n and t we have $E|\xi_n(t)|^p \leq C$. Suppose, finally, that for any finite set of points $t_1, \dots, t_k \subset [0, 1]$ the joint distribution of $\xi_n(t_1), \dots, \xi_n(t_k)$ converges to the joint distribution of $\xi(t_1), \dots, \xi(t_k)$ and $E|\xi_n(t)|^p \rightarrow E|\xi(t)|^p$ for all $t \in [0, 1]$. Then for any continuous functional f on $L_p[0, 1]$ the distribution of $f(\xi_n)$ converges to the distribution of $f(\xi)$.*

REFERENCES

1. V. D. Mil'man, *Geometric theory of Banach spaces*. II, Russian Math. Surveys **26** (1971), no. 6, 79–163.
2. L. S. Grinblat, *Compactifications of spaces and functions and integration of functionals*, Trans. Amer. Math. Soc. **217** (1976), 195–2.
3. ———, *Convergence of random processes without discontinuities of the second kind and limit theorems for sums of independent random variables*, Trans. Amer. Math. Soc. (to appear).
4. ———, *A limit theorem for measurable random processes and its applications*, Proc. Amer. Math. Soc. **61** (1976), 371–376.

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN, ISRAEL

Current address: Department of Mathematics, Harvard University, Cambridge, Massachusetts 02138