HYPERGEOMETRIC FUNCTIONS OF 2×2 MATRIX ARGUMENT ARE EXPRESSIBLE IN TERMS OF APPELL'S FUNCTIONS F_4

TOM KOORNWINDER AND IDA SPRINKHUIZEN-KUYPER

ABSTRACT. It is proved that the hypergeometric function of 2×2 matrix argument is expressible as a solution of the partial differential equations for Appell's function F_4 . As a result the first-mentioned function can be written as a sum of two F_4 -functions.

1. Introduction. Hypergeometric functions of matrix argument, introduced by Herz [6] are of considerable interest in multivariate statistics (cf., for instance, Constantine [2] and James [7]). Until now the hypergeometric functions of matrix argument on the one hand and of Appell and Lauricella type on the other hand seemed to be completely different generalizations of the one-variable case. In the present note it will be shown that at least in the two variable case some relationship exists between these generalized hypergeometric functions. It will be proved that both the hypergeometric function ${}_2F_1(a, b; c; W)$ of 2×2 matrix argument W and Appell's hypergeometric function in two variables $F_4(a, b; c - \frac{1}{2}, 1 + a + b - c; x, y)$ are solutions of the same set of partial differential equations. Since a fundamental system of solutions around $(0, 0)$ of the partial differential equations for F_4 is known, there follows an explicit expression of ${}_2F_1(a, b; c; W)$ as a linear combination of two F_4 functions. For literature about F_4 functions see Erdélyi [4, §5.7].

The idea that the results presented here would be valid, originated from our research on a class of rothogonal polynomials in two variables (cf. Koornwinder and Sprinkhuizen [9]). In a forthcoming paper we will extend the results presented here to the solutions of the partial differential equations of F_4 with arbitrary parameters, thus continuing the work initiated by Erdélyi [3].

2. Preliminaries. Constantine [2] gives a series expansion for ${}_pF_q(W)$ in terms of so-called zonal polynomials which were introduced by James [7]. If W is a 2×2 matrix then the zonal polynomials are explicitly known (cf. James [8, (7.9)]). This results in the explicit expansion

Received by the editors July 21, 1976 and, in revised form, August 24, 1977.

AMS (MOS) subject classifications (1970). Primary 33A30.

Key words and phrases. Hypergeometric functions of matrix argument, Appell's hypergeometric functions F_4 .

$$\begin{aligned}
{}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right) \\
= \sum_{m=0}^{\infty} \sum_{l=0}^m \frac{(a)_m (a - \frac{1}{2})_l (b)_m (b - \frac{1}{2})_l (\frac{3}{2})_{m-l}}{(c)_m (c - \frac{1}{2})_l (\frac{3}{2})_m l! (\frac{1}{2})_{m-l}} \\
\cdot (XY)^{(m+l)/2} P_{m-l}\left(\frac{1}{2} \frac{X+Y}{\sqrt{XY}}\right),
\end{aligned} \tag{2.1}$$

$a, b, c \in \mathbb{C}$, $c \neq \frac{1}{2}, 0, -\frac{1}{2}, -1, \dots$, $X, Y \in \mathbb{C}$, $|X|, |Y| < 1$, where $P_n(x)$ denotes the Legendre polynomial of degree n (cf. Erdélyi [5]). Note that

$${}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & 0 \end{bmatrix}\right) = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m}{(c)_m m!} X^m = {}_2F_1(a, b; c; X).$$

For

$$Z(X, Y) = {}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right),$$

Muirhead [10] gives the following set of partial differential equations:

$$\begin{aligned}
X(1-X)Z_{XX} + \left\{c - \frac{1}{2} - \left(a + b + \frac{1}{2}\right)X + \frac{1}{2} \frac{X(1-X)}{X-Y}\right\} Z_X \\
- \frac{1}{2} \frac{Y(1-Y)}{X-Y} Z_Y = abZ,
\end{aligned} \tag{2.2a}$$

$$\begin{aligned}
Y(1-Y)Z_{YY} + \frac{1}{2} \frac{X(1-X)}{X-Y} Z_X \\
+ \left\{c - \frac{1}{2} - \left(a + b + \frac{1}{2}\right)Y - \frac{1}{2} \frac{Y(1-Y)}{X-Y}\right\} Z_Y = abZ,
\end{aligned} \tag{2.2b}$$

and he proves that ${}_2F_1(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix})$ is the unique solution of each of these differential equations that satisfies the additional properties

- (a) $Z(X, Y)$ is a symmetrical function of X and Y , and
- (b) Z is regular at $(0, 0)$ and $Z(0, 0) = 1$.

The hypergeometric function $F_4(\alpha, \beta; \gamma, \gamma'; x, y)$ of two variables is defined by

$$F_4(\alpha, \beta; \gamma, \gamma'; x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_{m+n}}{(\gamma)_m (\gamma')_n m! n!} x^m y^n, \tag{2.3}$$

$\alpha, \beta, \gamma, \gamma' \in \mathbb{C}$, $\gamma, \gamma' \neq 0, -1, -2, \dots$, $x, y \in \mathbb{C}$, $|x|^{1/2} + |y|^{1/2} < 1$, cf. Appell and Kampé de Fériet [1] or Slater [11]. Note that

$$F_4(\alpha, \beta; \gamma, \gamma'; 0, y) = \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma')_n n!} y^n = {}_2F_1(\alpha, \beta; \gamma'; y).$$

The function $F_4(\alpha, \beta; \gamma, \gamma'; x, y)$ satisfies the following set of partial differential equations:

$$x(1-x)z_{xx} - 2xyz_{xy} - y^2z_{yy} + (\gamma - (\alpha + \beta + 1)x)z_x - (\alpha + \beta + 1)yz_y = \alpha\beta z, \quad (2.4a)$$

$$-x^2z_{xx} - 2xyz_{xy} + y(1-y)z_{yy} - (\alpha + \beta + 1)xz_x + (\gamma' - (\alpha + \beta + 1)y)z_y = \alpha\beta z. \quad (2.4b)$$

In any simply connected region included in $\{(x, y) \in \mathbb{C}^2 | x \neq 0, y \neq 0, |x|^{1/2} + |y|^{1/2} < 1\}$, the general solution of (2.4(a),(b)) reads

$$\begin{aligned} z(x, y) = & AF_4(\alpha, \beta; \gamma, \gamma'; x, y) \\ & + By^{1-\gamma'}F_4(1 + \alpha - \gamma', 1 + \beta - \gamma'; \gamma, 2 - \gamma'; x, y) \\ & + Cx^{1-\gamma}F_4(1 + \alpha - \gamma, 1 + \beta - \gamma; 2 - \gamma, \gamma'; x, y) \\ & + Dx^{1-\gamma}y^{1-\gamma'}F_4(2 + \alpha - \gamma - \gamma', 2 + \beta - \gamma - \gamma'; \\ & \quad 2 - \gamma, 2 - \gamma'; x, y), \end{aligned} \quad (2.5)$$

provided that γ and γ' are noninteger (cf. Appell and Kampé de Fériet [1]).

3. The main result. Consider the transformation of variables

$$x = XY, \quad y = (1 - X)(1 - Y). \quad (3.1)$$

THEOREM 3.1. *The function given by (2.1) is an analytic function of x and y in some neighbourhood of $(0, 1)$ and it satisfies the system of partial differential equations (2.4(a),(b)) with $\alpha := a$, $\beta := b$, $\gamma := c - \frac{1}{2}$, $\gamma' := 1 + a + b - c$.*

PROOF. The first statement is clear from (2.1). For proving the second statement let S and T denote the equations (2.2a) and (2.2b), respectively. Then (2.4a) and (2.4b) are obtained by transforming the equations

$$\frac{1}{2}(S + T) - \frac{2 - X - Y}{2(X - Y)}(S - T), \quad \frac{1}{2}(S + T) + \frac{X - Y}{2(X - Y)}(S - T)$$

in terms of x and y . $\square$

THEOREM 3.2.

$$\begin{aligned} {}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right) &= \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)} \\ &\times F_4\left(a, b; c - \frac{1}{2}, 1 + a + b - c; XY, (1 - X)(1 - Y)\right) \\ &+ \frac{\Gamma(c)\Gamma(a + b - c)}{\Gamma(a)\Gamma(b)} \{(1 - X)(1 - Y)\}^{c - a - b} \\ &\times F_4\left(c - b, c - a; c - \frac{1}{2}, 1 + c - a - b; XY, (1 - X)(1 - Y)\right), \end{aligned}$$

with X and Y such that all terms are well defined and provided that c is not an integer or half integer $\leq \frac{1}{2}$ and that $a + b - c$ is not an integer.

PROOF. Let $z(x, y)$ denote the function given by (2.1) considered as a function of x and y in a neighbourhood of $(0, 1)$. Then $z(x, y)$ is analytic in a neighbourhood of some point $(0, y)$, $0 < y < 1$. In view of (2.5) it follows that

$$z = AF_4\left(a, b; c - \frac{1}{2}, 1 + a + b - c; x, y\right) \\ + By^{c-a-b}F_4\left(c - b, c - a; c - \frac{1}{2}, 1 + c - a - b; x, y\right).$$

Restriction to $x = 0$ gives

$${}_2F_1(a, b; c; 1 - y) = A {}_2F_1(a, b; 1 + a + b - c; y) \\ + By^{c-a-b} {}_2F_1(c - b, c - a; 1 + c - a - b; y).$$

It follows from Erdélyi [4, §2.10] that

$$A = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}, \quad B = \frac{\Gamma(c)\Gamma(a + b - c)}{\Gamma(a)\Gamma(b)}. \quad \square$$

ACKNOWLEDGEMENT. We thank the referee for suggesting some improvements.

REFERENCES

1. P. Appell and J. Kampé de Fériet, *Fonctions hypergéométriques et hypersphériques, Polynômes d'Hermite*, Gauthier-Villars, Paris, 1926.
2. A. G. Constantine, *Some non-central distribution problems in multivariate analysis*, Ann. Math. Statist. **34** (1963), 1270–1285.
3. A. Erdélyi, *Integration of the differential equations of Appell's function F_4* , Quart. J. Math. Oxford Ser. **11** (1941), 68–77.
4. A. Erdélyi, W. Magnus, F. Obberhettinger and F. G. Tricomi, *Higher transcendental functions*, Vol. 1, McGraw-Hill, New York, 1953.
5. ———, *Higher transcendental functions*, Vol. 2, McGraw-Hill, New York, 1953.
6. C. S. Herz, *Bessel functions of matrix argument*, Ann. of Math. (2) **61** (1955), 474–523.
7. A. T. James, *Distribution of matrix variates and latent roots derived from normal samples*, Ann. Math. Statist. **35** (1964), 475–501.
8. ———, *Calculation of zonal polynomial coefficients by use of the Laplace-Beltrami operator*, Ann. Math. Statist. **39** (1968), 1711–1718.
9. T. H. Koornwinder and I. G. Sprinkhuizen-Kuyper, *Generalized power series expansions for a class of orthogonal polynomials in two variables*, SIAM J. Math. Anal. (to appear).
10. R. J. Muirhead, *Systems of partial differential equations for hypergeometric functions of matrix argument*, Ann. Math. Statist. **41** (1970), 991–1001.
11. L. J. Slater, *Generalized hypergeometric functions*, Cambridge Univ. Press, Cambridge, 1966.

MATHEMATISCH CENTRUM, 2^e BOERHAAVESTRAAT 49, AMSTERDAM, NETHERLANDS