AN IDENTITY ON ALGEBRAS OVER A HOPF ALGEBRA

STAVROS PAPASTAVRIDIS

ABSTRACT. Let A be a connected Hopf algebra which has an associative comultiplication $\psi: A \rightarrow A \otimes A$. Let $\chi: A \rightarrow A$ be the canonical conjugation on A . Let M be a graded algebra over the Hopf algebra A . If $x, y \in M$, $\psi(a) = \sum a' \otimes a''$, then we have the identity

$$ax \cdot y = \sum (-1)^{\deg x \cdot \deg a''} a'(x \cdot \chi(a''))y.$$

If X is a topological space and $x, y \in H^*(X; Z_2)$, then we have the identity

$$\text{Sq}^n x \cdot y = \sum_{i=0}^n \text{Sq}^i(x \cdot \chi(\text{Sq}^{n-i})y)$$

(for the proof see [1]). This formula has been proved useful in quite a few occasions (for example see [1], [2]).

In this note we give a generalization of this formula for algebras over a Hopf algebra.

Let A be a Hopf algebra over a commutative ring with unit. We assume that A is connected and that its comultiplication ψ is associative. Let $\chi: A \rightarrow A$ be the canonical conjugation of A . Let M be a graded algebra over the Hopf algebra A . For the terminology and the basic results, we refer to [3]. Under those assumptions, we have

THEOREM. Let $x, y \in M$ and $\psi(a) = \sum_i a'_i \otimes a''_i$. Then we have

$$ax \cdot y = \sum_i (-1)^{\deg x \cdot \deg a''_i} a'_i(x \cdot \chi(a''_i))y.$$

PROOF. Let

$$\psi(a'_i) = \sum_j b'_{ij} \otimes b''_{ij} \quad \text{and} \quad \psi(a''_i) = \sum_j c'_{ij} \otimes c''_{ij}.$$

Since ψ is associative, we will have

$$\sum_{i,j} b'_{ij} \otimes b''_{ij} \otimes a''_i = \sum_{i,j} a'_i \otimes c'_{ij} \otimes c''_{ij}.$$

From the basic properties of tensor products and the fact that χ is linear, we get that

$$\sum_{i,j} b'_{ij} \otimes b''_{ij} \cdot \chi(a''_i) = \sum_{i,j} a'_i \otimes c'_{ij} \cdot \chi(c''_{ij}).$$

Received by the editors June 28, 1977.

AMS (MOS) subject classifications (1970). Primary 57F05.

Key words and phrases. Hopf algebras.

© American Mathematical Society 1978

So we get (see [1, p. 171])

$$\begin{aligned} & \sum_{i,j} (-1)^{\deg x \cdot \deg b''_{ij}} b'_{ij} x \cdot (-1)^{\deg x \cdot \deg a''_{ij}} b''_{ij} \chi(a''_i) y \\ &= \sum_i (-1)^{\deg x \cdot \deg a''_i} a'_i x \cdot \left(\sum_j c'_{ij} \chi(c''_{ij}) \right) y. \end{aligned}$$

But $\sum_j c'_{ij} \chi(c''_{ij}) = 0$ unless $\deg a''_i = 0$, so the second member of the previous identity is $ax \cdot y$. Because of the “Cartan” formula (see [3, p. 173]), the first member of the previous identity is equal to the second member of the identity that we want to prove. That ends the proof.

REFERENCES

1. S. Papastavridis, *A formula for the obstruction to transversality*, *Topology* **11** (1972), 415–416.
2. ———, *The Arf invariant of manifolds with few non-zero Stiefel-Whitney classes*, Ph. D. Thesis, Princeton Univ., Princeton, N. J., 1974.
3. N. Steenrod, *The cohomology algebra of a space*, *Enseignement Math.* **7** (1961), 153–177.

UNIVERSITY OF ATHENS, MATHEMATICAL INSTITUTE, SOLONOS 57, ATHENS 143, GREECE

Current address: Department of Mathematics, University of Crete, Iraklion, Greece