MINIMAL H^2 INTERPOLATION IN THE CARATHÉODORY CLASS

E. BELLER AND B. PINCHUK

ABSTRACT. For $(c_1, \dots, c_n)$ in $\mathbb{C}^n$, let $C(c_1, \dots, c_n)$ denote the class of functions $f(z) = 1 + c_1z + \dots + c_nz^n + \sum_{k=n+1}^{\infty} a_kz^k$ which are analytic and satisfy $\operatorname{Re} f(z) > 0$ in the unit disc. The unique function of least H^2 norm in $C(c_1, \dots, c_n)$ is explicitly determined.

In this paper we consider a minimal interpolation problem at the origin, for the class $H^2 \cap C$, where H^2 denotes the well-known Hardy space for the unit disc, and C is the Carathéodory class of functions $f(z) = 1 + c_1z + c_2z^2 + \dots$ which are analytic and satisfy $\operatorname{Re} f(z) > 0$ in $|z| < 1$. In particular, given n complex numbers $c_1, \dots, c_n$, we wish to find the function f in $H^2 \cap C$, of the form

$$f(z) = 1 + c_1z + \dots + c_nz^n + \sum_{k=n+1}^{\infty} a_kz^k \quad (1)$$

with least H^2 norm.

For each n , expansion (1) defines a mapping

$$\nu_n: f \rightarrow (c_1, \dots, c_n)$$

of C onto some compact set $C_n \subset \mathbb{C}^n$, which is called the n th coefficient body for C . The following result is due to C. Carathéodory and O. Toeplitz (see [2]):

THEOREM A. C_n is a convex, compact body in $\mathbb{C}^n$. To each point in the interior of C_n there correspond infinitely many functions in C ; but each boundary point of C_n corresponds to only one f in C . The boundary points correspond to functions of the form

$$f(z) = \sum_{k=1}^m \frac{1 + \alpha_k z}{1 - \alpha_k z} \mu_k,$$

where $1 \leq m \leq n$; $|\alpha_k| = 1$, $\mu_k > 0$ ($k = 1, \dots, m$), and $\sum_1^m \mu_k = 1$.

From Theorem A it follows that a function corresponding to a boundary point of C_n has infinite H^2 norm. On the other hand, any interior point corresponds to infinitely many functions in $H^2 \cap C$. Indeed, if $(c_1, \dots, c_n)$ is interior to C_n , choose $0 < r < 1$ such that $(b_1, \dots, b_n) \in C_n$, where $b_k =$

Received by the editors November 21, 1977 and, in revised form, January 17, 1978.

AMS (MOS) subject classifications (1970). Primary 30A38, 30A76; Secondary 42A04, 49B35.

Key words and phrases. Positive real part, H^2 , minimal interpolation, Riesz-Herglotz representation, coefficient body, multiplier rule.

© American Mathematical Society 1978

c_k/r^k ($k = 1, \dots, n$). Let $g(z)$ be any function in C which corresponds to $(b_1, \dots, b_n)$. Then $f(z) \equiv g(rz)$ is still in C , is now in H^2 , and corresponds to $(c_1, \dots, c_n)$.

Thus, our interpolation problem is meaningful precisely for the interior of C_n . If we let $C(c_1, \dots, c_n)$ denote the set of all f in C of the form (1), we have the following

THEOREM. *For each $(c_1, \dots, c_n)$ in the interior of C_n , there exists a unique function f with least H^2 norm in $C(c_1, \dots, c_n)$. This f is of the form*

$$f(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{e^{it} + z}{e^{it} - z} u(t) dt, \quad (2)$$

where $u(t) = \max(0, P(t))$, $P(t) = a_0 + \sum_{k=1}^n (a_k \cos kt + b_k \sin kt)$, (a_k, b_k real).

PROOF. $C(c_1, \dots, c_n)$ is a normal and compact family, and $\|f\|_2$ is a continuous functional. Thus, we are assured of the existence of an extremal f .

It is classical that functions f in C have the Riesz-Herglotz representation

$$f(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{e^{it} + z}{e^{it} - z} d\mu(t), \quad (3)$$

where $d\mu(t)$ is a positive measure on $[-\pi, \pi]$ for which $(1/2\pi) \int_{-\pi}^{\pi} d\mu(t) = 1$. If we write

$$\hat{\mu}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-ikt} d\mu(t) \quad (k = 0, \pm 1, \pm 2, \dots),$$

then by comparing Taylor coefficients of both sides of (3), we obtain $c_k = 2\hat{\mu}(k)$ ($k = 1, 2, \dots$). We thus have a one-one correspondence between $C(c_1, \dots, c_n)$ and all positive measures $d\mu(t)$ which satisfy

$$\hat{\mu}(0) = 1, \quad \hat{\mu}(k) = \frac{1}{2} c_k \quad (k = 1, \dots, n).$$

Since

$$\|f\|_2 = \left(1 + 4 \sum_1^{\infty} |\hat{\mu}(k)|^2 \right)^{1/2},$$

and the L^2 norm of $d\mu(t)$ is $(1 + 2\sum_1^{\infty} |\hat{\mu}(k)|^2)^{1/2}$, it follows that $f \in H^2 \cap C$ corresponds to $u \in L^2[-\pi, \pi]$, $u(t) \geq 0$, where $u(t)dt = d\mu(t)$.

Thus, we must find $u_0 \in L^2[-\pi, \pi]$, $u_0 \geq 0$, which minimizes $\int_{-\pi}^{\pi} F_0(t, u(t)) dt$ under the constraints

$$\int_{-\pi}^{\pi} F_j(t, u(t)) dt = d_j \quad (j = 1, 2, \dots, 2^{n+1}), \quad (4)$$

where

$$\begin{aligned} F_0(u, t) &= u^2, \\ F_{2k+1}(t, u) &= u \cos kt \quad (k = 0, 1, \dots, n), \\ F_{2k}(t, u) &= u \sin kt \quad (k = 1, \dots, n), \\ d_1 &= 2\pi; \quad d_{2k+1} + id_{2k} = \pi c_k \quad (k = 1, \dots, n). \end{aligned}$$

Of course, the existence of an extremal f implies that a corresponding extremal u_0 exists. We now need the following

LEMMA. Suppose u_0 is the extremal function described above. Then there exist real multipliers $\lambda_0 = 1, \lambda_1, \dots, \lambda_{2n+1}$, for which the function

$$F = \sum_{k=0}^{2n+1} \lambda_k F_k$$

is such that the inequality

$$F(t, u) \geq F(t, u_0(t)) \quad (5)$$

holds for all pairs $(t, u) \in E \times [0, \infty)$, where $E \subset [-\pi, \pi]$ contains almost all points of $[-\pi, \pi]$.

Assume for a moment that this Lemma has been established. Then by elementary calculus, the only function $u_0(t) \geq 0$ which minimizes F in the sense of (5) is

$$u_0(t) = \max \left(0, -\frac{1}{2} \lambda_1 - \frac{1}{2} \sum_{k=1}^n \lambda_{2k} \sin kt + \lambda_{2k+1} \cos kt \right) \quad \text{a.e.} \quad (6)$$

For convenience, let $u_0(t)$ denote the continuous representative of the L^2 equivalence class (6). Then we have the strict inequality $F(t, u) > F(t, u_0(t))$ for all $(t, u) \in [-\pi, \pi] \times [0, \infty)$, $u \neq u_0(t)$. If we write

$$J_k(u) = \int_{-\pi}^{\pi} F_k(t, u(t)) dt - d_k \quad (k = 0, 1, \dots, 2n+1; d_0 = 0),$$

$$J = \sum_{k=0}^{2n+1} \lambda_k J_k,$$

then for any $u(t) \geq 0$ satisfying (4), we have $J(u) = J(u_0)$. Thus, if such a $u(t)$ differs from $u_0(t)$ on a set of positive measure, we have $J_0(u) = J(u) > J(u_0) = J_0(u_0)$. In other words, u_0 is the *unique* minimizing function. This in turn implies that the corresponding extremal f , given by (2), is unique. Thus, our Theorem is proven, subject to the Lemma.

PROOF OF LEMMA. This Lemma is essentially an L^p version of Theorem 5.1 in [1, p. 215]. That variational theorem deals with piecewise continuous functions, but in our case, we do not know *a priori* that the extremal u_0 is piecewise continuous. In the sequel, we follow the notation of Hestenes [1, Chapters 4, 5].

Let

$$G_i(t) = \int_{-\pi}^t F_i(s, u_0(s)) ds \quad (i = 0, 1, \dots, 2n+1),$$

where u_0 is a nonnegative L^2 function which minimizes $\int_{-\pi}^{\pi} F_0(t, u(t)) dt$ under the constraints (4). Let E be the set of all $t \in (-\pi, \pi)$ such that $G_i'(t) = F_i(t, u_0(t))$ ($i = 0, 1, \dots, 2n+1$). Note that $[-\pi, \pi] - E$ has measure zero.

Let K be the class of all vectors $k = (k^0, k^1, \dots, k^{2n+1})$ of the form

$$k^i = F_i(t, u) - F_i(t, u_0(t)) \quad (i = 0, 1, \dots, 2n+1), \quad (7)$$

where $t \in E$, $u \geq 0$. For any finite set of vectors $k_1, \dots, k_M$ in K , we define $u(\epsilon) = u(\epsilon, t)$ as in [1, p. 221]. At the point $\epsilon = 0$, we have

$$\partial J_i(u(\epsilon)) / \partial \epsilon_j = k_j^i \quad (i = 0, 1, \dots, 2n+1; j = 1, \dots, M).$$

Let K^* denote the cone generated by K , and K^- the cone generated by the vector $(-1, 0, \dots, 0)$. $K^+ = K^* - K^-$ denotes the set of all vectors of the form $k - k^-$, where $k \in K^*$, $k^- \in K^-$. Set $N = 2n+2$. Then Lemmas 8.1 and 8.2 follow as in [1, pp. 192–194]. What remains to be proven is that there exists a vector $k \neq 0$ which is not in the cone K^+ .

Suppose that K^+ contained all vectors. Select N vectors $k_1^*, \dots, k_N^*$ in K^* having the properties described in Lemma 8.2. Since K^* is generated by K , there exists a finite set of vectors $k_1, \dots, k_M$ in K such that

$$k_j^* = \sum_{i=1}^M k_i a_{ij}, \quad a_{ij} \geq 0 \quad (j = 1, \dots, N).$$

Writing $\beta = (\beta_1, \dots, \beta_N)$, ($\beta_j \geq 0$), we define ϵ_i by

$$\epsilon_i = \sum_{j=1}^N a_{ij} \beta_j \quad (i = 1, \dots, M),$$

or in matrix notation, $\epsilon = A\beta$.

Writing $u_A(\beta) = u(A\beta) = u(\epsilon)$, where $u(\epsilon)$ is as above, we have, at $\beta = 0$,

$$\frac{\partial J_{i-1}(u_A(\beta))}{\partial \beta_j} = k_j^{*i} \quad (i, j = 1, \dots, N).$$

If we now define $F_i(z, t)$ as in Lemma 8.3 [1, p. 195], (with k_j^* instead of k_j), then the $F_i(z, t)$ will be continuous, and we may proceed as in the proof of Lemma 8.3. Thus, we conclude that there exist multipliers $\lambda_0 \geq 0$, $\lambda_1, \dots, \lambda_{2n+1}$ such that $\sum_0^{2n+1} \lambda_i k^i \geq 0$ for all $k \in K$. In view of (7), this is exactly inequality (5).

In our specific case, $\lambda_0 \neq 0$. Indeed, suppose $\lambda_0 = 0$. Then we would have $F(t, u) = uT(t)$, where T is a trigonometric polynomial. If $T(t)$ were negative on some interval, then F could not have a minimum on that interval. If $T(t)$ were nonnegative, then the only $u_0(t) \geq 0$ which minimized F would be $u_0(t) \equiv 0$, but this u_0 does not satisfy the first side condition

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} u_0(t) dt = 1.$$

Thus, $\lambda_0 > 0$, and we may take $\lambda_0 = 1$.

REFERENCES

1. M. R. Hestenes, *Calculus of variations and optimal control theory*, Wiley, New York, 1966.
2. A. Pfluger, *Some coefficient problems of starlike functions*, Ann. Acad. Sci. Fenn. Ser. AI 2 (1976), 383–396.

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT GAN, ISRAEL