POSITIVELY CURVED COMPLEX SUBMANIFOLDS IMMERSED IN A COMPLEX SPACE FORM

TAKEHIRO ITOH

Dedicated to Professor T. Otsuki on his 60th birthday

ABSTRACT. The author gives the partial solution for the conjecture; a Kaehler submanifold in a complex space form of constant holomorphic sectional curvature 1 is totally geodesic if ever its holomorphic sectional curvature is greater than $\frac{1}{2}$.

Let $\tilde{M}_m(\tilde{c})$ be an m -dimensional complex space form of constant holomorphic sectional curvature $\tilde{c}$. Let M_n be an n -dimensional Kaehler submanifold of $\tilde{M}_m(\tilde{c})$. Then we have had the following conjecture:

If all holomorphic sectional curvatures of M_n are greater than $\tilde{c}/2$, then M_n is totally geodesic.

In [3], K. Ogiue gave some partial results for this conjecture.

In this paper, we shall prove the following partial solution for the above conjecture,

THEOREM. *Let M be an n -dimensional complete Kaehler submanifold immersed in an m -dimensional complex space form of constant holomorphic sectional curvature $\tilde{c}$. If every holomorphic sectional curvature of M (with respect to the induced metric) is greater than $3n\tilde{c}/(4n+2)$, then M is totally geodesic.*

1. Preliminaries. Let $\tilde{M}$ be an $(n+p)$ -dimensional complex space form of constant holomorphic sectional curvature $\tilde{c}$ and M be an n -dimensional Kaehler submanifold immersed in $\tilde{M}$ (i.e., complex submanifold with the induced Kaehler structure). Let J (resp. $\tilde{J}$) be the complex structure of M (resp. $\tilde{M}$) and g (resp. $\tilde{g}$) be the Kaehler metric of M (resp. $\tilde{M}$). We denote by ∇ (resp. $\tilde{\nabla}$) the covariant differentiation with respect to g (resp. $\tilde{g}$). Then the second fundamental form σ of the immersion is given by

$$\sigma(X, Y) = \tilde{\nabla}_X Y - \nabla_X Y \quad \text{for vector fields } X \text{ and } Y \text{ on } M, \quad (1.1)$$

and it satisfies

$$\sigma(X, Y) = \sigma(Y, X), \quad \sigma(JX, Y) = \sigma(X, JY) = \tilde{J}(\sigma(X, Y)). \quad (1.2)$$

We choose a local field of orthonormal frames $e_1, e_2, \dots, e_n, e_{1*} = Je_1, \dots, e_{n*} = Je_n, e_{\bar{1}}, e_{\bar{2}}, \dots, e_{\bar{p}}, e_{\bar{1}*} = \tilde{J}e_{\bar{1}}, \dots, e_{\bar{p}*} = \tilde{J}e_{\bar{p}}$ in $\tilde{M}$ in such a way that, restricted to M , $e_1, \dots, e_n$ are tangent to M . With respect to the

Received by the editors November 28, 1977.

AMS (MOS) subject classifications (1970). Primary 53C55, 53C40.

Key words and phrases. Kaehler submanifold, Holomorphic pinching.

© American Mathematical Society 1978

frame fields chosen above, let $\omega_1, \dots, \omega_n, \omega_{1^*}, \dots, \omega_{n^*}, \omega_{\bar{1}}, \dots, \omega_{\bar{p}}, \omega_{\bar{1}^*}, \dots, \omega_{\bar{p}^*}$ be the field of dual frames. Then, restricting these forms to M , we have the structure equations of M :¹

$$\begin{cases} \omega_\lambda = 0, & \omega_{i\lambda} = \sum h_{ij}^\lambda \omega_j, & h_{ij}^\lambda = h_{ji}^\lambda, \\ d\omega_i = \sum \omega_{ij} \wedge \omega_j, & \omega_{ij} + \omega_{ji} = 0, \\ d\omega_{ij} = \sum \omega_{ik} \wedge \omega_{kj} - \Omega_{ij}, & \Omega_{ij} = \frac{1}{2} \sum R_{ijkl} \omega_k \wedge \omega_l. \end{cases} \quad (1.3)$$

The equation of Gauss is written as

$$R_{ijkl} = \frac{\tilde{c}}{4} (\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk} + J_{ik} J_{jl} - J_{il} J_{jk} + 2J_{ij} J_{kl}) + \sum (h_{ik}^\lambda h_{jl}^\lambda - h_{il}^\lambda h_{jk}^\lambda), \quad (1.4)$$

where

$$J = (J_{ij}) = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix},$$

I_n being the identity matrix. Let $H_\alpha = (h_{ij}^\alpha)$, $H'_\alpha = (h_{ab}^\alpha)$ and $H''_\alpha = (h_{ab}^{\alpha*})$, then we can easily see that

$$H_\alpha = \begin{pmatrix} H'_\alpha & H''_\alpha \\ H''_\alpha & -H'_\alpha \end{pmatrix} \quad \text{and} \quad H_{\alpha^*} = \begin{pmatrix} -H''_\alpha & H'_\alpha \\ H'_\alpha & H''_\alpha \end{pmatrix}. \quad (1.5)$$

Now, we can consider the following nonnegative functions on M ;

$$K_N = \sum \left(\sum_k (h_{ik}^\lambda h_{kj}^\mu - h_{jk}^\lambda h_{ki}^\mu) \right)^2 \quad \text{and} \quad L_N = \sum h_{ij}^\lambda h_{ij}^\mu h_{kl}^\lambda h_{kl}^\mu.$$

Then we know the following differential equation [1]:

$$\frac{1}{2} \Delta \|\sigma\|^2 = \sum \|\nabla' \sigma\|^2 - K_N - L_N + \frac{n+2}{2} \tilde{c} \|\sigma\|^2, \quad (1.6)$$

where $\|\phi\|$ is the length of a form ϕ and ∇' is the covariant differentiation respect to the connection in (tangent bundle of M) $\oplus$ (normal bundle) induced naturally from $\tilde{\nabla}$.

2. Proof of Theorem. First of all we note that, by the theorem of Myers [3], M is compact. Let c be the minimum of all holomorphic sectional curvatures of M . The holomorphic sectional curvature $H(X)$ of M determined by a unit vector X is given by

$$H(X) = \tilde{c} - 2\|\sigma(X, X)\|^2. \quad (2.1)$$

¹We use the following convention on the range of indices unless otherwise stated: $a, b, c = 1, 2, \dots, n$; $i, j, k, l = 1, 2, \dots, n, 1^*, \dots, n^*$; $\alpha, \beta, \gamma = 1, \dots, \bar{p}$; $\lambda, \mu, \nu = 1, 2, \dots, \bar{p}, 1^*, 2^*, \dots, \bar{p}^*$; and we agree that repeated indices under a summation sign without indication are summed over the respective range.

It is clear from (2.1) that $\tilde{c} \geq c$. Setting $A_{ij} := \sigma(e_i, e_j)$, from (1.2) we have

$$A_{a^*b^*} = -A_{ab}, \quad A_{ab^*} = A_{a^*b} = \tilde{J}A_{ab}, \quad \text{for } a, b. \quad (2.2)$$

For a vector $X = \lambda e_i + \eta e_j$ on M , we get

$$\sigma(X, X) = \lambda^2 A_{ii} + 2\lambda\eta A_{ij} + \eta^2 A_{jj}.$$

Since $H(X) \geq c$ for all unit vector X , $\|\sigma(X, X)\| \leq (\tilde{c} - c)/2$, in particular,

$$\|A_{aa}\|^2 = \|\sigma(e_a, e_a)\|^2 \leq (\tilde{c} - c)/2 \quad \text{for } \forall a. \quad (2.3)$$

Since $\|\sigma(X, X)\| \leq (\tilde{c} - c)/2$ for all unit vectors $X = \lambda e_i + \eta e_j$ ($i \neq j$), given by

i	j	λ	η
a	b	λ	η
		λ	$-\eta$
a	b^*	λ	η
		λ	$-\eta$

i	j	λ	η
b	a	λ	η
		λ	$-\eta$
b	a^*	λ	η
		λ	$-\eta$

we have

$$(\lambda^4 + \eta^4)(\|A_{aa}\|^2 + \|A_{bb}\|^2) + 8\lambda^2\eta^2\|A_{ab}\|^2 \leq \tilde{c} - c. \quad (2.4)$$

Since $\lambda^4 + \eta^4 \geq 2\lambda^2\eta^2$, from (2.4) we get

$$\lambda^2\eta^2(\|A_{aa}\|^2 + \|A_{bb}\|^2 + 4\|A_{ab}\|^2) \leq (\tilde{c} - c)/2$$

in which, setting $\lambda^2 = \eta^2 = \frac{1}{2}$, we get

$$\|A_{aa}\|^2 + \|A_{bb}\|^2 + 4\|A_{ab}\|^2 \leq 2(\tilde{c} - c). \quad (2.5)$$

Since $\|A_{aa}\|^2 + \|A_{bb}\|^2 \geq 2\|A_{aa}\| \cdot \|A_{bb}\|$, (2.5) implies

$$\|A_{aa}\| \cdot \|A_{bb}\| + 2\|A_{ab}\|^2 \leq \tilde{c} - c. \quad (2.6)$$

Using (2.3), (2.6) and the fact that $|\langle A_{aa}, A_{bb} \rangle| \leq \|A_{aa}\| \cdot \|A_{bb}\|$, we can prove

LEMMA. *The following inequality holds on M ;*

$$K_N + L_N \leq (2n + 1)(\tilde{c} - c)\|\sigma\|^2. \quad (2.7)$$

PROOF OF LEMMA. For each α , we set

$$K_N^\alpha := \sum_{i,j,\mu} \left(\sum_k (h_{ik}^\alpha h_{kj}^\mu - h_{jk}^\alpha h_{ki}^\mu) \right)^2 + \sum_{i,j,\mu} \left(\sum_k (h_{ik}^{\alpha^*} h_{kj}^\mu - h_{jk}^{\alpha^*} h_{ki}^\mu) \right)^2,$$

$$L_N^\alpha := \sum_{i,j,k,l,\mu} h_{ij}^\alpha h_{ij}^\mu h_{kl}^\alpha h_{kl}^\mu + \sum_{i,j,k,l,\mu} h_{ij}^{\alpha^*} h_{ij}^\mu h_{kl}^{\alpha^*} h_{kl}^\mu.$$

Then $K_N = \sum_\alpha K_N^\alpha$ and $L_N = \sum_\alpha L_N^\alpha$. For fixed α , choosing a frame $\{e_1, \dots, e_n, e_{1^*}, \dots, e_{n^*}\}$ such that $h_{ij}^\alpha = \lambda_i^\alpha \delta_{ij}$, from (1.5) we have

$$H_\alpha = \begin{pmatrix} H'_\alpha & 0 \\ 0 & -H'_\alpha \end{pmatrix}, \quad H_{\alpha^*} = \begin{pmatrix} 0 & H'_\alpha \\ H'_\alpha & 0 \end{pmatrix} \quad \text{and} \quad H'_\alpha = \begin{pmatrix} \lambda_1^\alpha & & 0 \\ & \ddots & \\ & 0 & \lambda_n^\alpha \end{pmatrix}. \quad (2.8)$$

Using (2.8), we have

$$K_N^\alpha = 8 \sum_{a \neq b} \{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \|A_{ab}\|^2 + 16 \sum_a (\lambda_a^\alpha)^2 \|A_{aa}\|^2,$$

$$L_N^\alpha = 8 \sum_{a \neq b} \lambda_a^\alpha \lambda_b^\alpha \langle A_{aa}, A_{bb} \rangle + 8 \sum_a (\lambda_a^\alpha)^2 \|A_{aa}\|^2,$$

and so

$$\begin{aligned} K_N^\alpha + L_N^\alpha &= 8 \sum_{a \neq b} \left[\{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \|A_{ab}\|^2 + \lambda_a^\alpha \lambda_b^\alpha \langle A_{aa}, A_{bb} \rangle \right] \\ &\quad + 24 \sum_a (\lambda_a^\alpha)^2 \|A_{aa}\|^2. \end{aligned} \quad (2.9)$$

Since we have the following inequalities

$$2\lambda_a^\alpha \lambda_b^\alpha \langle A_{aa}, A_{bb} \rangle \leq -\{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \langle A_{aa}, A_{bb} \rangle \quad \text{if } \langle A_{aa}, A_{bb} \rangle < 0,$$

$$2\lambda_a^\alpha \lambda_b^\alpha \langle A_{aa}, A_{bb} \rangle \leq \{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \langle A_{aa}, A_{bb} \rangle \quad \text{if } \langle A_{aa}, A_{bb} \rangle \geq 0,$$

and $|\langle A_{aa}, A_{bb} \rangle| \leq \|A_{aa}\| \cdot \|A_{bb}\|$, we get

$$2\lambda_a^\alpha \lambda_b^\alpha \langle A_{aa}, A_{bb} \rangle \leq \{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \|A_{aa}\| \cdot \|A_{bb}\|. \quad (2.10)$$

It follows from (2.9) and (2.10) that we have

$$\begin{aligned} K_N^\alpha + L_N^\alpha &\leq 4 \sum_{a \neq b} \{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} \{2\|A_{ab}\|^2 + \|A_{aa}\| \cdot \|A_{bb}\|\} \\ &\quad + 24 \sum_a (\lambda_a^\alpha)^2 \|A_{aa}\|^2. \end{aligned} \quad (2.11)$$

Hence, using (2.3) and (2.6), from (2.11) we obtain

$$\begin{aligned} K_N^\alpha + L_N^\alpha &\leq 4(\tilde{c} - c) \left[\sum_{a \neq b} \{(\lambda_a^\alpha)^2 + (\lambda_b^\alpha)^2\} + 3 \sum_a (\lambda_a^\alpha)^2 \right] \\ &= 4(2n + 1)(\tilde{c} - c) \sum_a (\lambda_a^\alpha)^2 \\ &= (2n + 1)(\tilde{c} - c) (\text{Tr } H_\alpha^2 + \text{Tr } H_{\alpha^*}^2). \end{aligned} \quad (2.12)$$

Since $K_N + L_N = \sum_\alpha (K_N^\alpha + L_N^\alpha)$ and $\|\sigma\|^2 = \sum_\alpha (\text{Tr } H_\alpha^2 + \text{Tr } H_{\alpha^*}^2)$, (2.12) implies (2.7). Q.E.D.

PROOF OF THEOREM. Using (2.7), from (1.6) we have

$$\begin{aligned} \frac{1}{2} \Delta \|\sigma\|^2 &\geq \|\nabla' \sigma\|^2 + \frac{n+2}{2} \tilde{c} \|\sigma\|^2 - (2n+1)(\tilde{c} - c) \|\sigma\|^2 \\ &\geq (2n+1) \left(c - \frac{3n\tilde{c}}{4n+2} \right) \|\sigma\|^2 \quad \text{on } M. \end{aligned} \quad (2.13)$$

Since M is compact and $c > 3n\tilde{c}/(4n + 2)$, from (2.13) $\Delta\|\sigma\|^2 = 0$ on M . Hence we have $\|\sigma\| = 0$ identically on M , so that M is totally geodesic. Q.E.D.

BIBLIOGRAPHY

1. S. S. Chern, M. do Carmo and S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, Functional Analysis and Related Fields, Springer, New York, 1970, pp. 59–75.
2. T. Itoh, *Addendum to my paper "On Veronese manifolds"*, J. Math. Soc. Japan **30** (1978), 73–74.
3. K. Ogiue, *Differential geometry of Kaehler submanifolds*, Advances in Math. **13** (1974), 73–114.
4. S. Myer, *Riemannian manifolds with positive mean curvature*, Duke Math. J. **8** (1941), 401–404.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TSUKUBA, IBARAKI 300-31, JAPAN